

## AN OPTIMAL INEQUALITIES CHAIN FOR BIVARIATE MEANS

ZHEN-HANG YANG AND YU-MING CHU

**Abstract.** Let  $p \in \mathbb{R}$ ,  $M$  be a bivariate mean, and  $M_p$  be defined by  $M_p(a, b) = M^{1/p}(a^p, b^p)$  ( $p \neq 0$ ) and  $M_0(a, b) = \lim_{p \rightarrow 0} M_p(a, b)$ . In this paper, we prove that the sharp inequalities  $L_2(a, b) < P(a, b) < NS_{1/2}(a, b) < He(a, b) < A_{2/3}(a, b) < I(a, b) < Z_{1/3}(a, b) < Y_{1/2}(a, b)$  hold for all  $a, b > 0$  with  $a \neq b$ , where  $L(a, b) = (a - b)/(\log a - \log b)$ ,  $P(a, b) = (a - b)/[2\arcsin((a - b)/(a + b))]$ ,  $NS(a, b) = (a - b)/[2\operatorname{arcsinh}((a - b)/(a + b))]$ ,  $He(a, b) = (a + \sqrt{ab} + b)/3$ ,  $A(a, b) = (a + b)/2$ ,  $I(a, b) = 1/e(a^a/b^b)^{1/(a-b)}$ ,  $Z(a, b) = a^{a/(a+b)}b^{b/(a+b)}$  and  $Y(a, b) = I(a, b)e^{1-ab/L^2(a, b)}$  are respectively the logarithmic, first Seiffert, Neuman-Sándor, Heronian, arithmetic, identric, power-exponential and exponential-geometric means of  $a$  and  $b$ .

*Mathematics subject classification (2010):* 26E60.

*Keywords and phrases:* Logarithmic mean, first Seiffert mean, Neuman-Sándor mean, identric mean, power-exponential mean, exponential-geometric mean.

## REFERENCES

- [1] D. ACU, *Some inequalities for certain means in two arguments*, Gen. Math. **9**, 1-2 (2001), 11–14.
- [2] H. ALZER AND W. JANOUS, *Solution of problem 8\**, Crux Math. **13** (1987), 173–178.
- [3] P. S. BULLEN, D. S. MITRINović AND P. M. VASIĆ, *Means and Their Inequalities*, D. Reidel, Dordrecht, 1988.
- [4] Y.-M. CHU, M.-K. WANG AND Y.-F. QIU, *An optimal double inequality between power-type Heron and Seiffert means*, J. Inequal. Appl. **2010**, Article ID 146945, 11 pages.
- [5] Y.-M. CHU AND B.-Y. LONG, *Bounds of the Neuman-Sándor mean using power and identric means*, Abstr. Appl. Anal. **2013**, Article ID 832591, 6 pages.
- [6] Y.-M. CHU, B.-Y. LONG, W.-M. GONG AND Y.-Q. SONG, *Sharp bounds for Seiffert and Neuman-Sándor means in terms of generalized logarithmic means*, J. Inequal. Appl. **2013**, 10 (2013), 13 pages.
- [7] I. COSTIN AND G. TOADER, *Optimal evaluations of some Seiffert-type means by power means*, Appl. Math. Comput. **219**, 9 (2013), 4745–4754.
- [8] I. COSTIN AND G. TOADER, *A nice separation of some Seiffert-type means by power means*, Int. J. Math. Math. Sci. **2012**, Article ID 430692, 6 pages.
- [9] C. GINI, *Di una formula comprensiva delle media*, Metron **13**, 2 (1938), 3–22.
- [10] P. A. HÄSTÖ, *A monotonicity property of ratios of symmetric homogeneous means*, JIPAM. J. Inequal. Pure Appl. Math. **3**, 5 (2002), Article 71, 23 pages.
- [11] P. A. HÄSTÖ, *Optimal inequalities between Seiffert's mean and power mean*, Math. Inequal. Appl. **7**, 1 (2004), 47–53.
- [12] A. A. JAGERS, *Solution of problem 887*, Nieuw Arch. Wisk. (4) **12** (1994), 230–231.
- [13] G. JIA AND J.-D. CAO, *A new upper bound of the logarithmic mean*, JIPAM. J. Inequal. Pure Appl. Math. **4**, 4 (2003), Article 80, 4 pages.
- [14] T. P. LIN, *The power mean and the logarithmic mean*, Amer. Math. Monthly **81** (1974), 879–883.
- [15] E. NEUMAN AND J. SÁNDOR, *On the Schwab-Borchardt mean II*, Math. Pannon. **17**, 1 (2006), 49–59.
- [16] Z. PÁLES, *Inequalities for sums of powers*, J. Math. Anal. Appl. **131**, 1 (1988), 265–270.
- [17] A. O. PITTINGER, *Inequalities between arithmetic and logarithmic means*, Univ. Beograd Publ. Elektrotehn. Fak. Ser. Mat. Fiz. **678-715** (1980), 15–18.

- [18] K. B. STOLARSKY, *The power and generalized logarithmic means*, Amer. Math. Monthly **87**, 7 (1980), 545–548.
- [19] J. SÁNDOR, *On certain inequalities for means III*, Arch. Math. **76**, 1 (2001), 34–40.
- [20] J. SÁNDOR, *A note on the Gini means*, Gen. Math. **12**, 4 (2004), 17–21.
- [21] H.-J. SEIFFERT, *Werte zwischen dem geometrischen und dem arithmetischen Mittel zweier Zahlen*, Elem. Math. **42** (1987), 105–107.
- [22] H.-J. SEIFFERT, *Aufgabe β 16*, Die Wurzel **29** (1995), 221–222.
- [23] S. TOADER, *Derivatives of generalized means*, Math. Inequal. Appl. **5** (2002), 517–532.
- [24] ZH.-H. YANG, *On the homogeneous functions with two parameters and its monotonicity*, JIPAM. J. Inequal. Pure Appl. Math. **6**, 4 (2005), Article 101, 11 pages.
- [25] ZH.-H. YANG, *On the log-convexity of two-parameter homogeneous functions*, Math. Inequal. Appl. **10**, 3 (2007), 499–516.
- [26] ZH.-H. YANG, *On the monotonicity and log-convexity of a four-parameter homogeneous mean*, J. Inequal. Appl. **2008**, Article ID 149286, 12 pages.

- [27] ZH.-H. YANG, *Sharp bounds for the second Seiffert mean in terms of power means*, arXiv: 1206.5494v1 [math. CA].
- [28] ZH.-H. YANG, *Sharp power means bounds for Neuman-Sandor mean*, arXiv: 1208.0895v1 [math. CA].
- [29] ZH.-H. YANG, *Estimates for Neuman-Sandor mean by power means and their relative errors*, J. Math. Inequal. **7**, 4 (2013), 711–726.