

QUANTUM INTEGRAL INEQUALITIES FOR CONVEX FUNCTIONS

WEERAWAT SUDSUTAD, SOTIRIS K. NTOUYAS AND JESSADA TARIBOON

(Communicated by A. Aglić Aljinović)

Abstract. In this paper we establish some new quantum integral inequalities for convex functions.

1. Introduction

A function $f : I \rightarrow \mathbb{R}$, $\emptyset \neq I \subseteq \mathbb{R}$, is said to be convex on I if inequality

$$f(tx + (1-t)y) \leq t f(x) + (1-t)f(y),$$

holds for all $x, y \in I$ and $t \in [0, 1]$. Many inequalities have been established for convex functions but the most famous is the Hermite-Hadamard's inequality [1], due to its rich geometrical significance and applications, which is stated as follows:

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex mapping and $a, b \in I$ with $a < b$. Then

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}. \quad (1.1)$$

Both the inequalities hold in reversed direction if f is concave. Since its discovery, Hermite-Hadamard's inequality has been considered the most useful inequality in mathematical analysis. This inequality has been extended in a number of ways and a number of papers have been written. The main aim of this paper is to establish some new quantum integral inequalities for convex functions. Many consequences of Hermite-Hadamard type inequalities are obtained as special cases when $q \rightarrow 1$.

Mathematics subject classification (2010): 34A08, 26D10, 26D15.

Keywords and phrases: Hermite-Hadamard inequality, convex functions, integral inequalities.

2. Preliminaries

Let $J := [a, b] \subset \mathbb{R}$, $J^o := (a, b)$ be interval and $0 < q < 1$ be a constant. We define q -derivative of a function $f : J \rightarrow \mathbb{R}$ at a point $x \in J$ on $[a, b]$ as follows.

DEFINITION 2.1. Assume $f : J \rightarrow \mathbb{R}$ is a continuous function and let $x \in J$. Then the expression

$${}_aD_q f(x) = \frac{f(x) - f(qx + (1-q)a)}{(1-q)(x-a)}, \quad x \neq a, \quad {}_aD_q f(a) = \lim_{x \rightarrow a} {}_aD_q f(x), \quad (2.1)$$

is called the q -derivative on J of function f at x .

We say that f is q -differentiable on J provided ${}_aD_q f(x)$ exists for all $x \in J$. Note that if $a = 0$ in (2.1), then ${}_0D_q f = \mathcal{D}_q f$, where \mathcal{D}_q is the well-known q -derivative of the function $f(x)$ defined by

$$\mathcal{D}_q f(x) = \frac{f(x) - f(qx)}{(1-q)x}. \quad (2.2)$$

For more details, see [2].

LEMMA 2.1. [3] Let $\alpha \in \mathbb{R}$, then we have

$${}_aD_q (x-a)^\alpha = \left(\frac{1-q^\alpha}{1-q} \right) (x-a)^{\alpha-1}. \quad (2.3)$$

DEFINITION 2.2. Assume $f : J \rightarrow \mathbb{R}$ is a continuous function. Then the q -integral on J is defined by

$$\int_a^x f(t) {}_aD_q t = (1-q)(x-a) \sum_{n=0}^{\infty} q^n f(q^n x + (1-q^n)a), \quad (2.4)$$

for $x \in J$. Moreover, if $c \in (a, x)$ then the definite q -integral on J is defined by

$$\begin{aligned} \int_c^x f(t) {}_aD_q t &= \int_a^x f(t) {}_aD_q t - \int_a^c f(t) {}_aD_q t \\ &= (1-q)(x-a) \sum_{n=0}^{\infty} q^n f(q^n x + (1-q^n)a) \\ &\quad - (1-q)(c-a) \sum_{n=0}^{\infty} q^n f(q^n c + (1-q^n)a). \end{aligned}$$

Note that if $a = 0$, then (2.4) reduces to the classical q -integral of a function $f(x)$, defined by $\int_0^x f(t) {}_0D_q t = (1-q)x \sum_{n=0}^{\infty} q^n f(q^n x)$ for $x \in [0, \infty)$. For more details, see [2].

THEOREM 2.1. [3] Let $f : J \rightarrow \mathbb{R}$ be a continuous function. Then we have

- (i) ${}_aD_q \int_a^x f(t) {}_a d_q t = f(x)$;
- (ii) $\int_c^x {}_a D_q f(t) {}_a d_q t = f(x) - f(c)$ for $c \in (a, x)$.

THEOREM 2.2. [3] Assume $f, g : J \rightarrow \mathbb{R}$ are continuous functions, $\alpha \in \mathbb{R}$. Then, for $x \in J$,

- (i) $\int_a^x [f(t) + g(t)] {}_a d_q t = \int_a^x f(t) {}_a d_q t + \int_a^x g(t) {}_a d_q t$;
- (ii) $\int_a^x (\alpha f)(t) {}_a d_q t = \alpha \int_a^x f(t) {}_a d_q t$;
- (iii) $\int_c^x f(t) {}_a D_q g(t) {}_a d_q t = (fg)|_c^x - \int_c^x g(qt + (1-q)a) {}_a D_q f(t) {}_a d_q t$, for $c \in (a, x)$.

LEMMA 2.2. [4] For $\alpha \in \mathbb{R} \setminus \{-1\}$, the following formula holds:

$$\int_a^x (t-a)^\alpha {}_a d_q t = \left(\frac{1-q}{1-q^{\alpha+1}} \right) (x-a)^{\alpha+1}. \quad (2.5)$$

THEOREM 2.3. [4] (q -Hermite-Hadamard) Let $f : J \rightarrow \mathbb{R}$ be a convex continuous function on J and $0 < q < 1$. Then we have

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) {}_a d_q t \leq \frac{qf(a) + f(b)}{1+q}. \quad (2.6)$$

3. Auxiliary results

In this section, we present some auxiliary results which are used throughout this article.

LEMMA 3.1. Let $f : J \rightarrow \mathbb{R}$ be a continuous function and $0 < q < 1$. If ${}_a D_q f$ is an integrable function on J^o , then the following equality holds:

$$\begin{aligned} & \frac{1}{b-a} \int_a^b f(s) {}_a d_q s - \frac{qf(a) + f(b)}{1+q} \\ &= \frac{q(b-a)}{1+q} \int_0^1 (1-(1+q)s) {}_a D_q f(sb + (1-s)a) {}_0 d_q s. \end{aligned} \quad (3.1)$$

Proof. Using Definitions 2.1 and 2.2, we have

$$\begin{aligned} & \int_0^1 (1-(1+q)s) {}_a D_q f(sb + (1-s)a) {}_0 d_q s \\ &= \int_0^1 \left(\frac{f(sb + (1-s)a) - f(sqb + (1-sq)a)}{(1-q)(b-a)s} \right) {}_0 d_q s \\ &\quad - (1+q) \int_0^1 s \left(\frac{f(sb + (1-s)a) - f(sqb + (1-sq)a)}{(1-q)(b-a)s} \right) {}_0 d_q s \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{b-a} \left[\sum_{n=0}^{\infty} f(q^n b + (1-q^n)a) - \sum_{n=0}^{\infty} f(q^{n+1} b + (1-q^{n+1})a) \right] \\
&\quad - \frac{(1+q)}{b-a} \left[\sum_{n=0}^{\infty} q^n f(q^n b + (1-q^n)a) - \sum_{n=0}^{\infty} q^n f(q^{n+1} b + (1-q^{n+1})a) \right] \\
&= \frac{f(b)-f(a)}{b-a} - \frac{(1+q)}{b-a} \sum_{n=0}^{\infty} q^n f(q^n b + (1-q^n)a) \\
&\quad + \frac{(1+q)}{q(b-a)} \sum_{n=1}^{\infty} q^n f(q^n b + (1-q^n)a) \\
&= \frac{f(b)-f(a)}{b-a} - \frac{(1+q)}{b-a} \sum_{n=0}^{\infty} q^n f(q^n b + (1-q^n)a) \\
&\quad + \frac{(1+q)}{q(b-a)} \left[f(b) - f(b) + \sum_{n=1}^{\infty} q^n f(q^n b + (1-q^n)a) \right] \\
&= \frac{f(b)-f(a)}{b-a} - \frac{(1+q)}{q(b-a)} f(b) + \frac{(1+q)}{q(b-a)} \sum_{n=0}^{\infty} q^n f(q^n b + (1-q^n)a) \\
&\quad - \frac{(1+q)}{b-a} \sum_{n=0}^{\infty} q^n f(q^n b + (1-q^n)a) \\
&= -\frac{qf(a)+f(b)}{q(b-a)} + \frac{(1+q)(1-q)}{q(b-a)} \frac{(b-a)}{(b-a)} \sum_{n=0}^{\infty} q^n f(q^n b + (1-q^n)a) \\
&= -\frac{qf(a)+f(b)}{q(b-a)} + \frac{(1+q)}{q(b-a)^2} \int_a^b f(s)_a d_q s.
\end{aligned}$$

Therefore, we obtain the desired result in (3.1) as required. The proof is completed. \square

REMARK 3.1. If $q \rightarrow 1$, then (3.1) reduces to

$$\frac{f(a)+f(b)}{2} - \frac{1}{b-a} \int_a^b f(s) ds = \frac{b-a}{2} \int_0^1 (1-2s) f'(sb + (1-s)a) ds.$$

See also [5, Lemma 2.1, page 91].

LEMMA 3.2. Let $0 < q < 1$ be a constant. Then the following equality holds:

$$\int_0^1 s |1 - (1+q)s|_0 d_q s = \frac{q(1+4q+q^2)}{(1+q+q^2)(1+q)^3}. \quad (3.2)$$

Proof. By computing directly, we have

$$\begin{aligned}
&\int_0^1 s |1 - (1+q)s|_0 d_q s \\
&= \int_0^{\frac{1}{1+q}} s (1 - (1+q)s)_0 d_q s + \int_{\frac{1}{1+q}}^1 s ((1+q)s - 1)_0 d_q s
\end{aligned}$$

$$\begin{aligned}
&= \int_0^{\frac{1}{1+q}} s_0 d_q s - (1+q) \int_0^{\frac{1}{1+q}} s^2_0 d_q s + (1+q) \int_{\frac{1}{1+q}}^1 s^2_0 d_q s - \int_{\frac{1}{1+q}}^1 s_0 d_q s \\
&= \frac{1}{(1+q)^3} - \frac{1+q}{(1+q+q^2)(1+q)^3} + \frac{(1+q)(3q+3q^2+q^3)}{(1+q+q^2)(1+q)^3} - \frac{2q+q^2}{(1+q)^3} \\
&= \frac{q(1+4q+q^2)}{(1+q+q^2)(1+q)^3}.
\end{aligned}$$

The proof is completed. \square

LEMMA 3.3. Let $0 < q < 1$ be a constant. Then the following equality holds:

$$\int_0^1 (1-s)|1-(1+q)s|_0 d_q s = \frac{q(1+3q^2+2q^3)}{(1+q+q^2)(1+q)^3}. \quad (3.3)$$

Proof. Taking into account Lemma 3.2, we have

$$\begin{aligned}
&\int_0^1 (1-s)|1-(1+q)s|_0 d_q s \\
&= \int_0^1 |1-(1+q)s|_0 d_q s - \int_0^1 s|1-(1+q)s|_0 d_q s \\
&= \int_0^{\frac{1}{1+q}} (1-(1+q)s)_0 d_q s + \int_{\frac{1}{1+q}}^1 ((1+q)s-1)_0 d_q s - \int_0^1 s|1-(1+q)s|_0 d_q s \\
&= \int_0^{\frac{1}{1+q}} 1_0 d_q s - (1+q) \int_0^{\frac{1}{1+q}} s_0 d_q s + (1+q) \int_{\frac{1}{1+q}}^1 s_0 d_q s - \int_{\frac{1}{1+q}}^1 1_0 d_q s - \int_0^1 s|1-(1+q)s|_0 d_q s \\
&= \frac{1}{1+q} - \frac{1+q}{(1+q)^3} + \frac{(1+q)(2q+q^2)}{(1+q)^3} - \frac{q}{1+q} - \frac{q(1+4q+q^2)}{(1+q+q^2)(1+q)^3} \\
&= \frac{q(1+3q^2+2q^3)}{(1+q+q^2)(1+q)^3}.
\end{aligned}$$

The proof is completed. \square

4. Main results

In this section, we present some q -integral inequalities for convex functions on $[a, b]$.

THEOREM 4.1. Let $f : J \rightarrow \mathbb{R}$ be a continuous function. If $|{}_a D_q f|$ is convex and integrable on J^o , then the following inequality holds:

$$\begin{aligned}
&\left| \frac{qf(a)+f(b)}{1+q} - \frac{1}{b-a} \int_a^b f(s) {}_a d_q s \right| \\
&\leq \frac{q^2(b-a)}{(1+q+q^2)(1+q)^4} \left((1+4q+q^2)|{}_a D_q f(b)| + q(1+3q^2+2q^3)|{}_a D_q f(a)| \right).
\end{aligned} \quad (4.1)$$

Proof. Using Lemma 3.1 and the convexity of ${}_aD_q f$ on J^o , we have

$$\begin{aligned} & \left| \frac{qf(a) + f(b)}{1+q} - \frac{1}{b-a} \int_a^b f(s) {}_a d_q s \right| \\ &= \left| \frac{q(b-a)}{1+q} \int_0^1 (1-(1+q)s) {}_a D_q f(sb + (1-s)a) {}_0 d_q s \right| \\ &\leq \frac{q(b-a)}{1+q} \int_0^1 |1-(1+q)s| [s |{}_a D_q f(b)| + (1-s) |{}_a D_q f(a)|] {}_0 d_q s \\ &= \frac{q(b-a)}{1+q} \left[|{}_a D_q f(b)| \int_0^1 |1-(1+q)s| {}_0 d_q s + |{}_a D_q f(a)| \int_0^1 |1-(1+q)s|(1-s) {}_0 d_q s \right]. \end{aligned}$$

Applying Lemmas 3.2 and 3.3, we have

$$\begin{aligned} & \left| \frac{qf(a) + f(b)}{1+q} - \frac{1}{b-a} \int_a^b f(s) {}_a d_q s \right| \\ &\leq \frac{q(b-a)}{1+q} \left[\left(\frac{q(1+4q+q^2)}{(1+q+q^2)(1+q)^3} \right) |{}_a D_q f(b)| + \left(\frac{q(1+3q^2+2q^3)}{(1+q+q^2)(1+q)^3} |{}_a D_q f(a)| \right) \right] \\ &= \frac{q^2(b-a)}{(1+q+q^2)(1+q)^4} \left[(1+4q+q^2) |{}_a D_q f(b)| + ((1+3q^2+2q^3) |{}_a D_q f(a)|) \right]. \end{aligned}$$

The proof is completed. \square

REMARK 4.1. If $q \rightarrow 1$, then the inequality (4.1) reduces to the integral inequality for convex functions

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(s) ds \right| \leq \frac{(b-a)(|f'(a)| + |f'(b)|)}{8}.$$

See also [5, Theorem 2.2, page 92].

Next, we present the second result of q integral inequality for convex functions on $[a, b]$.

THEOREM 4.2. Let $f : J \rightarrow \mathbb{R}$ be a continuous function. If $|{}_a D_q f|^r$ is convex and integrable on J^o and $r \geq 1$ then the following inequality holds:

$$\begin{aligned} & \left| \frac{qf(a) + f(b)}{1+q} - \frac{1}{b-a} \int_a^b f(s) {}_a d_q s \right| \\ &\leq \frac{q^2(2+q+q^2)(b-a)}{(1+q)^4} \left[\frac{(1+4q+q^2)|{}_a D_q f(b)|^r + (1+3q^2+2q^3)|{}_a D_q f(a)|^r}{(1+q+q^2)(2+q+q^3)} \right]^{1/r}. \end{aligned} \tag{4.2}$$

Proof. From Lemma 3.1, we have

$$\begin{aligned} & \left| \frac{qf(a) + f(b)}{1+q} - \frac{1}{b-a} \int_a^b f(s)_a d_q s \right| \\ & \leqslant \frac{q(b-a)}{1+q} \int_0^1 |1-(1+q)s|_0 d_q s \end{aligned} \quad (4.3)$$

and by the power-mean inequality

$$\begin{aligned} & \int_0^1 |1-(1+q)s|_0 d_q s \leqslant \left(\int_0^1 |1-(1+q)s|_0 d_q s \right)^{1-\frac{1}{r}} \\ & \times \left(\int_0^1 |1-(1+q)s|_0 d_q s \right)^{\frac{1}{r}}. \end{aligned} \quad (4.4)$$

Using convexity of $|_a D_q f|^r$ and applying Lemmas 3.2 and 3.3, it follows that

$$\begin{aligned} & \int_0^1 |1-(1+q)s|_0 d_q s \leqslant \int_0^1 |1-(1+q)s| [s|_a D_q f(b)|^r + (1-s)|_a D_q f(a)|^r]_0 d_q s \\ & \leqslant |_a D_q f(b)|^r \int_0^1 s |1-(1+q)s|_0 d_q s + |_a D_q f(a)|^r \int_0^1 (1-s) |1-(1+q)s|_0 d_q s \\ & = \frac{q(1+4q+q^2)}{(1+q+q^2)(1+q)^3} |_a D_q f(b)|^r + \frac{q(1+3q^2+2q^3)}{(1+q+q^2)(1+q)^3} |_a D_q f(a)|^r \\ & = \frac{q}{(1+q+q^2)(1+q)^3} \left[(1+4q+q^2) |_a D_q f(b)|^r + (1+3q^2+2q^3) |_a D_q f(a)|^r \right]. \end{aligned} \quad (4.5)$$

Applying the fact that $\int_0^1 |1-(1+q)s|_0 d_q s = (q(2+q+q^3))/(1+q)^3$ and substituting (4.4) into (4.5) we get

$$\begin{aligned} & \int_0^1 |1-(1+q)s|_0 d_q s \leqslant \left(\frac{q(2+q+q^3)}{(1+q)^3} \right)^{1-\frac{1}{r}} \\ & \times \left(\frac{q}{(1+q+q^2)(1+q)^3} ((1+4q+q^2) |_a D_q f(b)|^q + (1+3q^2+2q^3) |_a D_q f(a)|^q) \right)^{\frac{1}{r}}, \end{aligned}$$

which yield the desired inequality in (4.2) as requested. This completes the proof. \square

REMARK 4.2. If $q \rightarrow 1$, then the inequality (4.2) reduces to integral inequality for convex function

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(s) ds \right| \leqslant \frac{(b-a)}{4} \left[\frac{|f'(a)|^r + |f'(b)|^r}{2} \right]^{\frac{1}{r}}.$$

See also [6, Theorem 1, page 52].

Next, two more q -integral inequalities for convex functions are established.

THEOREM 4.3. *Let f and g be real-valued, nonnegative and convex functions on J . Then the following inequalities hold:*

$$(i) \quad \frac{1}{b-a} \int_a^b f(x)g(x)_a d_q x \leq \frac{f(a)g(a)}{1+q+q^2} + \frac{q(1+q^2)f(b)g(b) + q^2N(a,b)}{(1+q)(1+q+q^2)}, \quad (4.6)$$

$$(ii) \quad 2f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)g(x)_a d_q x \\ + \frac{2q^2M(a,b) + (1+2q+q^2)N(a,b)}{2(1+q)(1+q+q^2)}. \quad (4.7)$$

where $M(a,b) = f(a)g(a) + f(b)g(b)$ and $N(a,b) = f(a)g(b) + f(b)g(a)$.

Proof. (i) Using the convexity of f and g , for all $s \in [0, 1]$, we have

$$f(sb + (1-s)a) \leq sf(b) + (1-s)f(a), \quad (4.8)$$

$$g(sb + (1-s)a) \leq sg(b) + (1-s)g(a). \quad (4.9)$$

Multiplying (4.8) with (4.9), we get

$$\begin{aligned} & f(sb + (1-s)a)g(sb + (1-s)a) \\ & \leq s^2f(b)g(b) + (1-s)^2f(a)g(a) + s(1-s)[f(a)g(b) + f(b)g(a)]. \end{aligned} \quad (4.10)$$

Taking q -integral for (4.10) with respect to s on $(0, 1)$, we obtain

$$\begin{aligned} & \int_0^1 f(sb + (1-s)a)g(sb + (1-s)a)_0 d_qs \\ & \leq f(b)g(b) \int_0^1 s^2{}_0 d_qs + f(a)g(a) \int_0^1 (1-s)^2{}_0 d_qs \\ & \quad + [f(a)g(b) + f(b)g(a)] \int_0^1 s(1-s)_0 d_qs \\ & = \frac{f(b)g(b)}{1+q+q^2} + \frac{q(1+q^2)f(a)g(a)}{(1+q)(1+q+q^2)} + \frac{q^2(f(a)g(b) + f(b)g(a))}{(1+q)(1+q+q^2)}. \end{aligned}$$

By substituting $sa + (1-s)b = x$, we deduce the desired inequality in (4.6).

(ii) Since f and g are convex functions on $[a, b]$, for $s \in [0, 1]$, we get

$$\begin{aligned} & f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \\ & = f\left(\frac{sb + (1-s)a}{2} + \frac{(1-s)b + sa}{2}\right)g\left(\frac{sb + (1-s)a}{2} + \frac{(1-s)b + sa}{2}\right) \end{aligned}$$

$$\begin{aligned}
&\leq \left(\frac{1}{2}f(sb + (1-s)a) + \frac{1}{2}f((1-s)b + sa) \right) \\
&\quad \times \left(\frac{1}{2}g(sb + (1-s)a) + \frac{1}{2}g((1-s)b + sa) \right) \\
&\leq \frac{1}{4} [f(sb + (1-s)a)g(sb + (1-s)a) + f((1-s)b + sa)g((1-s)b + sa)] \\
&\quad + \frac{1}{4} [(sf(b) + (1-s)f(a))((1-s)g(b) + sg(a)) + ((1-s)f(b) + sf(a)) \\
&\quad \times (sg(b) + (1-s)g(a))] \\
&= \frac{1}{4} (f(sb + (1-s)a)g(sb + (1-s)a) + f((1-s)b + sa)g((1-s)b + sa)) \\
&\quad + \frac{1}{4} [2s(1-s)(f(a)g(a) + f(b)g(b)) + (s^2 + (1-s)^2)(f(a)g(b) + f(b)g(a))].
\end{aligned}$$

Taking q -integral with respect to s on $[0, 1]$, we obtain

$$\begin{aligned}
&f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \\
&\leq \frac{1}{4} \int_0^1 (f(sb + (1-s)a)g(sb + (1-s)a) + f((1-s)b + sa)g((1-s)b + sa))_0 d_qs \\
&\quad + \frac{1}{4} [2(f(a)g(a) + f(b)g(b)) \int_0^1 s(1-s)_0 d_qs + (f(a)g(b) + f(b)g(a)) \\
&\quad \times \int_0^1 (s^2 + (1-s)^2)_0 d_qs] \\
&= \frac{2}{4(b-a)} \int_a^b f(x)g(x)_a d_qx + \frac{1}{4} \left[2(f(a)g(a) + f(b)g(b)) \left(\frac{1}{1+q} - \frac{1}{1+q+q^2} \right) \right. \\
&\quad \left. + (f(a)g(b) + f(b)g(a)) \left(\frac{1}{1+q+q^2} + 1 - \frac{2}{1+q} + \frac{1}{1+q+q^2} \right) \right] \\
&= \frac{1}{2(b-a)} \int_a^b f(x)g(x)_a d_qx + \frac{1}{4} \left[\frac{2q^2(f(a)g(a) + f(b)g(b))}{(1+q)(1+q+q^2)} \right. \\
&\quad \left. + \frac{(1+2q+q^3)(f(a)g(b) + f(b)g(a))}{(1+q)(1+q+q^2)} \right],
\end{aligned}$$

which leads to the estimate (4.7). This complete the proof. \square

REMARK 4.3. If $q \rightarrow 1$, then the inequalities (4.6) and (4.7) reduce to the integral inequalities for convex functions

$$\begin{aligned}
\frac{1}{b-a} \int_a^b f(x)g(x)dx &\leq \frac{1}{3}M(a,b) + \frac{1}{6}N(a,b), \\
2f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) &\leq \frac{1}{b-a} \int_a^b f(x)g(x)dx + \frac{1}{6}M(a,b) + \frac{1}{3}N(a,b),
\end{aligned}$$

respectively. See also [7, Theorem 5.2.1, pages 250-251].

Now, the last two q -integral inequalities for convex functions are established.

THEOREM 4.4. *Let f and g be real-valued, nonnegative and convex functions on J . Then the following inequalities hold:*

$$(i) \quad \frac{(1+q)(1+q+q^2)}{(b-a)^2} \int_a^b \int_a^b \int_0^1 f(sy + (1-s)x)g(sy + (1-s)x)_0 d_q s_a d_q x_a d_q y \\ \leq \frac{1+2q+q^2}{b-a} \int_a^b f(x)g(x)_a d_q x + \frac{2q^2}{(1+q)^2} ((q^2 f(a)g(a) + f(b)g(b)) + qN(a,b)), \quad (4.11)$$

$$(ii) \quad \frac{1+q+q^2}{b-a} \int_a^b \int_0^1 f\left(sy + (1-s)\frac{a+b}{2}\right) g\left(sy + (1-s)\frac{a+b}{2}\right)_0 d_q s_a d_q y \\ \leq \frac{1}{b-a} \int_a^b f(x)g(x)_a d_q x + \frac{q(1+q^2)}{4(1+q)} (M(a,b) + N(a,b)) \\ + \frac{q^2}{2(1+q)^2} (2(qf(a)g(a) + f(b)g(b)) + (1+q)N(a,b)). \quad (4.12)$$

where $M(a,b) = f(a)g(a) + f(b)g(b)$ and $N(a,b) = f(a)g(b) + f(b)g(a)$.

Proof. (i) From the convexity of f and g , for all $s \in [0, 1]$, $x, y \in J$, we have

$$f(sy + (1-s)x) \leq sf(y) + (1-s)f(x), \quad (4.13)$$

$$g(sy + (1-s)x) \leq sg(y) + (1-s)g(x). \quad (4.14)$$

Multiplying (4.13) with (4.14), we get

$$f(sy + (1-s)x)g(sy + (1-s)x) \\ \leq s^2 f(y)g(y) + (1-s)^2 f(x)g(x) + s(1-s)[f(x)g(y) + f(y)g(x)]. \quad (4.15)$$

Taking q -integral for (4.15) with respect to s on $(0, 1)$, one has

$$\int_0^1 f(sy + (1-s)x)g(sy + (1-s)x)_0 d_q s \\ \leq f(y)g(x) \int_0^1 s^2_0 d_q s + f(x)g(x) \int_0^1 (1-s)^2_0 d_q s \\ + [f(x)g(y) + f(y)g(x)] \int_0^1 s(1-s)_0 d_q s \\ = \frac{f(y)g(y)}{1+q+q^2} + \frac{q(1+q^2)f(x)g(x)}{(1+q)(1+q+q^2)} + \frac{q^2(f(x)g(y) + f(y)g(x))}{(1+q)(1+q+q^2)}. \quad (4.16)$$

Next, taking double q -integral to both sides of (4.16) with respect to x, y on (a, b) , we obtain

$$\begin{aligned} & \int_a^b \int_a^b \int_0^1 f(sy + (1-s)x)g(sy + (1-s)x)_0 d_q s_a d_q x_a d_q y \\ & \leq \frac{(b-a)}{1+q+q^2} \int_a^b f(y)g(y)_a d_q y + \frac{q(1+q^2)(b-a)}{(1+q)(1+q+q^2)} \int_a^b f(x)g(x)_a d_q x \\ & \quad + \frac{q^2}{(1+q)(1+q+q^2)} \left(\int_a^b g(y)_a d_q y \int_a^b f(x)_a d_q x + \int_a^b f(y)_a d_q y \int_a^b (g(x))_a d_q x \right). \end{aligned} \quad (4.17)$$

By applying Theorem 2.3 on the right hand side of (4.17), we have

$$\begin{aligned} & \int_a^b \int_a^b \int_0^1 f(sy + (1-s)x)g(sy + (1-s)x)_0 d_q s_a d_q x_a d_q y \\ & \leq \left(\frac{1}{1+q+q^2} + \frac{q(1+q^2)}{(1+q)(1+q+q^2)} \right) (b-a) \int_a^b f(x)g(x)_a d_q x \\ & \quad + \frac{2q^2(b-a)^2}{(1+q)^3(1+q+q^2)} ((q^2 f(a)g(a) + f(b(g(b))) + qN(a,b)). \end{aligned} \quad (4.18)$$

Multiplying both sides of (4.18) by $((1+q)(1+q+q^2))/(b-a)^2$, we deduce the desired inequality in (4.11).

(ii) Since f and g are convex functions on $[a, b]$, for $s \in [0, 1]$, we get

$$f\left(sy + (1-s)\left(\frac{a+b}{2}\right)\right) \leq sf(y) + (1-s)f\left(\frac{a+b}{2}\right), \quad (4.19)$$

$$g\left(sy + (1-s)\left(\frac{a+b}{2}\right)\right) \leq sg(y) + (1-s)g\left(\frac{a+b}{2}\right). \quad (4.20)$$

Multiplying (4.19) with (4.20), we obtain

$$\begin{aligned} & f\left(sy + (1-s)\left(\frac{a+b}{2}\right)\right)g\left(sy + (1-s)\left(\frac{a+b}{2}\right)\right) \\ & \leq s^2 f(y)g(y) + (1-s)^2 f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \\ & \quad + s(1-s) \left[f(y)g\left(\frac{a+b}{2}\right) + f\left(\frac{a+b}{2}\right)g(y) \right]. \end{aligned} \quad (4.21)$$

Taking q -integral for (4.21) with respect to s on $(0, 1)$, it follows that

$$\begin{aligned} & \int_0^1 f\left(sy + (1-s)\left(\frac{a+b}{2}\right)\right)g\left(sy + (1-s)\left(\frac{a+b}{2}\right)\right)_0 d_q s \\ & \leq f(y)g(y) \int_0^1 s^2_0 d_q s + f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \int_0^1 (1-s)^2_0 d_q s \\ & \quad + \left[f(y)g\left(\frac{a+b}{2}\right) + f\left(\frac{a+b}{2}\right)g(y) \right] \int_0^1 s(1-s)_0 d_q s \end{aligned}$$

$$\begin{aligned}
&= \frac{f(y)g(y)}{1+q+q^2} + f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right)\left[1 - \frac{2}{1+q} + \frac{1}{1+q+q^2}\right] \\
&\quad + \left[f(y)g\left(\frac{a+b}{2}\right) + f\left(\frac{a+b}{2}\right)g(y)\right]\left[\frac{1}{1+q} - \frac{1}{1+q+q^2}\right] \\
&= \frac{f(y)g(y)}{1+q+q^2} + \left[\frac{q+q^3}{(1+q)(1+q+q^2)}\right]f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \\
&\quad + \left[\frac{q^2}{(1+q)(1+q+q^2)}\right]\left[f(y)g\left(\frac{a+b}{2}\right) + f\left(\frac{a+b}{2}\right)g(y)\right]
\end{aligned}$$

Applying q -integral with respect to y on $[a, b]$ and using Theorem 2.3 with the convexity of the functions f, g , we observe that

$$\begin{aligned}
&\int_a^b \int_0^1 f\left(sy + (1-s)\left(\frac{a+b}{2}\right)\right)g\left(sy + (1-s)\left(\frac{a+b}{2}\right)\right)_0 d_qs_a d_q y \\
&\leqslant \frac{1}{1+q+q^2} \int_a^b f(y)g(y)_a d_q y + \frac{q+q^3}{(1+q)(1+q+q^2)} \int_a^b f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right)_a d_q y \\
&\quad + \frac{q^2}{(1+q)(1+q+q^2)} \left(g\left(\frac{a+b}{2}\right) \int_a^b f(y)_a d_q y + f\left(\frac{a+b}{2}\right) \int_a^b g(y)_a d_q y\right) \\
&\leqslant \frac{1}{1+q+q^2} \int_a^b f(y)g(y)_a d_q y + \frac{(q+q^3)(b-a)}{4(1+q)(1+q+q^2)} (f(a) + f(b))(g(a) + g(b)) \\
&\quad + \frac{q^2(b-a)}{(1+q)(1+q+q^2)} \left(\left(\frac{g(a) + g(b)}{2}\right) \left(\frac{qf(a) + f(b)}{1+q}\right)\right. \\
&\quad \left. + \left(\frac{f(a) + f(b)}{2}\right) \left(\frac{qg(a) + g(b)}{1+q}\right)\right) \\
&= \frac{1}{1+q+q^2} \int_a^b f(y)g(y)_a d_q y + \frac{q(1+q^2)(b-a)}{4(1+q)(1+q+q^2)} (M(a, b) + N(a, b)) \\
&\quad + \frac{q^2(b-a)}{2(1+q)^2(1+q+q^2)} (2(qf(a)g(a) + f(b)g(b)) + (1+q)N(a, b)). \tag{4.22}
\end{aligned}$$

Multiplying both sides of (4.22) by $((1+q)(1+q+q^2))/(b-a)$, we deduce the desired inequality in (4.12). This complete the proof. \square

REMARK 4.4. If $q \rightarrow 1$, then the inequalities (4.11) and (4.12) reduce to the integral inequalities for convex functions

$$\begin{aligned}
&\frac{3}{2(b-a)^2} \int_a^b \int_a^b \int_0^1 f(sy + (1-s)x)g(sy + (1-s)x)_0 d_qs dx dy \\
&\leqslant \frac{1}{b-a} \int_a^b f(x)g(x)dx + \frac{1}{8}(M(a, b) + N(a, b)),
\end{aligned}$$

$$\begin{aligned} & \frac{3}{b-a} \int_a^b \int_0^1 f\left(sy + (1-s)\frac{a+b}{2}\right) g\left(sy + (1-s)\frac{a+b}{2}\right) ds dy \\ & \leq \frac{1}{b-a} \int_a^b f(x)g(x)dx + \frac{1}{2}(M(a,b) + N(a,b)), \end{aligned}$$

respectively. See also [7, Theorem 5.2.2, pages 252–253].

Acknowledgement. The research of W. Sudsutad and J. Tariboon is supported by King Mongkut's University of Technology North Bangkok, Thailand.

REFERENCES

- [1] J. HADAMARD, *Etude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann*, J. Math. Pures Appl. **58** (1893) 171–215.
- [2] V. KAC, P. CHEUNG, *Quantum Calculus*, Springer, New York, 2002.
- [3] J. TARIBOON, S. K. NTOUYAS, *Quantum calculus on finite intervals and applications to impulsive difference equations*, Adv. Differ. Equ. 2013, **2013**:282.
- [4] J. TARIBOON, S. K. NTOUYAS, *Quantum integral inequalities on finite intervals*, J. Inequal. Appl. 2014, **2014**:121.
- [5] S. S. DRAGOMIR AND R. P. AGAWAL, *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, Appl. Math. Lett. **11** (5) (1998) 91–95.
- [6] C. E. M. PEARCE AND J. PEĆCARIĆ, *Inequalities for differentiable mappings with application to special means and quadrature formula*, Appl. Math. Lett. **13** (2000) 51–55.
- [7] B. G. PACHPATTE, *Analytic inequalities: Recent Advances*, Atlantic Press, Amsterdam-Paris, 2012.

(Received June 23, 2014)

Weerawat Sudsutad
*Nonlinear Dynamic Analysis Research Center
 Department of Mathematics, Faculty of Applied Science
 King Mongkut's University of Technology North Bangkok
 Bangkok, 10800 Thailand
 e-mail: wrw.sst@gmail.com*

Sotiris K. Ntouyas
*Department of Mathematics, University of Ioannina
 451 10 Ioannina, Greece
 and
 Nonlinear Analysis and Applied Mathematics (NAAM) – Research Group
 Department of Mathematics, Faculty of Science
 King Abdulaziz University
 P. O. Box 80203, Jeddah 21589, Saudi Arabia
 e-mail: sntouyas@uoi.gr*

Jessada Tariboon
*Nonlinear Dynamic Analysis Research Center
 Department of Mathematics, Faculty of Applied Science
 King Mongkut's University of Technology North Bangkok
 Bangkok, 10800 Thailand
 e-mail: jessadat@kmutnb.ac.th*