

A REFINEMENT OF THE DIFFERENCE BETWEEN TWO INTEGRAL MEANS IN TERMS OF THE CUMULATIVE VARIATION AND APPLICATIONS

WENJUN LIU, WANGSHU WEN AND JAEKEUN PARK

(Communicated by A. Aglić Aljinović)

Abstract. In this paper, we establish a refinement of the difference between two integral means for functions of bounded variation in terms of the cumulative variation function. The applications for probability density functions are also given.

1. Introduction

In the past few years, many authors have considered various generalizations of some kinds of integral inequalities, which give explicit error bounds for some known and some new quadrature formulae. For example, Dragomir [7] established the following Ostrowski's inequality for mappings of bounded variation:

THEOREM 1.1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a function of bounded variation on $[a, b]$. Then, for all $x \in [a, b]$, we have the following inequality*

$$\left| f(x)(b-a) - \int_a^b f(t)dt \right| \leqslant \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \nabla_a^b(f), \quad (1.1)$$

where $\nabla_a^b(f)$ denotes the total variation of f on $[a, b]$. The constant $\frac{1}{2}$ is the best possible.

Recently, the same author [6] considered the following refinements of the inequalities (1.1) in terms of the cumulative variation function and given some applications for selfadjoint operators on complex Hilbert spaces.

Mathematics subject classification (2010): 26D15, 26A45, 60E15.

Keywords and phrases: Ostrowski's inequality, integral mean, cumulative variation, function of bounded variation, probability density function.

THEOREM 1.2. *Let $f : [a, b] \rightarrow \mathbb{C}$ be a function of bounded variation on $[a, b]$. Then*

$$\begin{aligned} \left| f(x)(b-a) - \int_a^b f(t)dt \right| &\leq \int_a^x \left(\bigvee_t^x (f) \right) dt + \int_x^b \left(\bigvee_t^x (f) \right) dt \\ &\leq (x-a) \bigvee_a^x (f) + (b-x) \bigvee_x^b (f) \\ &\leq \begin{cases} \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b (f), \\ \left[\frac{1}{2} \bigvee_a^b (f) + \frac{1}{2} \left| \bigvee_a^x (f) - \bigvee_x^b (f) \right| \right] (b-a), \end{cases} \end{aligned} \quad (1.2)$$

and

$$\begin{aligned} \left| f(x)(b-a) - \int_a^b f(t)dt \right| &\leq \int_a^x \left(\bigvee_t^x (|f(x)-f|) \right) dt + \int_x^b \left(\bigvee_t^x (|f(x)-f|) \right) dt \\ &\leq (x-a) \bigvee_a^x (|f(x)-f|) + (b-x) \bigvee_x^b (|f(x)-f|) \\ &\leq (x-a) \bigvee_a^x (f) + (b-x) \bigvee_x^b (f) \\ &\leq \begin{cases} \left[\frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b (f), \\ \left[\frac{1}{2} \bigvee_a^b (f) + \frac{1}{2} \left| \bigvee_a^x (f) - \bigvee_x^b (f) \right| \right] (b-a), \end{cases} \end{aligned} \quad (1.3)$$

for any $x \in [a, b]$.

In [2], Barnett et al. compared the difference of two integral means for absolutely continuous mapping whose first derivatives are in $L_\infty[a, b]$ and gave several applications for probability density functions, special means, Jeffrey's divergence and continuous streams. In [11], Hwang and Dragomir compared the difference of two integral means as in the following Theorem 1.3 for the functions with bounded variation. The obtained result is also a generalization of (1.1) and has been applied to probability density functions. For related results, see [1], [3]–[5], [10], [12]–[25], [26] and the references therein.

THEOREM 1.3. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a function of bounded variation on $[a, b]$. Then, for all $a \leq x < y \leq b$, we have the inequality*

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(s) ds - \frac{1}{y-x} \int_x^y f(s) ds \right| \\ & \leq \frac{x-a}{b-a} \bigvee_a^x (f) + \max \left\{ \frac{x-a}{b-a}, \frac{b-y}{b-a} \right\} \bigvee_x^y (f) + \frac{b-y}{b-a} \bigvee_y^b (f). \end{aligned} \quad (1.4)$$

The inequality (1.4) is sharp.

Motivated by the above research, the purpose of this paper is to provide a refinement of the inequality (1.4) for functions of bounded variation in terms of the cumulative variation function and give several applications for probability density functions.

2. The refinement of the difference between two integral means

As in [6, 9], for a function of bounded variation $v : [a,b] \rightarrow \mathbb{C}$, we define the Cumulative Variation Function (CVF) $V : [a,b] \rightarrow [0,\infty]$ by

$$V(t) := \bigvee_a^t (v)$$

the total variation of v on the interval $[a,t]$ with $t \in [a,b]$.

It's known that the CVF is monotonic nondecreasing on $[a,b]$ and is continuous in a point $c \in [a,b]$ if and only if the generating function v is continuing in that point. If v is Lipschitzian with the constant $L > 0$, i.e.,

$$|v(t) - v(s)| \leq L|t-s| \text{ for any } t,s \in [a,b]$$

then V is also Lipschitzian with the same constant.

The following lemma is of interest in itself as well, see also [8, 9].

LEMMA 2.1. *Let $f, u : [a,b] \rightarrow \mathbb{R}$. If f is continuous on $[a,b]$ and u is of bounded variation on $[a,b]$, then*

$$\left| \int_a^b f(t) du(t) \right| \leq \int_a^b |f(t)| d \left(\bigvee_a^t (u) \right) \leq \max_{t \in [a,b]} |f(t)| \bigvee_a^b (u). \quad (2.1)$$

The following result may be stated.

THEOREM 2.1. *Let $f : [a,b] \rightarrow \mathbb{R}$ be a function of bounded variation on $[a,b]$. Then, for all $a \leq x < y \leq b$, we have the inequalities*

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(s) ds - \frac{1}{y-x} \int_x^y f(s) ds \right| \\ & \leq \frac{1}{b-a} \int_a^x \left(\bigvee_s^x (f) \right) ds + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \left[\int_x^{s_0} \left(\bigvee_s^s (f) \right) ds + \int_{s_0}^y \left(\bigvee_s^y (f) \right) ds \right] \\ & \quad + \frac{1}{b-a} \int_y^b \left(\bigvee_s^y (f) \right) ds \end{aligned} \quad (2.2)$$

$$\begin{aligned}
&\leq \frac{x-a}{b-a} \bigvee_a^x (f) + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \left[(s_0-x) \bigvee_x^{s_0} (f) + (y-s_0) \bigvee_{s_0}^y (f) \right] + \frac{b-y}{b-a} \bigvee_y^b (f) \\
&= \frac{x-a}{b-a} \bigvee_a^x (f) + \left[\frac{x-a}{b-a} \bigvee_x^{s_0} (f) + \frac{b-y}{b-a} \bigvee_{s_0}^y (f) \right] + \frac{b-y}{b-a} \bigvee_y^b (f) \\
&\leq \begin{cases} \frac{x-a}{b-a} \bigvee_a^x (f) + \max \left\{ \frac{x-a}{b-a}, \frac{b-y}{b-a} \right\} \bigvee_x^y (f) + \frac{b-y}{b-a} \bigvee_y^b (f), \\ \frac{x-a}{b-a} \bigvee_a^x (f) + \left(1 - \frac{y-x}{b-a} \right) \max \left\{ \bigvee_x^{s_0} (f), \bigvee_{s_0}^y (f) \right\} + \frac{b-y}{b-a} \bigvee_y^b (f), \end{cases}
\end{aligned}$$

where $s_0 = \frac{bx-ay}{(b-a)-(y-x)}$ is the solution of $\frac{s_0-x}{y-x} + \frac{a-s_0}{b-a} = 0$. The inequalities (2.2) are sharp.

Proof. We start with the identity

$$\begin{aligned}
&\frac{1}{b-a} \int_a^b f(s) ds - \frac{1}{y-x} \int_x^y f(s) ds \\
&= \int_a^x \frac{a-s}{b-a} df(s) + \int_x^y \left(\frac{s-x}{y-x} + \frac{a-s}{b-a} \right) df(s) + \int_y^b \frac{b-s}{b-a} df(s),
\end{aligned} \tag{2.3}$$

which holds for all $a \leq x < y \leq b$ and $f : [a, b] \rightarrow \mathbb{R}$ is a function of bounded variation on $[a, b]$ (see [11]). We denote

$$p(s) = \frac{s-x}{y-x} + \frac{a-s}{b-a}, \quad s \in [x, y],$$

then we get that

$$p(x) = \frac{a-x}{b-a} \leq 0, \quad p(y) = \frac{b-y}{b-a} \geq 0, \quad p'(s) = \frac{1}{y-x} - \frac{1}{b-a} \geq 0,$$

for all $a \leq x < y \leq b$, which implies that $p(s)$ is nondecreasing on $[x, y]$.

Taking the modulus in (2.3) and using the property (2.1), we have

$$\begin{aligned}
&\left| \frac{1}{b-a} \int_a^b f(s) ds - \frac{1}{y-x} \int_x^y f(s) ds \right| \\
&\leq \left| \int_a^x \frac{a-s}{b-a} df(s) \right| + \left| \int_x^y \left(\frac{s-x}{y-x} + \frac{a-s}{b-a} \right) df(s) \right| + \left| \int_y^b \frac{b-s}{b-a} df(s) \right| \\
&\leq \int_a^x \left| \frac{a-s}{b-a} \right| d \left(\bigvee_a^s (f) \right) + \int_x^y \left| \frac{s-x}{y-x} + \frac{a-s}{b-a} \right| d \left(\bigvee_a^s (f) \right) + \int_y^b \left| \frac{b-s}{b-a} \right| d \left(\bigvee_a^s (f) \right) \\
&= \int_a^x \frac{s-a}{b-a} d \left(\bigvee_a^s (f) \right) - \int_x^{s_0} \left(\frac{s-x}{y-x} + \frac{a-s}{b-a} \right) d \left(\bigvee_a^s (f) \right) \\
&\quad + \int_{s_0}^y \left(\frac{s-x}{y-x} + \frac{a-s}{b-a} \right) d \left(\bigvee_a^s (f) \right) + \int_y^b \frac{b-s}{b-a} d \left(\bigvee_a^s (f) \right)
\end{aligned} \tag{2.4}$$

for all $a \leq x < y \leq b$.

Utilising the integration by parts for Riemann-Stieltjes integral and the fact that $\frac{s_0 - x}{y - x} + \frac{a - s_0}{b - a} = 0$, we get

$$\begin{aligned} \int_a^x \frac{s-a}{b-a} d\left(\bigvee_a^s(f)\right) &= \left(\frac{s-a}{b-a}\right) \bigvee_a^s(f) \Big|_{s=a}^x - \frac{1}{b-a} \int_a^x \left(\bigvee_a^s(f)\right) ds \quad (2.5) \\ &= \frac{x-a}{b-a} \bigvee_a^x(f) - \frac{1}{b-a} \int_a^x \left(\bigvee_a^s(f)\right) ds \\ &= \frac{1}{b-a} \int_a^x \left(\bigvee_a^x(f)\right) ds - \frac{1}{b-a} \int_a^x \left(\bigvee_a^s(f)\right) ds \\ &= \frac{1}{b-a} \int_a^x \left(\bigvee_s^x(f)\right) ds, \end{aligned}$$

$$\begin{aligned} &- \int_x^{s_0} \left(\frac{s-x}{y-x} + \frac{a-s}{b-a} \right) d\left(\bigvee_a^s(f)\right) + \int_{s_0}^y \left(\frac{s-x}{y-x} + \frac{a-s}{b-a} \right) d\left(\bigvee_a^s(f)\right) \quad (2.6) \\ &= - \left(\frac{s-x}{y-x} + \frac{a-s}{b-a} \right) \bigvee_a^s(f) \Big|_{s=x}^{s_0} + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \int_x^{s_0} \left(\bigvee_a^s(f)\right) ds \\ &\quad + \left(\frac{s-x}{y-x} + \frac{a-s}{b-a} \right) \bigvee_a^s(f) \Big|_{s=s_0}^y - \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \int_{s_0}^y \left(\bigvee_a^s(f)\right) ds \\ &= \frac{a-x}{b-a} \bigvee_a^x(f) + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \int_x^{s_0} \left(\bigvee_a^s(f)\right) ds \\ &\quad + \frac{b-y}{b-a} \bigvee_a^y(f) - \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \int_{s_0}^y \left(\bigvee_a^s(f)\right) ds \\ &= \left(\frac{a-x}{b-a} - \frac{s_0-x}{y-x} - \frac{a-s_0}{b-a} \right) \bigvee_a^x(f) + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \int_x^{s_0} \left(\bigvee_a^s(f)\right) ds \\ &\quad + \left(\frac{b-y}{b-a} - \frac{s_0-x}{y-x} + 1 - \frac{a-s_0}{b-a} - 1 \right) \bigvee_a^y(f) - \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \int_{s_0}^y \left(\bigvee_a^s(f)\right) ds \\ &= - \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \int_x^{s_0} \left(\bigvee_a^x(f)\right) ds + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \int_x^{s_0} \left(\bigvee_a^s(f)\right) ds \\ &\quad + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \int_{s_0}^y \left(\bigvee_a^y(f)\right) ds - \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \int_{s_0}^y \left(\bigvee_a^s(f)\right) ds \\ &= \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \left[\int_x^{s_0} \left(\bigvee_a^s(f)\right) ds + \int_{s_0}^y \left(\bigvee_a^s(f)\right) ds \right] \end{aligned}$$

and

$$\begin{aligned}
 \int_y^b \frac{b-s}{b-a} d\left(\bigvee_a^s(f)\right) &= \frac{b-s}{b-a} \bigvee_a^s(f) \Big|_{s=y}^b + \frac{1}{b-a} \int_y^b \left(\bigvee_a^s(f)\right) ds \\
 &= -\frac{b-y}{b-a} \bigvee_a^y(f) + \frac{1}{b-a} \int_y^b \left(\bigvee_a^s(f)\right) ds \\
 &= \frac{1}{b-a} \int_y^b \left(\bigvee_a^s(f)\right) ds - \frac{1}{b-a} \int_y^b \left(\bigvee_a^y(f)\right) ds \\
 &= \frac{1}{b-a} \int_y^b \left(\bigvee_a^s(f)\right) ds
 \end{aligned} \tag{2.7}$$

for all $a \leq x < y \leq b$.

Using (2.4)-(2.7), we deduce the first inequality in (2.2).

Since

$$\begin{aligned}
 \bigvee_s^x(f) &\leq \bigvee_a^x(f) \text{ for } s \in [a, x], & \bigvee_x^s(f) &\leq \bigvee_x^{s_0}(f) \text{ for } s \in [x, s_0], \\
 \bigvee_s^y(f) &\leq \bigvee_{s_0}^y(f) \text{ for } s \in [s_0, y], & \bigvee_y^s(f) &\leq \bigvee_y^b(f) \text{ for } s \in [y, b]
 \end{aligned}$$

and

$$\left(\frac{1}{y-x} - \frac{1}{b-a}\right)(s_0-x) = \frac{x-a}{b-a}, \quad \left(\frac{1}{y-x} - \frac{1}{b-a}\right)(y-s_0) = \frac{b-y}{b-a},$$

then

$$\begin{aligned}
 &\frac{1}{b-a} \int_a^x \left(\bigvee_s^x(f)\right) ds + \left(\frac{1}{y-x} - \frac{1}{b-a}\right) \left[\int_x^{s_0} \left(\bigvee_x^s(f)\right) ds + \int_{s_0}^y \left(\bigvee_s^y(f)\right) ds \right] \\
 &+ \frac{1}{b-a} \int_y^b \left(\bigvee_s^y(f)\right) ds \\
 &\leq \frac{x-a}{b-a} \bigvee_a^x(f) + \left(\frac{1}{y-x} - \frac{1}{b-a}\right) \left[(s_0-x) \bigvee_x^{s_0}(f) + (y-s_0) \bigvee_{s_0}^y(f) \right] + \frac{b-y}{b-a} \bigvee_y^b(f) \\
 &= \frac{x-a}{b-a} \bigvee_a^x(f) + \left[\frac{x-a}{b-a} \bigvee_x^{s_0}(f) + \frac{b-y}{b-a} \bigvee_{s_0}^y(f) \right] + \frac{b-y}{b-a} \bigvee_y^b(f)
 \end{aligned}$$

for all $a \leq x < y \leq b$, which proves the second inequality and the subsequence in (2.2).

The last part is obvious by the max properties and the fact that for $c, d \in \mathbb{R}$ we have

$$\max\{c, d\} = \frac{c+d+|c-d|}{2}.$$

The details are omitted.

To prove the sharpness, we consider the mapping $f : [a, b] \rightarrow \mathbb{R}$ (see [11]) given by

$$f(s) = \begin{cases} 0, & \text{if } s \in [a, x], \\ 1, & \text{if } s \in [x, y], \\ 0, & \text{if } s \in (y, b]. \end{cases}$$

Then f is bounded variation on $[a, b]$, and $\mathcal{V}_a^x(f) = 1$, $\mathcal{V}_x^y(f) = 0$, $\mathcal{V}_x^{s_0}(f) = 0$, $\mathcal{V}_{s_0}^y(f) = 0$, $\mathcal{V}_y^b(f) = 1$, $\int_a^b f(s)ds = y - x$, $\int_x^y f(s)ds = y - x$. Obviously, all the identities hold in (2.2). This completes the proofs. \square

The following corollary holds.

COROLLARY 2.1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a monotonic mapping on $[a, b]$, then we have the inequalities*

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(s)ds - \frac{1}{y-x} \int_x^y f(s)ds \right| \\ & \leq \frac{1}{b-a} \int_a^x |f(x) - f(s)|ds + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \left[\int_x^{s_0} |f(s) - f(x)|ds + \int_{s_0}^y |f(y) - f(s)|ds \right] \\ & \quad + \frac{1}{b-a} \int_y^b |f(s) - f(y)|ds \\ & \leq \frac{x-a}{b-a} |f(x) - f(a)| + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) [(s_0 - x)|f(s_0) - f(x)| + (y - s_0)|f(y) - f(s_0)|] \\ & \quad + \frac{b-y}{b-a} |f(b) - f(y)| \\ & = \frac{x-a}{b-a} |f(x) - f(a)| + \left[\frac{x-a}{b-a} |f(s_0) - f(x)| + \frac{b-y}{b-a} |f(y) - f(s_0)| \right] + \frac{b-y}{b-a} |f(b) - f(y)| \\ & \leq \begin{cases} \frac{x-a}{b-a} |f(x) - f(a)| + \max \left\{ \frac{x-a}{b-a}, \frac{b-y}{b-a} \right\} |f(y) - f(x)| + \frac{b-y}{b-a} |f(b) - f(y)| \\ \frac{x-a}{b-a} |f(x) - f(a)| + \max \{|f(s_0) - f(x)|, |f(y) - f(s_0)|\} + \frac{b-y}{b-a} |f(b) - f(y)| \end{cases} \end{aligned} \tag{2.8}$$

for all $a \leq x < y \leq b$.

REMARK 2.1. Corollary 2.1 gives a refinement of [11, Corollary 1].

The case of Lipschitzian mapping is embodied in the following corollary.

COROLLARY 2.2. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a L -Lipschitzian mapping on $[a, b]$, that is,*

$$|f(x) - f(y)| \leq L|x - y|$$

for positive number L and for all $x, y \in [a, b]$, then we have the inequalities

$$\begin{aligned}
& \left| \frac{1}{b-a} \int_a^b f(s) ds - \frac{1}{y-x} \int_x^y f(s) ds \right| \\
& \leqslant \frac{(x-a)^2 L}{b-a} + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) [(s_0-x)^2 + (y-s_0)^2] L + \frac{(b-y)^2 L}{b-a} \\
& = \frac{L}{b-a} [(x-a)^2 + (x-a)(s_0-x) + (b-y)(y-s_0) + (b-y)^2] \\
& \leqslant \begin{cases} \frac{L}{b-a} [(x-a)^2 + \max\{x-a, b-y\}(y-x) + (b-y)^2] \\ \frac{L}{b-a} [(x-a)^2 + \max\{s_0-x, y-s_0\}(b-a) + (b-y)^2] \end{cases}
\end{aligned} \tag{2.9}$$

for all $a \leqslant x < y \leqslant b$.

REMARK 2.2. Corollary 2.2 gives a refinement of [11, Corollary 2].

Denote $\|f'\|_{1,[s,t]} = \int_s^t |f'(u)| du$, for $s, t \in [a, b]$. Using Theorem 2.1, we have the following corollary immediately.

COROLLARY 2.3. Let $f : [a, b] \rightarrow \mathbb{R}$ be a continuous differentiable on (a, b) , and f' is integrable on (a, b) , then we have the inequalities

$$\begin{aligned}
& \left| \frac{1}{b-a} \int_a^b f(s) ds - \frac{1}{y-x} \int_x^y f(s) ds \right| \\
& \leqslant \frac{1}{b-a} \int_a^x \|f'\|_{1,[s,x]} ds + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) \left[\int_x^{s_0} \|f'\|_{1,[x,s]} ds + \int_{s_0}^y \|f'\|_{1,[s,y]} ds \right] \\
& \quad + \frac{1}{b-a} \int_y^b \|f'\|_{1,[y,s]} ds \\
& \leqslant \frac{x-a}{b-a} \|f'\|_{1,[a,x]} + \left(\frac{1}{y-x} - \frac{1}{b-a} \right) [(s_0-x) \|f'\|_{1,[x,s_0]} + (y-s_0) \|f'\|_{1,[s_0,y]}] \\
& \quad + \frac{b-y}{b-a} \|f'\|_{1,[y,b]} \\
& = \frac{x-a}{b-a} \|f'\|_{1,[a,x]} + \left[\frac{x-a}{b-a} \|f'\|_{1,[x,s_0]} + \frac{b-y}{b-a} \|f'\|_{1,[s_0,y]} \right] + \frac{b-y}{b-a} \|f'\|_{1,[y,b]} \\
& \leqslant \begin{cases} \frac{x-a}{b-a} \|f'\|_{1,[a,x]} + \max \left\{ \frac{x-a}{b-a}, \frac{b-y}{b-a} \right\} \|f'\|_{1,[x,y]} + \frac{b-y}{b-a} \|f'\|_{1,[y,b]} \\ \frac{x-a}{b-a} \|f'\|_{1,[a,x]} + \max \{ \|f'\|_{1,[x,s_0]}, \|f'\|_{1,[s_0,y]} \} + \frac{b-y}{b-a} \|f'\|_{1,[y,b]} \end{cases}
\end{aligned} \tag{2.10}$$

for all $a \leqslant x < y \leqslant b$.

REMARK 2.3. Corollary 2.3 gives a refinement of [11, Corollary 3].

3. Applications for probability density functions

In the following, assume that $f : [a, b] \rightarrow \mathbb{R}^+$ is a probability density function of a certain continuous random variable X and $F(t) = \int_a^t f(x)dx$ is its cumulative distribution function. We shall give some refinements of [11, Propositions 1–4], respectively.

PROPOSITION 3.1. *Let f and F be as above, then we have*

$$\begin{aligned} \left| F(t) - \frac{t-a}{b-a} \right| &\leqslant \frac{b-t}{b-a} \int_a^t \left(\bigvee_s^t (f) \right) ds + \frac{t-a}{b-a} \int_t^b \left(\bigvee_s^b (f) \right) ds \\ &\leqslant \frac{(b-t)(t-a)}{b-a} \bigvee_a^b (f) \end{aligned} \quad (3.1)$$

provided that f is bounded variation on $[a, b]$.

Proof. Taking $x = a$ and $y = t$ in (2.2), we have the desired inequalities. \square

PROPOSITION 3.2. *Let f and F be as above, then we have*

$$\begin{aligned} \left| F(t) - \frac{t-a}{b-a} \right| &\leqslant \frac{b-t}{b-a} \int_a^t |f(t) - f(s)| ds + \frac{t-a}{b-a} \int_t^b |f(s) - f(t)| ds \\ &\leqslant \frac{(b-t)(t-a)}{b-a} (|f(t) - f(a)| + |f(b) - f(t)|) \end{aligned} \quad (3.2)$$

provided that f is a monotonic mapping on $[a, b]$.

Proof. Taking $x = a$ and $y = t$ in (2.8), we have the desired inequalities. \square

PROPOSITION 3.3. *Let f and F be as above, then we have*

$$\left| F(t) - \frac{t-a}{b-a} \right| \leqslant \frac{L}{2} (b-t)(t-a) \quad (3.3)$$

provided that f is a L -Lipschitzian mapping on $[a, b]$.

Proof. Taking $x = a$ and $y = t$ directly in the first inequality of (2.8) (so we have $s_0 = a$), and after calculating two integrals, we get inequality (3.3). \square

PROPOSITION 3.4. *Let f and F be as above, then we have*

$$\begin{aligned} \left| F(t) - \frac{t-a}{b-a} \right| &\leqslant (t-a) \left[\frac{b-t}{b-a} \|f'\|_{1,[a,t]} + \frac{b-t}{b-a} \|f'\|_{1,[t,b]} \right] \\ &\leqslant \frac{(b-t)(t-a)}{b-a} \|f'\|_{1,[a,b]} \end{aligned} \quad (3.4)$$

provided that f is a continuous differentiable on (a, b) and f' is integrable on (a, b) .

Proof. Taking $x = a$ and $y = t$ in (2.10), we have the desired inequality. \square

Acknowledgements. The authors wish to thank the anonymous referees and the editor for their valuable comments.

REFERENCES

- [1] M. W. ALOMARI, *A companion of Ostrowski's inequality for the Riemann-Stieltjes integral $\int_a^b f(t)du(t)$, where f is of bounded variation and u is of r -H-Hölder type and applications*, Appl. Math. Comput. **219** (2013), no. 1, 4792–4799.
- [2] N. S. BARNETT, P. CERONE, S. S. DRAGOMIR AND A. M. FINK, *Comparing two integral means for absolutely continuous mappings whose derivatives are in $L_\infty[a,b]$ and applications*, Comput. Math. Appl. **44** (2002), no. 1–2, 241–251.
- [3] N. S. BARNETT, S. S. DRAGOMIR AND I. GOMM, *A companion for the Ostrowski and the generalized trapezoid inequalities*, Math. Comput. Modelling **50** (2009), no. 1–2, 179–187.
- [4] P. CERONE, W. S. CHEUNG AND S. S. DRAGOMIR, *On Ostrowski type inequalities for Stieltjes integrals with absolutely continuous integrands and integrators of bounded variation*, Comput. Math. Appl. **54** (2007), no. 2, 183–191.
- [5] P. CERONE AND S. S. DRAGOMIR, *Differences between means with bounds from a Riemann-Stieltjes integral*, Comput. Math. Appl. **46** (2003), no. 2–3, 445–453.
- [6] S. S. DRAGOMIR, *Refinements of the generalized trapezoid inequality in terms of the cumulative variation and applications*, RGMIA Res. Rep. Coll. **16** (2013), Article 30, 15 pp.
- [7] S. S. DRAGOMIR, *The Ostrowski integral inequality for mappings of bounded variation*, Bull. Austral. Math. Soc. **60** (1999), no. 3, 495–508.
- [8] S. S. DRAGOMIR, *Refinements of the generalised trapezoid and Ostrowski inequalities for functions of bounded variation*, Arch. Math. (Basel) **91** (2008), no. 5, 450–460.
- [9] S. S. DRAGOMIR, *Refinements of the Ostrowski inequality in terms of the cumulative variation and applications*, RGMIA Res. Rep. Coll., **16** (2013), Article 29, 15 pp.
- [10] V. N. HUY AND Q.-A. NGÔ, *New bounds for the Ostrowski-like type inequalities*, Bull. Korean Math. Soc. **48** (2011), no. 1, 95–104.
- [11] D.-Y. HWANG AND S. S. DRAGOMIR, *Comparing two integral means for mapping of bounded variation and applications*, RGMIA Res. Rep. Coll. **16** (2013), Article 11, 8 pp.
- [12] D.-Y. HWANG AND S. S. DRAGOMIR, *Comparing two integral means for absolutely continuous functions whose absolute value of the derivative are convex and applications*, Appl. Math. Comput. **230** (2014), 259–266.
- [13] D.-Y. HWANG AND S. S. DRAGOMIR, *Some results on comparing two integral means for absolutely continuous functions and applications*, Bull. Aust. Math. Soc. (2014), in press, DOI: 10.1017/S0004972714000021
- [14] D.-Y. HWANG AND S. S. DRAGOMIR, *Comparing two integral means for functions whose absolute value of the derivative are quasi-convex and applications*, RGMIA Res. Rep. Coll., **15** (2012), Article 73, 13 pp.
- [15] D.-Y. HWANG AND S. S. DRAGOMIR, *Difference between the Integral Means Arising from Montgomery's Identity and Applications*, J. Comput. Anal. Appl. **17** (2014), no. 3, 562–573.
- [16] W. J. LIU AND X. GAO, *Approximating the finite Hilbert transform via a companion of Ostrowski's inequality for function of bounded variation and applications*, Appl. Math. Comput. **247** (2014), 373–385.
- [17] W. J. LIU, Q. A. NGO AND W. B. CHEN, *On new Ostrowski type inequalities for double integrals on time scales*, Dynam. Systems Appl. **19** (2010), no. 1, 189–198.
- [18] W. J. LIU, Q. A. NGO AND W. B. CHEN, *Ostrowski type inequalities on time scales for double integrals*, Acta Appl. Math. **110** (2010), no. 1, 477–497.
- [19] Z. LIU, *Another generalization of weighted Ostrowski type inequality for mappings of bounded variation*, Appl. Math. Lett. **25** (2012), no. 3, 393–397.
- [20] M. MATIĆ AND J. PEČARIĆ, *Two-point Ostrowski inequality*, Math. Inequal. Appl. **4** (2001), no. 2, 215–221.

- [21] J. PARK, *Some companions of perturbed Ostrowski type inequalities for functions whose second derivatives are bounded*, Int. J. Appl. Math. Stat. **36** (2013), no. 6, 95–103.
- [22] M. Z. SARIKAYA, *On the Ostrowski type integral inequality*, Acta Math. Univ. Comenian. (N.S.) **79** (2010), no. 1, 129–134.
- [23] K.-L. TSENG, S.-R. HWANG, G.-S. YANG AND Y.-M. CHOU, *Weighted Ostrowski integral inequality for mappings of bounded variation*, Taiwanese J. Math. **15** (2011), no. 2, 573–585.
- [24] K.-L. TSENG, G.-S. YANG AND K.-C. HSU, *Some inequalities for differentiable mappings and applications to Fejér inequality and weighted trapezoidal formula*, Taiwanese J. Math. **15** (2011), no. 4, 1737–1747.
- [25] S. W. VONG, *A note on some Ostrowski-like type inequalities*, Comput. Math. Appl. **62** (2011), no. 1, 532–535.
- [26] W. XU AND H. LIU, *A modified general modulus-based matrix splitting method for linear complementarity problems of H -matrices*, Linear Algebra Appl. **458** (2014), 626–637.

(Received May 27, 2014)

Wenjun Liu

College of Mathematics and Statistics
Nanjing University of Information Science and Technology
Nanjing 210044, China
e-mail: wjliu@nuist.edu.cn

Wangshu Wen

College of Mathematics and Statistics
Nanjing University of Information Science and Technology
Nanjing 210044, China
e-mail: vekestom@gmail.com

Jaekeun Park

Department of Mathematics
Hanseo University
Chungnam-do, Seosan-si 356-706, Republic of Korea
e-mail: jkpark@hanseo.ac.kr