

NEW WEIGHTED ČEBYŠEV-OSTROWSKI TYPE INTEGRAL INEQUALITIES ON TIME SCALES

ADNAN TUNA AND WENJUN LIU

(Communicated by A. Peterson)

Abstract. In this paper we obtain some weighted Čebyšev-Ostrowski type integral inequalities on time scales involving functions whose first derivatives belong to $L_p(a,b)$ ($1 \leq p \leq \infty$). We also give some other interesting inequalities as special cases.

1. Introduction

In [21], Rafiq et al. obtained some weighted Čebyšev-Ostrowski type integral inequalities, involving functions whose first derivatives belong to $L_\infty(a,b)$, as follows.

THEOREM 1.1. *Let $f, g : [a,b] \rightarrow \mathbb{R}$ be absolutely continuous functions whose first derivatives f', g' belong to $L_\infty(a,b)$. Then for all $x \in [a,b]$, we have*

$$\left| \frac{1}{m(a,b)} \int_a^b w(t) f(t) g(t) dt - \left(\frac{1}{m(a,b)} \int_a^b w(t) f(t) dt \right) \left(\frac{1}{m(a,b)} \int_a^b w(t) g(t) dt \right) \right| \\ \leq \frac{\|f'\|_\infty \|g'\|_\infty}{m^3(a,b)} \int_a^b w(x) \left(\int_a^b sgn(t-x) (t-x) w(t) dt \right)^2 dx,$$

and

$$\left| \frac{1}{m(a,b)} \int_a^b w(t) f(t) g(t) dt - \left(\frac{1}{m(a,b)} \int_a^b w(t) f(t) dt \right) \left(\frac{1}{m(a,b)} \int_a^b w(t) g(t) dt \right) \right| \\ \leq \frac{1}{2m^2(a,b)} \int_a^b w(x) [|g(x)| \|f'\|_\infty + |f(x)| \|g'\|_\infty] \left(\int_a^b sgn(t-x) (t-x) w(t) dt \right) dx.$$

Mathematics subject classification (2010): 26D15, 26E70.

Keywords and phrases: Weighted Čebyšev-Ostrowski type inequality, weighted Čebyšev type inequality, time scales.

THEOREM 1.2. Let $f, g : [a, b] \rightarrow \mathbb{R}$ be absolutely continuous functions whose first derivatives f', g' belong to $L_\infty(a, b)$. Then for all $x \in [a, b]$, we have the inequalities

$$\begin{aligned} & \left| FG - \frac{G}{m(a, b)} \int_a^b w(t) f(t) dt - \frac{F}{m(a, b)} \int_a^b w(t) g(t) dt \right. \\ & \quad \left. + \left(\frac{1}{m(a, b)} \int_a^b w(t) f(t) dt \right) \left(\frac{1}{m(a, b)} \int_a^b w(t) g(t) dt \right) \right| \\ & \leq \frac{\|f'\|_\infty \|g'\|_\infty}{m^2(a, b)} \left(\int_a^{\frac{a+b}{2}} sgn\left(x - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, x\right) dx \right. \\ & \quad \left. + \int_{\frac{a+b}{2}}^b sgn\left(x - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, x\right) dx \right)^2, \end{aligned}$$

and

$$\begin{aligned} & \left| \frac{1}{m(a, b)} \int_a^b w(t) [Fg(t) + Gf(t)] dt - 2 \left(\frac{1}{m(a, b)} \int_a^b w(t) f(t) dt \right) \right. \\ & \quad \times \left. \left(\frac{1}{m(a, b)} \int_a^b w(t) g(t) dt \right) \right| \\ & \leq \frac{1}{m^2(a, b)} \int_a^b w(x) (|g(x)| \|f'\|_\infty + |f(x)| \|g'\|_\infty) \\ & \quad \times \left(\int_a^{\frac{a+b}{2}} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) dt + \int_{\frac{a+b}{2}}^b sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) dt \right) dx \end{aligned}$$

where

$$F = m\left(a, \frac{5a+b}{6}\right) f(a) + m\left(\frac{5a+b}{6}, \frac{a+5b}{6}\right) f\left(\frac{a+b}{2}\right) + m\left(\frac{a+5b}{6}, b\right) f(b),$$

$$G = m\left(a, \frac{5a+b}{6}\right) g(a) + m\left(\frac{5a+b}{6}, \frac{a+5b}{6}\right) g\left(\frac{a+b}{2}\right) + m\left(\frac{a+5b}{6}, b\right) g(b),$$

and

$$m(x) = \begin{cases} \int_a^x w(\tau) d\tau, & a \leq x < \frac{a+b}{2}, \\ \int_{\frac{5a+b}{6}}^x w(\tau) d\tau, & \frac{a+b}{2} \leq x \leq b. \end{cases}$$

Ahmad, Mir and Rafiq [2] established weighted Čebyšev-Ostrowski type integral inequalities involving functions whose first derivatives belong to $L_p(a, b)$, ($1 \leq p < \infty$) as follows.

THEOREM 1.3. *Let $f, g : [a, b] \rightarrow \mathbb{R}$ be absolutely continuous functions whose first derivatives f' , g' belong to $L_p(a, b)$ space where $1 \leq p < \infty$. Then we have*

$$\begin{aligned} & \left| \frac{1}{m(a, b)} \int_a^b w(t) f(t) g(t) dt - \frac{1}{m^2(a, b)} \left(\int_a^b w(t) f(t) dt \right) \left(\int_a^b w(t) g(t) dt \right) \right| \\ & \leq \frac{\|f'\|_p \|g'\|_p}{m^3(a, b)} \int_a^b w(x) \left(\int_a^x m^q(a, t) dt + \int_x^b m^q(t, b) dt \right)^{\frac{2}{q}} dx \end{aligned}$$

and

$$\begin{aligned} & \left| \frac{1}{m(a, b)} \int_a^b w(t) f(t) g(t) dt - \frac{1}{m^2(a, b)} \left(\int_a^b w(t) f(t) dt \right) \left(\int_a^b w(t) g(t) dt \right) \right| \\ & \leq \frac{1}{2m^2(a, b)} \left[\int_a^b w(x) \left[|g(x)| \|f'\|_p + |f(x)| \|g'\|_p \right] \left(\int_a^x m^q(a, t) dt + \int_x^b m^q(t, b) dt \right)^{\frac{1}{q}} dx \right] \end{aligned}$$

for all $x \in [a, b]$.

THEOREM 1.4. *Let $f, g : [a, b] \rightarrow \mathbb{R}$ be absolutely continuous functions whose first derivatives f' , g' belong to $L_p(a, b)$ space where $1 \leq p < \infty$. Then we have*

$$\begin{aligned} & \left| FG - \frac{G}{m(a, b)} \int_a^b w(t) f(t) dt - \frac{F}{m(a, b)} \int_a^b w(t) g(t) dt \right. \\ & \quad \left. + \left(\frac{1}{m(a, b)} \int_a^b w(t) f(t) dt \right) \left(\frac{1}{m(a, b)} \int_a^b w(t) g(t) dt \right) \right| \\ & \leq \frac{\|f'\|_p \|g'\|_p}{m^2(a, b)} \left(\int_a^{\frac{5a+b}{6}} m^q \left(t, \frac{5a+b}{6} \right) dt + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} m^q \left(\frac{5a+b}{6}, t \right) dt \right. \\ & \quad \left. + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} m^q \left(t, \frac{a+5b}{6} \right) dt + \int_{\frac{a+5b}{6}}^b m^q \left(\frac{a+5b}{6}, t \right) dt \right)^{\frac{1}{q}}, \end{aligned}$$

and

$$\begin{aligned}
& \left| \frac{1}{m(a,b)} \int_a^b w(t) [Fg(t) + Gf(t)] dt - 2 \frac{1}{m^2(a,b)} \left(\int_a^b w(t) g(t) dt \right) \left(\int_a^b w(t) f(t) dt \right) \right| \\
& \leqslant \frac{1}{m^2(a,b)} \int_a^b \left[w(x) \left(|g(x)| \|f'\|_p + |f(x)| \|g'\|_p \right) \right. \\
& \quad \times \left(\int_a^{\frac{5a+b}{6}} m^q \left(t, \frac{5a+b}{6} \right) dt + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} m^q \left(\frac{5a+b}{6}, t \right) dt \right. \\
& \quad \left. \left. + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} m^q \left(t, \frac{a+5b}{6} \right) dt + \int_{\frac{a+5b}{6}}^b m^q \left(\frac{a+5b}{6}, t \right) dt \right)^{\frac{1}{q}} \right] dx
\end{aligned}$$

where

$$F = m \left(a, \frac{5a+b}{6} \right) f(a) + m \left(\frac{5a+b}{6}, \frac{a+5b}{6} \right) f \left(\frac{a+b}{2} \right) + m \left(\frac{a+5b}{6}, b \right) f(b),$$

$$G = m \left(a, \frac{5a+b}{6} \right) g(a) + m \left(\frac{5a+b}{6}, \frac{a+5b}{6} \right) g \left(\frac{a+b}{2} \right) + m \left(\frac{a+5b}{6}, b \right) g(b),$$

and

$$m(x) = \begin{cases} \int_a^x w(\tau) d\tau, & a \leq x < \frac{a+b}{2} \\ \int_{\frac{5a+b}{6}}^x w(\tau) d\tau, & \frac{a+b}{2} \leq x \leq b. \end{cases}$$

In 1988, Hilger [14] initiated the development of the theory of time scales. Recently, many authors studied the weighted Čebyšev type inequalities, the Grüss inequality, Ostrowski inequalities, Diamond-alpha Grüss type inequalities, the weighted Ostrowski–Grüss inequalities, Ostrowski type inequalities, a generalized Ostrowski's inequality, an Ostrowski-Grüss type inequality, a perturbed Ostrowski-type inequality, the weighted Ostrowski, trapezoid and Grüss type inequalities, Grüss type inequalities, the weighted Ostrowski and Čebyšev type inequalities, the Čebyšev's inequality on time scales (see [1, 3, 6, 7, 8, 9, 10, 12, 14, 15, 16, 17, 18, 19, 22, 23, 21, 24, 25, 26, 27, 28]).

The purpose of this paper is to obtain some weighted Čebyšev-Ostrowski type integral inequalities on time scales involving functions whose first derivatives belong to $L_p(a,b)$ ($1 \leq p \leq \infty$). We also give some other interesting inequalities as special cases.

2. General definitions

In this section we briefly introduce the time scales theory. For further details and proofs we refer the reader to Hilger's Ph. D. thesis [14], the books [5, 6, 13], and the survey [1].

DEFINITION 1. A time scale \mathbb{T} is an arbitrary nonempty closed subset of \mathbb{R} .

We assume throughout that \mathbb{T} has the topology that is inherited from the standard topology on \mathbb{R} . It also assumed throughout that in \mathbb{T} the interval $[a, b]$ means the set $\{t \in \mathbb{T} : a \leq t \leq b\}$ for the points $a < b$ in \mathbb{T} . Since a time scale may not be connected, we need the following concept of jump operators.

DEFINITION 2. For $t \in \mathbb{T}$, we define the forward jump operator $\sigma : \mathbb{T} \rightarrow \mathbb{T}$ by $\sigma(t) = \inf\{s \in \mathbb{T} : s > t\}$, while the backward jump operator $\rho : \mathbb{T} \rightarrow \mathbb{T}$ is defined by $\rho(t) = \inf\{s \in \mathbb{T} : s < t\}$.

The jump operators σ and ρ allow the classification of points in \mathbb{T} as follows.

DEFINITION 3. If $\sigma(t) > t$, then we say that t is right-scattered, while if $\rho(t) < t$ then we say that t is left-scattered. Points that are right-scattered and left-scattered at the same time are called isolated. If $\sigma(t) = t$, then t is called right-dense, and if $\rho(t) = t$ then t is called left-dense. Points that are both right-dense and left-dense are called dense.

DEFINITION 4. The mapping $\mu : \mathbb{T} \rightarrow [0, \infty)$ defined by $\mu(t) = \sigma(t) - t$ is called the graininess function. The set \mathbb{T}^k is defined as follows: if \mathbb{T} has a left-scattered maximum m , then $\mathbb{T}^k = \mathbb{T} - \{m\}$; otherwise, $\mathbb{T}^k = \mathbb{T}$.

If $\mathbb{T} = \mathbb{R}$, then $\mu(t) = 0$, and when $\mathbb{T} = \mathbb{Z}$, we have $\mu(t) = 1$.

DEFINITION 5. Let $f : \mathbb{T} \rightarrow \mathbb{R}$. f is called differentiable at $t \in \mathbb{T}^k$, with (delta) derivative $f^\Delta(t) \in \mathbb{R}$, if for any given $\varepsilon > 0$ there exists a neighborhood U of t such that

$$\left| f(\sigma(t)) - f(s) - f^\Delta(t)[\sigma(t) - s] \right| \leq \varepsilon |\sigma(t) - s|, \quad \forall s \in U.$$

If $\mathbb{T} = \mathbb{R}$, then $f^\Delta(t) = \frac{df(t)}{dt}$, and if $\mathbb{T} = \mathbb{Z}$, then $f^\Delta(t) = f(t+1) - f(t)$.

THEOREM 2.1. Assume $f, g : \mathbb{T} \rightarrow \mathbb{R}$ are differentiable at $t \in \mathbb{T}^k$. Then the product $fg : \mathbb{T} \rightarrow \mathbb{R}$ is differentiable at t with

$$(fg)^\Delta(t) = f^\Delta(t)g(t) + f(\sigma(t))g^\Delta(t).$$

DEFINITION 6. The function $f : \mathbb{T} \rightarrow \mathbb{R}$ is said to be rd-continuous (denote $f \in C_{rd}(\mathbb{T}, \mathbb{R})$), if it is continuous at all right-dense points $t \in \mathbb{T}$ and its left-sided limits exist at all left-dense points $t \in \mathbb{T}$.

It follows from [5, Theorem 1.74] that every rd-continuous function has an anti-derivative.

DEFINITION 7. Let $f \in C_{rd}(\mathbb{T}, \mathbb{R})$. Then $F : \mathbb{T} \rightarrow \mathbb{R}$ is called the antiderivative of f on \mathbb{T} if it satisfies $F^\Delta(t) = f(t)$ for any $t \in \mathbb{T}^k$. In this case, we define the Δ -integral of f as

$$\int_a^b f(s) \Delta s = F(b) - F(a), \quad t \in \mathbb{T}.$$

THEOREM 2.2. Let f, g be rd-continuous, $a, b, c \in \mathbb{T}$ and $\alpha, \beta \in \mathbb{R}$. Then

$$(1) \int_a^b [\alpha f(t) + \beta g(t)] \Delta t = \alpha \int_a^b f(t) \Delta t + \beta \int_a^b g(t) \Delta t,$$

$$(2) \int_a^b f(t) \Delta t = - \int_b^a f(t) \Delta t,$$

$$(3) \int_a^b f(t) \Delta t = \int_a^c f(t) \Delta t + \int_c^b f(t) \Delta t$$

$$(4) \int_a^b f(t) g^\Delta(t) \Delta t = (fg)(b) - (fg)(a) - \int_a^b f^\Delta(t) g(t) \Delta t,$$

THEOREM 2.3. If f is Δ -integrable on $[a, b]$, then so is $|f|$, and

$$\left| \int_a^b f(t) \Delta t \right| \leq \int_a^b |f(t)| \Delta t.$$

DEFINITION 8. Let $h_k : \mathbb{T}^2 \rightarrow \mathbb{R}$, $k \in \mathbb{N}_0$ be defined by $h_0(t, s) = 1$, for all $s, t \in \mathbb{T}$ and then recursively by $h_{k+1}(t, s) = \int_s^t h_k(\tau, s) \Delta \tau$, for all $s, t \in \mathbb{T}$.

THEOREM 2.4. (Hölder's Inequality) Let $a, b \in \mathbb{T}$. For rd-continuous functions $f : [a, b] \rightarrow \mathbb{R}$ we have

$$\int_a^b |f(t)g(t)| \Delta t \leq \left\{ \int_a^b |f(t)|^p \Delta t \right\}^{\frac{1}{p}} \left\{ \int_a^b |g(t)|^q \Delta t \right\}^{\frac{1}{q}}$$

where $p > 1$ and $q = \frac{p}{p-1}$.

3. Main results

THEOREM 3.1. *Let \mathbb{T} be a time scale and $a, b \in \mathbb{T}$ $a < b$ and $w : [a, b] \rightarrow [0, \infty)$ be non-negative and integrable. If $f, g \in C_{rd}^1([a, b], \mathbb{R})$ such that $f^\Delta, g^\Delta \in L^\infty((a, b))$, then for all $t \in [a, b]$, we have*

$$\begin{aligned} & \left| \frac{1}{m(a, b)} \int_a^b w(t) f(t) g(t) \Delta t - \frac{1}{m^2(a, b)} \left(\int_a^b w(t) f(t) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. - \frac{1}{m^2(a, b)} \left(\int_a^b w(t) g(t) \Delta t \right) \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. + \frac{1}{m^2(a, b)} \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\ & \leqslant \frac{MN}{m^3(a, b)} \int_a^b w(x) \left(\int_a^b sgn(t-x)(\sigma(t)-x)w(t) \Delta t \right)^2 \Delta x \end{aligned} \quad (3.1)$$

and

$$\begin{aligned} & \left| \frac{1}{m(a, b)} \int_a^b w(t) f(t) g(t) \Delta t - \frac{1}{2m^2(a, b)} \left(\int_a^b w(t) g(t) \Delta t \right) \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. - \frac{1}{2m^2(a, b)} \left(\int_a^b w(t) f(t) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\ & \leqslant \frac{1}{2m^2(a, b)} \int_a^b w(x) [|g(x)|M + |f(x)|N] \left(\int_a^b sgn(t-x)(\sigma(t)-x)w(s) \Delta t \right) \Delta x \end{aligned} \quad (3.2)$$

where $M = \sup_{a < t < b} |f^\Delta(t)| < \infty$ and $N = \sup_{a < t < b} |g^\Delta(t)| < \infty$.

Proof. We have (see also [9])

$$f(x) - \frac{1}{m(a, b)} \int_a^b w(t) f(\sigma(t)) \Delta t = \frac{1}{m(a, b)} \int_a^b P_w(x, t) f^\Delta(t) \Delta t \quad (3.3)$$

Replacing f by g in (3.3), we have

$$g(x) - \frac{1}{m(a, b)} \int_a^b w(t) g(\sigma(t)) \Delta t = \frac{1}{m(a, b)} \int_a^b P_w(x, t) g^\Delta(t) \Delta t \quad (3.4)$$

where

$$P_w(x, t) = \begin{cases} \int_a^t w(\tau) \Delta\tau, & a \leq t < x, \\ \int_t^b w(\tau) \Delta\tau, & x \leq t \leq b. \end{cases} \quad (3.5)$$

We also have (see also [9])

$$\int_a^b |P_w(x, t)| \Delta s = \int_a^b (\sigma(t) - x) w(t) \operatorname{sgn}(t - x) \Delta t. \quad (3.6)$$

Now, multiplying both sides of (3.3) and (3.4), we obtain

$$\begin{aligned} f(x)g(x) - \frac{f(x)}{m(a, b)} \int_a^b w(t) g(\sigma(t)) \Delta t - \frac{g(x)}{m(a, b)} \int_a^b w(t) f(\sigma(t)) \Delta t \\ + \frac{1}{m^2(a, b)} \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \\ = \frac{1}{m^2(a, b)} \left(\int_a^b P_w(x, t) f^\Delta(t) \Delta t \right) \left(\int_a^b P_w(x, t) g^\Delta(t) \Delta t \right) \end{aligned} \quad (3.7)$$

Multiplying with $\frac{w(x)}{m(a, b)}$ and integrating over $[a, b]$ and taking absolute values, we get

$$\begin{aligned} & \left| \frac{1}{m(a, b)} \int_a^b w(t) f(t) g(t) \Delta t - \frac{1}{m^2(a, b)} \left(\int_a^b w(t) f(t) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. - \frac{1}{m^2(a, b)} \left(\int_a^b w(t) g(t) \Delta t \right) \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. + \frac{1}{m^2(a, b)} \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\ & \leq \frac{1}{m^3(a, b)} \int_a^b w(x) \left(\int_a^b |P_w(x, t)| |f^\Delta(t)| \Delta t \right) \left(\int_a^b |P_w(x, t)| |g^\Delta(t)| \Delta t \right) \Delta x \\ & \leq \frac{MN}{m^3(a, b)} \int_a^b w(x) \left(\int_a^b \operatorname{sgn}(t - x) (\sigma(t) - x) w(t) \Delta t \right)^2 \Delta x \end{aligned} \quad (3.8)$$

which completes the proof of inequality (3.1).

Next, multiplying (3.3) by $g(x)$ and (3.4) by $f(x)$, adding the resulting identities and multiplying the final result with $\frac{w(x)}{2m(a,b)}$ and integrating over $[a,b]$, we have

$$\begin{aligned} & \frac{1}{m(a,b)} \int_a^b w(t) f(t) g(t) \Delta t - \frac{1}{2m^2(a,b)} \left(\int_a^b w(t) g(t) \Delta t \right) \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \\ & - \frac{1}{2m^2(a,b)} \left(\int_a^b w(t) f(t) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \\ = & \frac{1}{2m^2(a,b)} \int_a^b w(x) \left[g(x) \left(\int_a^b P_w(x,t) f^\Delta(t) \Delta t \right) + f(x) \left(\int_a^b P_w(x,t) g^\Delta(t) \Delta t \right) \right] \Delta x \end{aligned} \quad (3.9)$$

taking absolute values, we get

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \int_a^b w(t) f(t) g(t) \Delta t - \frac{1}{2m^2(a,b)} \left(\int_a^b w(t) g(t) \Delta t \right) \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \right. \\ & \left. - \frac{1}{2m^2(a,b)} \left(\int_a^b w(t) f(t) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\ \leqslant & \frac{1}{2m^2(a,b)} \int_a^b w(x) \left[|g(x)| \left(\int_a^b |P_w(x,t)| |f^\Delta(t)| \Delta t \right) \right. \\ & \left. + |f(x)| \left(\int_a^b |P_w(x,t)| |g^\Delta(t)| \Delta t \right) \right] \Delta x \\ \leqslant & \frac{1}{2m^2(a,b)} \int_a^b w(x) [|g(x)| M + |f(x)| N] \left(\int_a^b |P_w(x,t)| \Delta t \right) \Delta x \\ \leqslant & \frac{1}{2m^2(a,b)} \int_a^b w(x) [|g(x)| M + |f(x)| N] \left(\int_a^b sgn(t-x) (\sigma(t)-x) w(t) \Delta t \right) \Delta x \end{aligned} \quad (3.10)$$

Thus, we get the inequality (3.2). \square

COROLLARY 3.1. In the case $\mathbb{T} = \mathbb{R}$ in Theorem 3.1, we have

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \int_a^b w(t) f(t) g(t) dt - \left(\frac{1}{m(a,b)} \int_a^b w(t) f(t) dt \right) \left(\frac{1}{m(a,b)} \int_a^b w(t) g(t) dt \right) \right| \\ \leqslant & \frac{MN}{m^3(a,b)} \int_a^b w(x) \left(\int_a^b sgn(t-x) (t-x) w(t) dt \right)^2 dx \end{aligned} \quad (3.11)$$

and

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \int_a^b w(t) f(t) g(t) dt - \left(\frac{1}{m(a,b)} \int_a^b w(t) g(t) dt \right) \left(\frac{1}{m(a,b)} \int_a^b w(t) f(t) dt \right) \right| \\ & \leq \frac{1}{2m^2(a,b)} \int_a^b w(x) [|g(x)|M + |f(x)|N] \left(\int_a^b sgn(t-x)(t-x)w(t)dt \right) dx \quad (3.12) \end{aligned}$$

where $M = \sup_{a < t < b} |f'(t)| < \infty$ and $N = \sup_{a < t < b} |g'(t)| < \infty$. This inequality can be found in [21] as Theorem 1.

COROLLARY 3.2. In the case $\mathbb{T} = \mathbb{Z}$ in Theorem 3.1, we have

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \sum_{t=a}^{b-1} w(t) f(t) g(t) - \frac{1}{m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) f(t) \right) \left(\sum_{t=a}^{b-1} w(t) g(t+1) \right) \right. \\ & \quad \left. - \frac{1}{m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) g(t) \right) \left(\sum_{t=a}^{b-1} w(t) f(t+1) \right) \right. \\ & \quad \left. + \frac{1}{m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) f(t+1) \right) \left(\sum_{t=a}^{b-1} w(t) g(t+1) \right) \right| \\ & \leq \frac{MN}{m^3(a,b)} \sum_{x=a}^{b-1} w(x) \left(\sum_{t=a}^{b-1} sgn(t-x)(t+1-x)w(t) \right)^2 \quad (3.13) \end{aligned}$$

and

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \sum_{t=a}^{b-1} w(t) f(t) g(t) - \frac{1}{2m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) g(t) \right) \left(\sum_{t=a}^{b-1} w(t) f(t+1) \right) \right. \\ & \quad \left. - \frac{1}{2m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) f(t) \right) \left(\sum_{t=a}^{b-1} w(t) g(t+1) \right) \right| \\ & \leq \frac{1}{2m^2(a,b)} \sum_{x=a}^{b-1} w(x) [|g(x)|M + |f(x)|N] \left(\sum_{t=a}^{b-1} sgn(t-x)(t+1-x)w(t) \right) \quad (3.14) \end{aligned}$$

where $M = \sup_{a < t < b-1} |\Delta f(t)| < \infty$ and $N = \sup_{a < t < b-1} |\Delta g(t)| < \infty$.

COROLLARY 3.3. In the case $\mathbb{T} = q\mathbb{Z} \cup \{0\}$ ($q > 1$) in Theorem 3.1, we have

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \int_{q^m}^{q^n} w(t) f(t) g(t) \Delta t - \frac{1}{m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) f(t) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) g(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. - \frac{1}{m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) g(t) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) f(\sigma(t)) \Delta t \right) \right| \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) f(\sigma(t)) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) g(\sigma(t)) \Delta t \right) \\
& \leq \frac{MN}{m^3(a,b)} \int_{q^m}^{q^n} w(x) \left(\int_{q^m}^{q^n} sgn(t-x)(\sigma(t)-x)w(s) \Delta t \right)^2 \Delta x
\end{aligned} \tag{3.15}$$

and

$$\begin{aligned}
& \left| \frac{1}{m(a,b)} \int_{q^m}^{q^n} w(t) f(t) g(t) \Delta t - \frac{1}{2m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) g(t) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) f(\sigma(t)) \Delta t \right) \right. \\
& \quad \left. - \frac{1}{2m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) f(t) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) g(\sigma(t)) \Delta t \right) \right| \\
& \leq \frac{1}{2m^2(a,b)} \int_{q^m}^{q^n} w(x) [|g(x)|M + |f(x)|N] \left(\int_{q^m}^{q^n} sgn(t-x)(\sigma(t)-x)w(s) \right) \Delta t
\end{aligned} \tag{3.16}$$

where $M = \sup_{q^m < t < q^n} \left| \frac{f(q^{k+1}) - f(q^k)}{(q-1)q^k} \right| < \infty$ and $N = \sup_{q^m < t < q^n} \left| \frac{g(q^{k+1}) - g(q^k)}{(q-1)q^k} \right| < \infty$.

THEOREM 3.2. Let \mathbb{T} be a time scale and $a, b \in \mathbb{T}$ $a < b$ and $w : [a, b] \rightarrow [0, \infty)$ be non-negative and integrable. If $f, g \in C_{rd}^1([a, b], \mathbb{R})$ such that $f^\Delta, g^\Delta \in L^\infty((a, b))$, then for all $t \in [a, b]$, we have

$$\begin{aligned}
& \left| FG - \frac{G}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t - \frac{F}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \right. \\
& \quad \left. + \left(\frac{1}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t \right) \left(\frac{1}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\
& \leq \frac{MN}{m^2(a,b)} \left[\int_a^{\frac{a+b}{2}} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) \Delta t \right. \\
& \quad \left. + \int_{\frac{a+b}{2}}^b sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) \Delta t \right]^2,
\end{aligned} \tag{3.17}$$

and

$$\begin{aligned}
& \left| \frac{F}{m(a,b)} \int_a^b w(t)g(t)\Delta t + \frac{G}{m(a,b)} \int_a^b w(t)f(t)\Delta t \right. \\
& - \frac{1}{m(a,b)} \left(\int_a^b w(t)g(t)\Delta t \right) \left(\int_a^b w(t)f(\sigma(t))\Delta t \right) \\
& \left. - \frac{1}{m(a,b)} \left(\int_a^b w(t)f(t)\Delta t \right) \left(\int_a^b w(t)g(\sigma(t))\Delta t \right) \right| \\
& \leq \frac{1}{m^2(a,b)} \int_a^b w(x) (|g(x)|M + |f(x)|N) \left(\int_a^{\frac{a+b}{2}} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) \Delta t \right. \\
& \quad \left. + \int_{\frac{a+b}{2}}^b sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) \Delta t \right) \Delta x \tag{3.18}
\end{aligned}$$

where $M = \sup_{a < t < b} |f^\Delta(t)| < \infty$, $N = \sup_{a < t < b} |g^\Delta(t)| < \infty$,

$$\begin{aligned}
F &= m\left(a, \frac{5a+b}{6}\right) f(a) + m\left(\frac{5a+b}{6}, \frac{a+5b}{6}\right) f\left(\frac{a+b}{2}\right) + m\left(\frac{a+5b}{6}, b\right) f(b), \\
G &= m\left(a, \frac{5a+b}{6}\right) g(a) + m\left(\frac{5a+b}{6}, \frac{a+5b}{6}\right) g\left(\frac{a+b}{2}\right) + m\left(\frac{a+5b}{6}, b\right) g(b)
\end{aligned}$$

and

$$m(t) = \begin{cases} \int_a^t w(\tau)\Delta\tau, & a \leq t < \frac{a+b}{2} \\ \int_{\frac{5a+b}{6}}^t w(\tau)\Delta\tau, & \frac{a+b}{2} \leq t \leq b. \end{cases}$$

Proof. Using item 4 of Theorem 2.2, we have

$$F - \frac{1}{m(a,b)} \int_a^b w(t)f(\sigma(t))\Delta t = \frac{1}{m(a,b)} \int_a^b m(t)f^\Delta(t)\Delta t \tag{3.19}$$

and

$$G - \frac{1}{m(a,b)} \int_a^b w(t)g(\sigma(t))\Delta t = \frac{1}{m(a,b)} \int_a^b m(t)g^\Delta(t)\Delta t \tag{3.20}$$

where

$$F = m\left(a, \frac{5a+b}{6}\right) f(a) + m\left(\frac{5a+b}{6}, \frac{a+5b}{6}\right) f\left(\frac{a+b}{2}\right) + m\left(\frac{a+5b}{6}, b\right) f(b),$$

$$G = m\left(a, \frac{5a+b}{6}\right) g(a) + m\left(\frac{5a+b}{6}, \frac{a+5b}{6}\right) g\left(\frac{a+b}{2}\right) + m\left(\frac{a+5b}{6}, b\right) g(b)$$

and

$$m(t) = \begin{cases} \int_0^t w(\tau) \Delta \tau, & a \leq t \leq \frac{a+b}{2} \\ \int_{\frac{5a+b}{6}}^t w(\tau) \Delta \tau, & \frac{a+b}{2} < t \leq b. \end{cases} \quad (3.21)$$

We also have

$$\begin{aligned} \int_a^b |m(t)| \Delta t &= \int_a^{\frac{a+b}{2}} \left| \int_{\frac{5a+b}{6}}^t w(\tau) \Delta \tau \right| \Delta t + \int_{\frac{a+b}{2}}^b \left| \int_{\frac{a+5b}{6}}^t w(\tau) \Delta \tau \right| \Delta t \\ &= \int_a^{\frac{5a+b}{6}} \left(\int_t^{\frac{5a+b}{6}} w(\tau) \Delta \tau \right) \Delta t + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} \left(\int_{\frac{5a+b}{6}}^t w(\tau) \Delta \tau \right) \Delta t \\ &\quad + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} \left(\int_t^{\frac{a+5b}{6}} w(\tau) \Delta \tau \right) \Delta t + \int_{\frac{a+5b}{6}}^b \left(\int_{\frac{a+5b}{6}}^t w(\tau) \Delta \tau \right) \Delta t \\ &= \int_a^{\frac{a+b}{2}} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) \Delta t + \int_{\frac{a+b}{2}}^b sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) \Delta t \end{aligned} \quad (3.22)$$

Now, multiplying both sides of (3.19) and (3.20), we obtain

$$\begin{aligned} FG - \frac{G}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t - \frac{F}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \\ + \left(\frac{1}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t \right) \left(\frac{1}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \right) \\ = \frac{1}{m^2(a,b)} \left(\int_a^b m(t) f^\Delta(t) \Delta t \right) \left(\int_a^b m(t) g^\Delta(t) \Delta t \right) \end{aligned} \quad (3.23)$$

taking absolute values and using (3.22), we get the inequality (3.17).

Next, multiplying (3.19) by $g(x)$ and (3.20) by $f(x)$, adding the resulting identities, we have

$$\begin{aligned} & Fg(x) + Gf(x) - \frac{g(x)}{m(a,b)} \int_a^b w(t)f(\sigma(t)) \Delta t - \frac{f(x)}{m(a,b)} \int_a^b w(t)g(\sigma(t)) \Delta t \\ &= \frac{g(x)}{m(a,b)} \int_a^b m(t)f^\Delta(t) \Delta t + \frac{f(x)}{m(a,b)} \int_a^b m(t)g^\Delta(t) \Delta t. \end{aligned} \quad (3.24)$$

Multiplying the final result with $\frac{w(x)}{m(a,b)}$, integrating over $[a,b]$ and taking absolute values, we get

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \int_a^b w(t) [Fg(t) + Gf(t)] \Delta t - \frac{1}{m^2(a,b)} \left(\int_a^b w(t)g(t) \Delta t \right) \left(\int_a^b w(t)f(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. - \frac{1}{m^2(a,b)} \left(\int_a^b w(t)f(t) \Delta t \right) \left(\int_a^b w(t)g(\sigma(t)) \Delta t \right) \right| \\ & \leqslant \frac{1}{m^2(a,b)} \int_a^b w(x) \left(|g(x)| \int_a^b |m(t)| |f^\Delta(t)| \Delta t + |f(x)| \int_a^b |m(t)| |g^\Delta(t)| \Delta t \right) \Delta x \\ & \leqslant \frac{1}{m^2(a,b)} \int_a^b w(x) (|g(x)|M + |f(x)|N) \left(\int_a^b |m(t)| \Delta t \right) \Delta x \\ & \leqslant \frac{1}{m^2(a,b)} \int_a^b w(x) (|g(x)|M + |f(x)|N) \left(\int_a^{\frac{a+b}{2}} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) \Delta t \right. \\ & \quad \left. + \int_{\frac{a+b}{2}}^b sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) \Delta t \right) \Delta x \end{aligned} \quad (3.25)$$

Thus, we obtain the inequality (3.18). \square

COROLLARY 3.4. *In the case $\mathbb{T} = \mathbb{R}$ in Theorem 3.2, we have*

$$\begin{aligned} & FG - \frac{G}{m(a,b)} \int_a^b w(t)f(t) dt - \frac{F}{m(a,b)} \int_a^b w(t)g(t) dt \\ &+ \left(\frac{1}{m(a,b)} \int_a^b w(t)f(t) dt \right) \left(\frac{1}{m(a,b)} \int_a^b w(t)g(t) dt \right) \end{aligned}$$

$$\leq \frac{MN}{m^3(a,b)} \left(\int_a^{\frac{a+b}{2}} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) dt + \int_{\frac{a+b}{2}}^b sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) dt \right)^2 \quad (3.26)$$

and

$$\begin{aligned} & \left| \frac{F}{m(a,b)} \int_a^b w(t)g(t)dt + \frac{G}{m(a,b)} \int_a^b w(t)f(t)dt \right. \\ & \left. - \frac{2}{m(a,b)} \left(\int_a^b w(t)g(t)dt \right) \left(\int_a^b w(t)f(t)dt \right) \right| \\ & \leq \frac{1}{m^2(a,b)} \int_a^b w(x) (|g(x)|M + |f(x)|N) \left(\int_a^{\frac{a+b}{2}} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) dt \right. \\ & \left. + \int_{\frac{a+b}{2}}^b sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) dt \right) dx \end{aligned} \quad (3.27)$$

where $M = \sup_{a < t < b} |f'(t)| < \infty$ and $N = \sup_{a < t < b} |g'(t)| < \infty$.

$$F = m\left(a, \frac{5a+b}{6}\right) f(a) + m\left(\frac{5a+b}{6}, \frac{a+5b}{6}\right) f\left(\frac{a+b}{2}\right) + m\left(\frac{a+5b}{6}, b\right) f(b),$$

$$G = m\left(a, \frac{5a+b}{6}\right) g(a) + m\left(\frac{5a+b}{6}, \frac{a+5b}{6}\right) g\left(\frac{a+b}{2}\right) + m\left(\frac{a+5b}{6}, b\right) g(b).$$

This inequality can be found in [21] as Theorem 2.

COROLLARY 3.5. In the case $\mathbb{T} = \mathbb{Z}$ in Theorem 3.2, we have

$$\begin{aligned} & \left| FG - \frac{G}{m(a,b)} \sum_{t=a}^{b-1} w(t)f(t+1) - \frac{F}{m(a,b)} \sum_{t=a}^{b-1} w(t)g(t+1) \right. \\ & \left. + \left(\frac{1}{m(a,b)} \sum_{t=a}^{b-1} w(t)f(t+1) \right) \left(\frac{1}{m(a,b)} \sum_{t=a}^{b-1} w(t)g(t+1) \right) \right| \end{aligned}$$

$$\begin{aligned} &\leq \frac{MN}{m^3(a,b)} \left(\sum_{t=a}^{\frac{a+b}{2}-1} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) \right. \\ &\quad \left. + \sum_{t=\frac{a+b}{2}-1}^{b-1} sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) \right)^2 \end{aligned} \quad (3.28)$$

and

$$\begin{aligned} &\left| \frac{F}{m(a,b)} \sum_{t=a}^{b-1} w(t)g(t) + \frac{G}{m(a,b)} \sum_{t=a}^{b-1} w(t)f(t) \right. \\ &\quad \left. - \frac{1}{m(a,b)} \left(\sum_{t=a}^{b-1} w(t)g(t) \right) \left(\sum_{t=a}^{b-1} w(t)f(t+1) \right) \right. \\ &\quad \left. - \frac{1}{m(a,b)} \left(\sum_{t=a}^{b-1} w(t)f(t) \right) \left(\sum_{t=a}^{b-1} w(t)g(t+1) \right) \right| \\ &\leq \frac{1}{m^2(a,b)} \sum_{x=a}^{b-1} w(x) (|g(x)|M + |f(x)|N) \left(\sum_{t=a}^{\frac{a+b}{2}-1} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) \right. \\ &\quad \left. + \sum_{t=\frac{a+b}{2}}^{b-1} sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) \right) \end{aligned} \quad (3.29)$$

where $M = \sup_{a < t < b-1} |\Delta f(t)| < \infty$ and $N = \sup_{a < t < b-1} |\Delta g(t)| < \infty$.

$$F = m\left(a, \frac{5a+b}{6}\right) f(a) + m\left(\frac{5a+b}{6}, \frac{a+5b}{6}\right) f\left(\frac{a+b}{2}\right) + m\left(\frac{a+5b}{6}, b\right) f(b),$$

$$G = m\left(a, \frac{5a+b}{6}\right) g(a) + m\left(\frac{5a+b}{6}, \frac{a+5b}{6}\right) g\left(\frac{a+b}{2}\right) + m\left(\frac{a+5b}{6}, b\right) g(b).$$

COROLLARY 3.6. In the case $\mathbb{T} = q^{\mathbb{Z}} \cup \{0\}$ ($q > 1$) in Theorem 3.2, we have

$$\begin{aligned} &\left| FG - \frac{G}{m(a,b)} \int_{q^m}^{q^n} w(t)f(\sigma(t)) \Delta t - \frac{F}{m(a,b)} \int_{q^m}^{q^n} w(t)g(\sigma(t)) \Delta t \right. \\ &\quad \left. + \left(\frac{1}{m(a,b)} \int_{q^m}^{q^n} w(t)f(\sigma(t)) \Delta t \right) \left(\frac{1}{m(a,b)} \int_{q^m}^{q^n} w(t)g(\sigma(t)) \Delta t \right) \right| \end{aligned}$$

$$\begin{aligned} &\leq \frac{MN}{2m^3(a,b)} \left(\int_{q^m}^{\frac{q^m+q^n}{2}} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) \Delta t \right. \\ &\quad \left. + \int_{\frac{q^m+q^n}{2}}^{q^n} sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) \Delta t \right)^2 \end{aligned} \quad (3.30)$$

and

$$\begin{aligned} &\left| \frac{F}{m(a,b)} \int_{q^m}^{q^n} w(t)g(t)\Delta t + \frac{G}{m(a,b)} \int_{q^m}^{q^n} w(t)f(t)\Delta t \right. \\ &\quad \left. - \frac{1}{m(a,b)} \left(\int_{q^m}^{q^n} w(t)g(t)\Delta t \right) \left(\int_{q^m}^{q^n} w(t)f(\sigma(t))\Delta t \right) \right. \\ &\quad \left. - \frac{1}{m(a,b)} \left(\int_{q^m}^{q^n} w(t)f(t)\Delta t \right) \left(\int_{q^m}^{q^n} w(t)g(\sigma(t))\Delta t \right) \right| \\ &\leq \frac{1}{m^2(a,b)} \int_{q^m}^{q^n} w(x) (|g(x)|M + |f(x)|N) \left(\int_{q^m}^{\frac{q^m+q^n}{2}} sgn\left(t - \frac{5a+b}{6}\right) m\left(\frac{5a+b}{6}, t\right) \Delta t \right. \\ &\quad \left. + \int_{\frac{q^m+q^n}{2}}^{q^n} sgn\left(t - \frac{a+5b}{6}\right) m\left(\frac{a+5b}{6}, t\right) \Delta t \right) \Delta x \end{aligned} \quad (3.31)$$

where $M = \sup_{q^m < t < q^n} \left| \frac{f(q^{k+1}) - f(q^k)}{(q-1)q^k} \right| < \infty$ and $N = \sup_{q^m < t < q^n} \left| \frac{g(q^{k+1}) - g(q^k)}{(q-1)q^k} \right| < \infty$.

$$\begin{aligned} F &= m\left(q^m, \frac{5q^m+q^n}{6}\right) f(q^m) + m\left(\frac{5q^m+q^n}{6}, \frac{q^m+5q^n}{6}\right) f\left(\frac{q^m+q^n}{2}\right) \\ &\quad + m\left(\frac{q^m+5q^n}{6}, q^n\right) f(q^n), \\ G &= m\left(q^m, \frac{5q^m+q^n}{6}\right) g(q^m) + m\left(\frac{5q^m+q^n}{6}, \frac{q^m+5q^n}{6}\right) g\left(\frac{q^m+q^n}{2}\right) \\ &\quad + m\left(\frac{q^m+5q^n}{6}, q^n\right) g(q^n). \end{aligned}$$

THEOREM 3.3. Let \mathbb{T} be a time scale and $a, b \in \mathbb{T}$ $a < b$ and $w : [a, b] \rightarrow [0, \infty)$ be non-negative and integrable. If $f, g \in C_{rd}^1([a, b], \mathbb{R})$ such that $f^\Delta, g^\Delta \in L^p([a, b])$,

then for all $t \in [a, b]$, we have

$$\begin{aligned} & \left| \frac{1}{m(a, b)} \int_a^b w(t) f(t) g(t) \Delta t - \frac{1}{m^2(a, b)} \left(\int_a^b w(t) f(t) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. - \frac{1}{m^2(a, b)} \left(\int_a^b w(t) g(t) \Delta t \right) \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. + \frac{1}{m^2(a, b)} \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\ & \leqslant \frac{\|f^\Delta\|_p \|g^\Delta\|_p}{m^3(a, b)} \int_a^b w(x) \left(\int_a^x m^q(a, t) \Delta t + \int_x^b m^q(t, b) \Delta t \right)^{\frac{2}{q}} \Delta x \end{aligned} \quad (3.32)$$

and

$$\begin{aligned} & \left| \frac{1}{m(a, b)} \int_a^b w(t) f(t) g(t) \Delta t - \frac{1}{2m^2(a, b)} \left(\int_a^b w(t) g(t) \Delta t \right) \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. - \frac{1}{2m^2(a, b)} \left(\int_a^b w(t) f(t) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\ & \leqslant \frac{1}{2m^2(a, b)} \int_a^b w(x) \left[|g(x)| \|f^\Delta\|_p + |f(x)| \|g^\Delta\|_p \right] \\ & \quad \times \left(\int_a^x m^q(a, t) \Delta t + \int_x^b m^q(t, b) \Delta t \right)^{\frac{1}{q}} \Delta x \end{aligned} \quad (3.33)$$

where $\|f^\Delta\|_p = \left(\int_a^b |f^\Delta(t)|^p \Delta t \right)^{\frac{1}{p}} < \infty$ and $\|g^\Delta\|_p = \left(\int_a^b |g^\Delta(t)|^p \Delta t \right)^{\frac{1}{p}} < \infty$.

Proof. Using (3.5), we obtain

$$\begin{aligned} \left(\int_a^b |p_w(x, t)|^q \Delta t \right)^{\frac{1}{q}} &= \left(\int_a^x \left| \int_a^t w(u) \Delta u \right|^q \Delta t + \int_x^b \left| \int_b^t w(u) \Delta u \right|^q \Delta t \right)^{\frac{1}{q}} \\ &= \left(\int_a^x \left(\int_a^t w(u) \Delta u \right)^q \Delta t + \int_x^b \left(\int_t^b w(u) \Delta u \right)^q \Delta t \right)^{\frac{1}{q}} \end{aligned}$$

$$= \left(\int_a^x m^q(a, t) \Delta t + \int_x^b m^q(t, b) \Delta t \right)^{\frac{1}{q}}. \quad (3.34)$$

From (3.7), we have

$$\begin{aligned} & f(x)g(x) - \frac{f(x)}{m(a, b)} \int_a^b w(t)g(\sigma(t)) \Delta t - \frac{g(x)}{m(a, b)} \int_a^b w(t)f(\sigma(t)) \Delta t \\ & + \frac{1}{m^2(a, b)} \left(\int_a^b w(t)f(\sigma(t)) \Delta t \right) \left(\int_a^b w(t)g(\sigma(t)) \Delta t \right) \\ & = \frac{1}{m^2(a, b)} \left(\int_a^b P_w(x, t) f^\Delta(t) \Delta t \right) \left(\int_a^b P_w(x, t) g^\Delta(t) \Delta t \right) \end{aligned} \quad (3.35)$$

Multiplying with $\frac{w(x)}{m(a, b)}$, integrating over $[a, b]$ and taking absolute values and using Hölder's inequality on time scales, we have

$$\begin{aligned} & \left| \frac{1}{m(a, b)} \int_a^b w(t)f(t)g(t) \Delta t - \frac{1}{m^2(a, b)} \left(\int_a^b w(t)f(t) \Delta t \right) \left(\int_a^b w(t)g(\sigma(t)) \Delta t \right) \right. \\ & \left. - \frac{1}{m^2(a, b)} \left(\int_a^b w(t)g(t) \Delta t \right) \left(\int_a^b w(t)f(\sigma(t)) \Delta t \right) \right. \\ & \left. + \frac{1}{m^2(a, b)} \left(\int_a^b w(t)f(\sigma(t)) \Delta t \right) \left(\int_a^b w(t)g(\sigma(t)) \Delta t \right) \right| \\ & \leqslant \frac{1}{m^3(a, b)} \int_a^b w(x) \left(\int_a^b |P_w(x, t)| |f^\Delta(t)| \Delta t \right) \left(\int_a^b |P_w(x, t)| |g^\Delta(t)| \Delta t \right) \Delta x \\ & \leqslant \frac{\|f^\Delta\|_p \|g^\Delta\|_p}{m^3(a, b)} \int_a^b w(x) \left(\int_a^b |P_w(x, t)|^q \Delta t \right)^{\frac{2}{q}} \Delta x \end{aligned} \quad (3.36)$$

From (3.36) and (3.34), we get the desired inequality (3.32).

From (3.9), we have

$$\begin{aligned} & \frac{1}{m(a, b)} \int_a^b w(t)f(t)g(t) \Delta t - \frac{1}{2m^2(a, b)} \left(\int_a^b w(t)g(t) \Delta t \right) \left(\int_a^b w(t)f(\sigma(t)) \Delta t \right) \\ & - \frac{1}{2m^2(a, b)} \left(\int_a^b w(t)f(t) \Delta t \right) \left(\int_a^b w(t)g(\sigma(t)) \Delta t \right) \end{aligned}$$

$$= \frac{1}{2m^2(a,b)} \int_a^b w(x) \left[g(x) \left(\int_a^b P_w(x,t) f^\Delta(t) \Delta t \right) + f(x) \left(\int_a^b P_w(x,t) g^\Delta(t) \Delta t \right) \right] \Delta x \quad (3.37)$$

taking absolute values and using Hölder's inequality on time scales, we get

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \int_a^b w(t) f(t) g(t) \Delta t - \frac{1}{2m^2(a,b)} \left(\int_a^b w(t) g(t) \Delta t \right) \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \right. \\ & \quad \left. - \frac{1}{2m^2(a,b)} \left(\int_a^b w(t) f(t) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\ & \leqslant \frac{1}{2m^2(a,b)} \int_a^b w(x) \left[|g(x)| \left(\int_a^b |P_w(x,t)| |f^\Delta(t)| \Delta t \right) \right. \\ & \quad \left. + |f(x)| \left(\int_a^b |P_w(x,t)| |g^\Delta(t)| \Delta t \right) \right] \Delta x \\ & \leqslant \frac{1}{2m^2(a,b)} \int_a^b \left[w(x) |g(x)| \|f^\Delta\|_p \left(\int_a^b |p_w(x,t)|^q \Delta t \right)^{\frac{1}{q}} \right. \\ & \quad \left. + w(x) |f(x)| \|g^\Delta\|_p \left(\int_a^b |p_w(x,t)|^q \Delta t \right)^{\frac{1}{q}} \right] \Delta x \\ & \leqslant \frac{1}{2m^2(a,b)} \int_a^b w(x) \left[|g(x)| \|f^\Delta\|_p + |f(x)| \|g^\Delta\|_p |f(x)| \right] \\ & \quad \times \left(\int_a^x m^q(a,t) \Delta t + \int_x^b m^q(t,b) \Delta t \right)^{\frac{1}{q}} \Delta x \end{aligned}$$

Consequently, we get the inequality (3.33). \square

COROLLARY 3.7. *In the case $\mathbb{T} = \mathbb{R}$ in Theorem 3.3, we have*

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \int_a^b w(t) f(t) g(t) dt - \frac{1}{m^2(a,b)} \left(\int_a^b w(t) f(t) dt \right) \left(\int_a^b w(t) g(t) dt \right) \right| \\ & \leqslant \frac{\|f'\|_p \|g'\|_p}{m^3(a,b)} \int_a^b w(x) \left(\int_a^x m^q(a,t) dt + \int_x^b m^q(t,b) dt \right)^{\frac{2}{q}} dx \quad (3.38) \end{aligned}$$

and

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \int_a^b w(t) f(t) g(t) dt - \frac{1}{m^2(a,b)} \left(\int_a^b w(t) g(t) dt \right) \left(\int_a^b w(t) f(t) dt \right) \right| \\ & \leqslant \frac{1}{2m^2(a,b)} \int_a^b w(x) \left[|g(x)| \|f'\|_p + |f(x)| \|g'\|_p \right] \left(\int_a^x m^q(a,t) dt + \int_x^b m^q(t,b) dt \right)^{\frac{1}{q}} dx \end{aligned} \quad (3.39)$$

where $\|f'\|_p = \left(\int_a^b |f'(t)|^p dt \right)^{\frac{1}{p}} < \infty$ and $\|g'\|_p = \left(\int_a^b |g'(t)|^p dt \right)^{\frac{1}{p}} < \infty$. This inequality can be found in [2] as Theorem 3.

COROLLARY 3.8. In the case $\mathbb{T} = \mathbb{Z}$ in Theorem 3.3, we have

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \sum_{t=a}^{b-1} w(t) f(t) g(t) - \frac{1}{m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) f(t) \right) \left(\sum_{t=a}^{b-1} w(t) g(t+1) \right) \right. \\ & \quad \left. - \frac{1}{m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) g(t) \right) \left(\sum_{t=a}^{b-1} w(t) f(t+1) \right) \right. \\ & \quad \left. + \frac{1}{m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) f(t+1) \right) \left(\sum_{t=a}^{b-1} w(t) g(t+1) \right) \right| \\ & \leqslant \frac{\|\Delta f\|_p \|\Delta g\|_p}{m^3(a,b)} \sum_{x=a}^{b-1} w(x) \left(\sum_{t=a}^{x-1} m^q(a,t) + \sum_{t=x}^{b-1} m^q(t,b) \right)^{\frac{2}{q}} \end{aligned} \quad (3.40)$$

and

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \sum_{t=a}^{b-1} w(t) f(t) g(t) - \frac{1}{2m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) g(t) \right) \left(\sum_{t=a}^{b-1} w(t) f(t+1) \right) \right. \\ & \quad \left. - \frac{1}{2m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) f(t) \right) \left(\sum_{t=a}^{b-1} w(t) g(t+1) \right) \right| \\ & \leqslant \frac{1}{2m^2(a,b)} \sum_{x=a}^{b-1} w(x) \left[|g(x)| \|\Delta f\|_p + |f(x)| \|\Delta g\|_p \right] \left(\sum_{t=a}^{x-1} m^q(a,t) + \sum_{t=x}^{b-1} m^q(t,b) \right)^{\frac{1}{q}} \end{aligned} \quad (3.41)$$

where $\|\Delta f\|_p = \left(\sum_{t=a}^{b-1} |\Delta f(t)|^p \right)^{\frac{1}{p}} < \infty$ and $\|\Delta g\|_p = \left(\sum_{t=a}^{b-1} |\Delta g(t)|^p \right)^{\frac{1}{p}} < \infty$.

COROLLARY 3.9. In the case $\mathbb{T} = q^{\mathbb{Z}} \cup \{0\}$ ($q > 1$) in Theorem 3.3, we have

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \int_{q^m}^{q^n} w(t) f(t) g(t) \Delta t - \frac{1}{m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) f(t) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) g(\sigma(t)) \Delta t \right) \right. \\ & \quad - \frac{1}{m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) g(t) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) f(\sigma(t)) \Delta t \right) \\ & \quad \left. + \frac{1}{m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) f(\sigma(t)) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) g(\sigma(t)) \Delta t \right) \right| \\ & \leqslant \frac{\|f^\Delta\|_p \|g^\Delta\|_p}{m^3(a,b)} \int_{q^m}^{q^n} w(x) \left(\int_{q^m}^x m^q(a,t) \Delta t + \int_x^{q^n} m^q(t,b) \Delta t \right)^{\frac{2}{q}} \Delta x \end{aligned} \quad (3.42)$$

and

$$\begin{aligned} & \left| \frac{1}{m(a,b)} \int_{q^m}^{q^n} w(t) f(t) g(t) \Delta t - \frac{1}{2m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) g(t) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) f(\sigma(t)) \Delta t \right) \right. \\ & \quad - \frac{1}{2m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) f(t) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) g(\sigma(t)) \Delta t \right) \left. \right| \\ & \leqslant \frac{1}{2m^2(a,b)} \int_{q^m}^{q^n} w(x) \left[|g(x)| \|f^\Delta\|_p + |f(x)| \|g^\Delta\|_p \right] \\ & \quad \times \left(\int_{q^m}^x m^q(a,t) \Delta t + \int_x^{q^n} m^q(t,b) \Delta t \right)^{\frac{1}{q}} \Delta x. \end{aligned} \quad (3.43)$$

THEOREM 3.4. Let \mathbb{T} be a time scale and $a, b \in \mathbb{T}$ $a < b$ and $w : [a, b] \rightarrow [0, \infty)$ be non-negative and integrable. If $f, g \in C_{rd}^1([a, b], \mathbb{R})$ such that $f^\Delta, g^\Delta \in L^p([a, b])$, then for all $t \in [a, b]$, we have

$$\begin{aligned} & \left| FG - \frac{G}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t - \frac{F}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \right. \\ & \quad \left. + \left(\frac{1}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t \right) \left(\frac{1}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \end{aligned}$$

$$\begin{aligned}
&\leq \frac{\|f^\Delta\|_p \|g^\Delta\|_p}{m^2(a,b)} \left(\int_a^{\frac{5a+b}{6}} m^q \left(t, \frac{5a+b}{6} \right) \Delta t + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} m^q \left(\frac{5a+b}{6}, t \right) \Delta t \right. \\
&\quad \left. + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} m^q \left(t, \frac{a+5b}{6} \right) \Delta t + \int_{\frac{a+5b}{6}}^b m^q \left(\frac{a+5b}{6}, t \right) \Delta t \right)^{\frac{2}{q}} \tag{3.44}
\end{aligned}$$

and

$$\begin{aligned}
&\left| \frac{1}{m(a,b)} \int_a^b w(t) [Fg(t) + Gf(t)] \Delta t - \frac{1}{m^2(a,b)} \left(\int_a^b w(t) g(t) \Delta t \right) \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \right. \\
&\quad \left. - \frac{1}{m^2(a,b)} \left(\int_a^b w(t) f(t) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\
&\leq \frac{1}{m^2(a,b)} \int_a^b w(x) \left(|g(x)| \|f^\Delta\|_p + |f(x)| \|g^\Delta\|_p \right) \\
&\quad \times \left(\int_a^{\frac{5a+b}{6}} m^q \left(t, \frac{5a+b}{6} \right) \Delta t + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} m^q \left(\frac{5a+b}{6}, t \right) \Delta t \right. \\
&\quad \left. + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} m^q \left(t, \frac{a+5b}{6} \right) \Delta t + \int_{\frac{a+5b}{6}}^b m^q \left(\frac{a+5b}{6}, t \right) \Delta t \right)^{\frac{1}{q}} \Delta x \tag{3.45}
\end{aligned}$$

where $\|f^\Delta\|_p = \left(\int_a^b |f^\Delta(t)|^p \Delta t \right)^{\frac{1}{p}} < \infty$ and $\|g^\Delta\|_p = \left(\int_a^b |g^\Delta(t)|^p \Delta t \right)^{\frac{1}{p}} < \infty$.

Proof. Using (3.21), we get

$$\begin{aligned}
\left(\int_a^b |m(t)|^q \Delta t \right)^{\frac{1}{q}} &= \left(\int_a^{\frac{a+b}{2}} \left| \int_{\frac{5a+b}{6}}^t w(\tau) \Delta \tau \right|^q \Delta t + \int_{\frac{a+b}{2}}^b \left| \int_{\frac{a+5b}{6}}^t w(\tau) \Delta \tau \right|^q \Delta t \right)^{\frac{1}{q}} \\
&= \left(\int_a^{\frac{5a+b}{6}} m^q \left(t, \frac{5a+b}{6} \right) \Delta t + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} m^q \left(\frac{5a+b}{6}, t \right) \Delta t \right. \\
&\quad \left. + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} m^q \left(t, \frac{a+5b}{6} \right) \Delta t + \int_{\frac{a+5b}{6}}^b m^q \left(\frac{a+5b}{6}, t \right) \Delta t \right)^{\frac{1}{q}} \tag{3.46}
\end{aligned}$$

Using (3.23), we have

$$\begin{aligned}
& FG - \frac{G}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t - \frac{F}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \\
& + \left(\frac{1}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t \right) \left(\frac{1}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \right) \\
& = \frac{1}{m^2(a,b)} \left(\int_a^b m(t) f^\Delta(t) \Delta t \right) \left(\int_a^b m(t) g^\Delta(t) \Delta t \right)
\end{aligned} \tag{3.47}$$

taking absolute values and using Hölder's inequality on time scales, we get

$$\begin{aligned}
& \left| FG - \frac{G}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t - \frac{F}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \right. \\
& \quad \left. + \left(\frac{1}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t \right) \left(\frac{1}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\
& \leq \frac{1}{m^2(a,b)} \left(\int_a^b |m(t)| |f^\Delta(t)| \Delta t \right) \left(\int_a^b |m(t)| |g^\Delta(t)| \Delta t \right) \\
& \leq \frac{\|f^\Delta\|_p \|g^\Delta\|_p}{m^2(a,b)} \left(\int_a^b |m(t)|^q \Delta t \right)^{\frac{2}{q}} \\
& \leq \frac{\|f^\Delta\|_p \|g^\Delta\|_p}{m^2(a,b)} \left(\int_a^{\frac{5a+b}{6}} m^q \left(t, \frac{5a+b}{6} \right) \Delta t + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} m^q \left(\frac{5a+b}{6}, t \right) \Delta t \right. \\
& \quad \left. + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} m^q \left(t, \frac{a+5b}{6} \right) \Delta t + \int_{\frac{a+5b}{6}}^b m^q \left(\frac{a+5b}{6}, t \right) \Delta t \right)^{\frac{2}{q}}
\end{aligned}$$

which completes the proof of inequality (3.44).

Using (3.24), we have

$$\begin{aligned}
& Fg(x) + Gf(x) - \frac{g(x)}{m(a,b)} \int_a^b w(t) f(\sigma(t)) \Delta t - \frac{f(x)}{m(a,b)} \int_a^b w(t) g(\sigma(t)) \Delta t \\
& = \frac{g(x)}{m(a,b)} \int_a^b m(t) f^\Delta(t) \Delta t + \frac{f(x)}{m(a,b)} \int_a^b m(t) g^\Delta(t) \Delta t.
\end{aligned} \tag{3.48}$$

Multiplying with $\frac{w(x)}{m(a,b)}$ and integrating over $[a, b]$ and taking absolute values and using Hölder's inequality on time scales, we obtain

$$\begin{aligned}
& \left| \frac{1}{m(a,b)} \int_a^b w(t) [Fg(t) + Gf(t)] \Delta t \right. \\
& \quad - \frac{1}{m^2(a,b)} \left(\int_a^b w(t) g(t) \Delta t \right) \left(\int_a^b w(t) f(\sigma(t)) \Delta t \right) \\
& \quad \left. - \frac{1}{m^2(a,b)} \left(\int_a^b w(t) f(t) \Delta t \right) \left(\int_a^b w(t) g(\sigma(t)) \Delta t \right) \right| \\
& \leq \frac{1}{m^2(a,b)} \int_a^b w(x) \left(|g(x)| \int_a^b |m(t)| |f^\Delta(t)| \Delta t + |f(x)| \int_a^b |m(t)| |g^\Delta(t)| \Delta t \right) \Delta x \\
& \leq \frac{1}{m^2(a,b)} \int_a^b w(x) \left(|g(x)| \|f^\Delta\|_p + |f(x)| \|g^\Delta\|_p \right) \left(\int_a^b |m(t)|^q \Delta t \right)^{\frac{1}{q}} \Delta x \\
& \leq \frac{1}{m^2(a,b)} \int_a^b w(x) \left(|g(x)| \|f^\Delta\|_p + |f(x)| \|g^\Delta\|_p \right) \\
& \quad \times \left(\int_a^{\frac{5a+b}{6}} m^q \left(t, \frac{5a+b}{6} \right) \Delta t + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} m^q \left(\frac{5a+b}{6}, t \right) \Delta t \right. \\
& \quad \left. + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} m^q \left(t, \frac{a+5b}{6} \right) \Delta t + \int_{\frac{a+5b}{6}}^b m^q \left(\frac{a+5b}{6}, t \right) \Delta t \right)^{\frac{1}{q}} \Delta x \tag{3.49}
\end{aligned}$$

□

COROLLARY 3.10. *In the case $\mathbb{T} = \mathbb{R}$ in Theorem 3.4, we have*

$$\begin{aligned}
& \left| FG - \frac{G}{m(a,b)} \int_a^b w(t) f(t) dt - \frac{F}{m(a,b)} \int_a^b w(t) g(t) dt \right. \\
& \quad \left. + \left(\frac{1}{m(a,b)} \int_a^b w(t) f(t) dt \right) \left(\frac{1}{m(a,b)} \int_a^b w(t) g(t) dt \right) \right|
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{\|f'\|_p \|g'\|_p}{m^2(a,b)} \left(\int_a^{\frac{5a+b}{6}} m^q \left(t, \frac{5a+b}{6} \right) dt + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} m^q \left(\frac{5a+b}{6}, t \right) dt \right. \\
&\quad \left. + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} m^q \left(t, \frac{a+5b}{6} \right) dt + \int_{\frac{a+5b}{6}}^b m^q \left(\frac{a+5b}{6}, t \right) dt \right)^{\frac{2}{q}}
\end{aligned} \tag{3.50}$$

and

$$\begin{aligned}
&\left| \frac{1}{m(a,b)} \int_a^b w(t) [Fg(t) + Gf(t)] dt - \frac{2}{m^2(a,b)} \left(\int_a^b w(t) g(t) dt \right) \left(\int_a^b w(t) f(t) dt \right) \right| \\
&\leq \frac{1}{m^2(a,b)} \int_a^b w(x) \left(|g(x)| \|f'\|_p + |f(x)| \|g'\|_p \right) \\
&\quad \times \left(\int_a^{\frac{5a+b}{6}} m^q \left(t, \frac{5a+b}{6} \right) dt + \int_{\frac{5a+b}{6}}^{\frac{a+b}{2}} m^q \left(\frac{5a+b}{6}, t \right) dt \right. \\
&\quad \left. + \int_{\frac{a+b}{2}}^{\frac{a+5b}{6}} m^q \left(t, \frac{a+5b}{6} \right) dt + \int_{\frac{a+5b}{6}}^b m^q \left(\frac{a+5b}{6}, t \right) dt \right)^{\frac{1}{q}} dx
\end{aligned} \tag{3.51}$$

where $\|f'\|_p = \left(\int_a^b |f'(t)|^p dt \right)^{\frac{1}{p}} < \infty$ and $\|g'\|_p = \left(\int_a^b |g'(t)|^p dt \right)^{\frac{1}{p}} < \infty$. This inequality can be found in [2] as Theorem 4.

COROLLARY 3.11. In the case $\mathbb{T} = \mathbb{Z}$ in Theorem 3.4, we have

$$\begin{aligned}
&\left| FG - \frac{G}{m(a,b)} \sum_{t=a}^{b-1} w(t) f(t+1) - \frac{F}{m(a,b)} \sum_{t=a}^{b-1} w(t) g(t+1) \right. \\
&\quad \left. + \left(\frac{1}{m(a,b)} \sum_{t=a}^{b-1} w(t) f(t+1) \right) \left(\frac{1}{m(a,b)} \sum_{t=a}^{b-1} w(t) g(t+1) \right) \right| \\
&\leq \frac{\|\Delta f\|_p \|\Delta g\|_p}{m^2(a,b)} \left(\sum_{t=a}^{\frac{5a+b}{6}-1} m^q \left(t, \frac{5a+b}{6} \right) + \sum_{t=\frac{5a+b}{6}}^{\frac{a+b}{2}-1} m^q \left(\frac{5a+b}{6}, t \right) \right. \\
&\quad \left. + \sum_{t=\frac{a+b}{2}}^{\frac{a+5b}{6}-1} m^q \left(t, \frac{a+5b}{6} \right) + \sum_{t=\frac{a+5b}{6}}^{b-1} m^q \left(\frac{a+5b}{6}, t \right) \right)^{\frac{2}{q}}
\end{aligned} \tag{3.52}$$

and

$$\begin{aligned}
& \left| \frac{1}{m(a,b)} \sum_{t=a}^{b-1} w(t) [Fg(t) + Gf(t)] \Delta t - \frac{1}{m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) g(t) \right) \left(\sum_{t=a}^{b-1} w(t) f(t+1) \right) \right. \\
& \quad \left. - \frac{1}{m^2(a,b)} \left(\sum_{t=a}^{b-1} w(t) f(t) \right) \left(\sum_{t=a}^{b-1} w(t) g(t+1) \right) \right| \\
& \leqslant \frac{1}{m^2(a,b)} \sum_{x=a}^{b-1} w(x) \left(|g(x)| \|\Delta f\|_p + |f(x)| \|\Delta g\|_p \right) \\
& \quad \times \left(\sum_{t=a}^{\frac{5a+b}{6}-1} m^q \left(t, \frac{5a+b}{6} \right) + \sum_{t=\frac{5a+b}{6}}^{\frac{a+b}{2}-1} m^q \left(\frac{5a+b}{6}, t \right) \right. \\
& \quad \left. + \sum_{t=\frac{a+b}{2}}^{\frac{a+5b}{6}-1} m^q \left(t, \frac{a+5b}{6} \right) + \sum_{t=\frac{a+5b}{6}}^{b-1} m^q \left(\frac{a+5b}{6}, t \right) \right)^{\frac{1}{q}} \tag{3.53}
\end{aligned}$$

where $\|\Delta f\|_p = \left(\sum_{t=a}^{b-1} |\Delta f(t)|^p \right)^{\frac{1}{p}} < \infty$ and $\|\Delta g\|_p = \left(\sum_{t=a}^{b-1} |\Delta g(t)|^p \right)^{\frac{1}{p}} < \infty$,

$$F = m \left(a, \frac{5a+b}{6} \right) f(a) + m \left(\frac{5a+b}{6}, \frac{a+5b}{6} \right) f \left(\frac{a+b}{2} \right) + m \left(\frac{a+5b}{6}, b \right) f(b),$$

$$G = m \left(a, \frac{5a+b}{6} \right) g(a) + m \left(\frac{5a+b}{6}, \frac{a+5b}{6} \right) g \left(\frac{a+b}{2} \right) + m \left(\frac{a+5b}{6}, b \right) g(b).$$

COROLLARY 3.12. In the case $\mathbb{T} = q^{\mathbb{Z}} \cup \{0\}$ ($q > 1$) in Theorem 3.4, we have

$$\begin{aligned}
& \left| FG - \frac{G}{m(a,b)} \int_{q^m}^{q^n} w(t) f(\sigma(t)) \Delta t - \frac{F}{m(a,b)} \int_{q^m}^{q^n} w(t) g(\sigma(t)) \Delta t \right. \\
& \quad \left. + \left(\frac{1}{m(a,b)} \int_{q^m}^{q^n} w(t) f(\sigma(t)) \Delta t \right) \left(\frac{1}{m(a,b)} \int_{q^m}^{q^n} w(t) g(\sigma(t)) \Delta t \right) \right| \\
& \leqslant \frac{\|f^\Delta\|_p \|g^\Delta\|_p}{m^2(a,b)} \left(\int_{q^m}^{\frac{5q^m+q^n}{6}} m^q \left(t, \frac{5q^m+q^n}{6} \right) \Delta t + \int_{\frac{5q^m+q^n}{6}}^{\frac{q^m+q^n}{2}} m^q \left(\frac{5q^m+q^n}{6}, t \right) \Delta t \right. \\
& \quad \left. + \int_{\frac{q^m+q^n}{2}}^{\frac{q^m+5q^n}{6}} m^q \left(t, \frac{q^m+5q^n}{6} \right) \Delta t + \int_{\frac{q^m+5q^n}{6}}^{q^n} m^q \left(\frac{q^m+5q^n}{6}, t \right) \Delta t \right)^{\frac{2}{q}} \tag{3.54}
\end{aligned}$$

and

$$\begin{aligned}
& \left| \frac{1}{m(a,b)} \int_{q^m}^{q^n} w(t) [Fg(t) + Gf(t)] \Delta t \right. \\
& \quad - \frac{1}{m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) g(t) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) f(\sigma(t)) \Delta t \right) \\
& \quad - \frac{1}{m^2(a,b)} \left(\int_{q^m}^{q^n} w(t) f(t) \Delta t \right) \left(\int_{q^m}^{q^n} w(t) g(\sigma(t)) \Delta t \right) \left. \right| \\
& \leq \frac{1}{m^2(a,b)} \int_{q^m}^{q^n} w(x) \left(|g(x)| \|f^\Delta\|_p + |f(x)| \|g^\Delta\|_p \right) \\
& \quad \times \left(\int_{q^m}^{\frac{5q^m+q^n}{6}} m^q \left(t, \frac{5q^m+q^n}{6} \right) \Delta t + \int_{\frac{5q^m+q^n}{6}}^{\frac{q^m+q^n}{2}} m^q \left(\frac{5q^m+q^n}{6}, t \right) \Delta t \right. \\
& \quad \left. + \int_{\frac{q^m+q^n}{2}}^{\frac{q^m+5q^n}{6}} m^q \left(t, \frac{q^m+5q^n}{6} \right) \Delta t + \int_{\frac{q^m+5q^n}{6}}^{q^n} m^q \left(\frac{q^m+5q^n}{6}, t \right) \Delta t \right)^{\frac{1}{q}} \Delta x \quad (3.55)
\end{aligned}$$

where $\|f^\Delta\|_p = \left(\int_{q^m}^{q^n} |f^\Delta(t)|^p \Delta t \right)^{\frac{1}{p}} < \infty$ and $\|g^\Delta\|_p = \left(\int_{q^m}^{q^n} |g^\Delta(t)|^p \Delta t \right)^{\frac{1}{p}} < \infty$,

$$\begin{aligned}
F &= m \left(q^m, \frac{5q^m+q^n}{6} \right) f(q^m) + m \left(\frac{5q^m+q^n}{6}, \frac{q^m+5q^n}{6} \right) f \left(\frac{q^m+q^n}{2} \right) \\
&\quad + m \left(\frac{q^m+5q^n}{6}, q^n \right) f(q^n), \\
G &= m \left(q^m, \frac{5q^m+q^n}{6} \right) g(q^m) + m \left(\frac{5q^m+q^n}{6}, \frac{q^m+5q^n}{6} \right) g \left(\frac{q^m+q^n}{2} \right) \\
&\quad + m \left(\frac{q^m+5q^n}{6}, q^n \right) g(q^n).
\end{aligned}$$

REFERENCES

- [1] R. AGARWAL, M. BOHNER, A. PETERSON, *Inequalities on time scales: a survey*, Math. Inequal. Appl., **4** (2001), 535–557.
- [2] F. AHMAD, N. M. MIR, A. RAFIQ, *Weighted Čebyšev Type inequalities involving functions whose first derivatives belong to $L_p(a,b)$, $\forall (1 \leq p < \infty)$* , Kragujevac J. Math. **32** (2009), 13–26.
- [3] G. ANASTASSIOU, *Duality principle of time scales and inequalities*, Appl. Anal., **89** (2010), 1837–1854.
- [4] M. BOHNER, A. PETERSON, *Dynamic Equations on Time Scales*, An Introduction with Applications, Birkhäuser Boston, Inc., Boston, MA, 2001.
- [5] M. BOHNER, A. PETERSON, *Advances in Dynamic Equations on Time Scales*, Birkhäuser Boston, Boston, MA, 2003.
- [6] M. BOHNER, T. MATTHEWS, *The Grüss inequality on time scales*, Commun. Math. Anal., **3** (2007), 1–8.
- [7] M. BOHNER, T. MATTHEWS, *Ostrowski inequalities on time scales*, JIPAM. J. Inequal. Pure Appl. Math., **9** (2008), Article 6, 8 pp.
- [8] M. BOHNER, T. MATTHEWS, A. TUNA, *Diamond-alpha Grüss type inequalities on time scales*, Int. J. Dyn. Syst. Differ. Equ., **3** (2011), 234–247.
- [9] M. BOHNER, T. MATTHEWS, A. TUNA, *Weighted Ostrowski–Grüss inequalities on time scales*, African Diaspora Journal of Mathematics, Volume 12, Number 1, 2011, 89–99.
- [10] C. DINU, *Ostrowski type inequalities on time scales*, An. Univ. Craiova Ser. Mat. Inform., **34** (2007), 43–58.
- [11] S. HILGER, *Ein Maßkettenkalkül mit Anwendung auf Zentrumsmannigfaltigkeiten*, Ph. D. thesis, Universität Würzburg, Würzburg, Germany, 1988.
- [12] B. KARPUZ, U. M. OZKAN, *Generalized Ostrowski's inequality on time scales*, JIPAM. J. Inequal. Pure Appl. Math., **9** (2008), Article 112, 7 pp.
- [13] V. LAKSHMIKANTHAM, S. SIVASUNDARAM, B. KAYMAKCALAN, *Dynamic Systems on Measure Chains*, Mathematics and its Applications, 370, Kluwer Academic Publishers Group, Dordrecht, 1996.
- [14] W. J. LIU, Q.-A. NGÔ, *An Ostrowski–Grüss type inequality on time scales*, Comput. Math. Appl., **58** (2009), 1207–1210.
- [15] W. J. LIU, Q.-A. NGÔ, *A generalization of Ostrowski inequality on time scales for k points*, Appl. Math. Comput., **203** (2008), 754–760.
- [16] W. J. LIU, Q.-A. NGÔ, W. B. CHEN, *A perturbed Ostrowski-type inequality on time scales for k points for functions whose second derivatives are bounded*, J. Inequal. Appl., 2008, Art. ID 597241, 12 pp.
- [17] W. J. LIU, Q.-A. NGÔ, W. B. CHEN, *A new generalization of Ostrowski type inequality on time scales*, An. Științ. Univ. "Ovidius" Constanța, **17** (2009), 101–114.
- [18] W. J. LIU, A. TUNA, *Weighted Ostrowski, trapezoid and Grüss type inequalities on time scales*, J. Math. Inequal., **6** (2012), 381–399.
- [19] Q.-A. NGÔ, W. J. LIU, *A sharp Grüss type inequality on time scales and application to the sharp Ostrowski–Grüss inequality*, Commun. Math. Anal., **6** (2009), 33–41.
- [20] A. OSTROWSKI, *Über die Absolutabweichung einer differentierbaren Funktion von ihrem Integralmittelwert*, Comment. Math. Helv., **10** (1937), 226–227.
- [21] A. RAFIQ, A. M. MIR, F. AHMAD, *Weighted Čebyšev–Ostrowski type inequalities*, Applied mathematics and Mechanics (English Edition), **28** (7) (2007), 901–906.
- [22] M. Z. SARIKAYA, *A note on Grüss type inequalities on time scales*, Dynam. Systems Appl., **17** (2008), 663–666.
- [23] M. Z. SARIKAYA, *New weighted Ostrowski and Čebyšev type inequalities on time scales*, Computers & Mathematics with Applications, Volume 60 Issue 5, September, 2010, 1510–1514.
- [24] M. Z. SARIKAYA, N. AKTAN, H. YILDIRIM, *On weighted Čebyšev–Grüss type inequalities on time scales*, J. Math. Inequal., **2** (2008), 185–195.
- [25] M. R. SIDI AMMI, D. F. M. TORRES, *Combined dynamic Grüss inequalities on time scales*, J. Math. Sci. (N. Y.), **161** (2009), 792–802.
- [26] A. TUNA, D. DAGHAN, *Generalization of Ostrowski and Ostrowski–Grüss type inequalities on time scales*, Comput. Math. Appl., **60** (2010), 803–811.

- [27] A. TUNA, Y. JIANG, W. J. LIU, *Weighted Ostrowski, Ostrowski-Griess and Ostrowski-Čebyšev Type Inequalities on Time Scales*, Publ. Math. Debrecen, **81** (2012), 81–102.
- [28] C. C. YEH, F. H. WONG, H. J. LI, *Čebyšev's inequality on time scales*, Journal of Inequalities in Pure and Applied Mathematics, Volume 6, Issue 1, Article 7, 2005.

(Received August 14, 2014)

Adnan Tuna

Department of Mathematics, Faculty of Science and Arts
University of Niğde
51240, Niğde, Turkey
e-mail: atuna@nigde.edu.tr

Wenjun Liu

College of Mathematics and Statistics
Nanjing University of Information Science and Technology
Nanjing 210044, China
e-mail: wjliu@nuist.edu.cn