

ON THE JAMES TYPE CONSTANT AND VON NEUMANN–JORDAN CONSTANT FOR A CLASS OF BANAŚ–FRĄCZIECK TYPE SPACES

CHANGSEN YANG AND XIANGZHAO YANG

Abstract. As a generalization of Banaś–Frączieck space, the space $X_{\lambda,p}$ that denotes \mathbb{R}^2 endowed with the norm

$$\|x\|_{\lambda,p} = \max\{\lambda|x_1|, \|x\|_p\}$$

for $\lambda > 1$, $p \geq 1$ and $x = (x_1, x_2) \in \mathbb{R}^2$ is well defined. In this note, the exact value of the the James type constants $J_{X_{\lambda,p},t}(1)$ and von Neumann–Jordan constant $C_{NJ}(X_{\lambda,p})$ about this space for $p \geq 2$ are investigated.

Mathematics subject classification (2010): Primary 46B20; Secondary 46B99.

Keywords and phrases: Jordan–von Neumann constant, modulus of convexity, Banaś–Frączieck type space, James type constant.

REFERENCES

- [1] J. A. CLARKSON, *The von Neumann–Jordan constant for the Lebesgue space*, Ann. of Math., **38** (1937), 114–115.
- [2] R. C. JAMES, *Uniformly nonsquare spaces*, Ann. of Math., **80** 1964, 542–550.
- [3] J. GAO, K. S. LAU, *On two classes of Banach spaces with uniform normal structure*, Studia Math. **99**(1991), 41–56.
- [4] Y. TAKAHASHI, *Some geometric constants of Banach spaces – a unified approach*, Banach and function spaces II, Yokohama Publ., Yokohama, (2008), 191–220.
- [5] J. ALONSO, E. LLORENS-FUSTER, *Geometric mean and triangles inscribed in a semicircle in Banach spaces*, J. Math. Anal. Appl. **340** (2008), 1271–1283.
- [6] J. BANAŚ AND K. FRĄCZEK, *Deformation of Banach spaces*, Comment. Math. Univ. Carolinae **34** (1993), no. 1, 47–53.
- [7] C. YANG, *Jordan–von Neumann constant for Banaś–Frączek space*, Banach J. Math. Anal. **8** (2014), no. 2, 185–192.
- [8] C. YANG, Y. WANG, *Some properties of James type constant*, Appl. Math. Lett. **25** (2012), 538–544.