

THE BÉZIER VARIANT OF LUPAS KANTOROVICH OPERATORS BASED ON POLYA DISTRIBUTION

BO-YONG LIAN AND QING-BO CAI

(Communicated by V. Gupta)

Abstract. In this paper we introduce the Bézier variant of Lupas Kantorovich operators based on Polya distribution. We establish a direct approximation by means of the Ditzian-Totik modulus of smoothness and a global approximation theorem in terms of second order modulus of continuity. Furthermore, we give the rate of convergence for absolutely continuous functions having a derivative equivalent to a bounded function. Our results extend the work of Agrawal [P. N. Agrawal, N. Ispir and A. Kajla, Approximation properties of Lupas-Kantorovich operators based on polya distribution, Rendiconti del Circolo Matematico di Palermo Series 2, 2016, 65 (2): 185–208] and Ispir [N. Ispir, P. N. Agrawal and A. Kajla, Rate of convergence of Lupas Kantorovich operators based on Polya distribution, Appl. Math. Comput., 2015, 261: 323–329].

1. Introduction

In the year 1987, Lupas [1] introduced Polya-Bernstein operators

$$P_n^{(1/n)}(f, x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) p_{n,k}^{(1/n)}(x), \quad (1)$$

where $f \in C[0, 1]$, $p_{n,k}^{(1/n)}(x) = \frac{2(n!)}{(2n)!} \binom{n}{k} (nx)_k (n-nx)_{n-k}$, $(n)_k = n(n+1) \cdots (n+k-1)$.

In [2], Miclaus studied some approximation properties of Bernstein-Stancu type operators based on Polya distribution. Recently, Gupta and Rassias [3] introduced the Durrmeyer variant of the operators (1) as follows:

$$D_n^{(1/n)}(f, x) = (n+1) \sum_{k=0}^n p_{n,k}^{(1/n)}(x) \int_0^1 p_{n,k}(t) f(t) dt, \quad (2)$$

where $p_{n,k}(t) = \binom{n}{k} t^k (1-t)^{n-k}$. They established some direct results which include an asymptotic formula, local and global approximation results. To approximate Lebesgue

Mathematics subject classification (2010): 41A36, 41A25, 41A10.

Keywords and phrases: Bézier operators, Polya distribution, rate of convergence, absolutely continuous functions.

This work is supported by the National Natural Science Foundation of China (Grant Nos. 11601266), the Natural Science Foundation of Fujian Province of China (Grant No. 2016J05017), the Program for New Century Excellent Talents in Fujian Province University and the Program for Outstanding Youth Scientific Research Talents in Fujian Province University. We also thank Fujian Provincial Key Laboratory of Data Intensive Computing and Key Laboratory of Intelligent Computing and Information Processing of Fujian Province University.

integrable functions, Agrawal et al. [4] introduced the following integral modification of the operators (1):

$$D_n^{*(1/n)}(f, x) = (n + 1) \sum_{k=0}^n p_{n,k}^{(1/n)}(x) \int_{\frac{k}{n+1}}^{\frac{k+1}{n+1}} f(t) dt. \tag{3}$$

In [4], Agrawal et al. studied the voronovskaja type theorem, local approximation, pointwise estimates and global approximation results. Later, Ispir et al. [5] estimated the rate of convergence for absolutely continuous functions having a derivative coinciding a.e. with a function of bounded variation. For the related research work, we can see [6–9].

It is well known that Bézier curves play an important role in computer aided designs and computer graphics systems. Zeng and Piriou [10] pioneered the study of two Bernstein-Bézier type operators for bounded variation functions. Then many scholars [11–14] have done research work in related fields. Agrawal et al. [15] and Neer et al. [16] introduced the Bézier variant of Lupas-Durrmeyer type operators and genuine-Durrmeyer type operators based on Polya distribution. They obtained some direct approximation theorem and the rate of convergence for absolutely continuous functions having a derivative equivalent to a function of bounded variation.

Inspired by the idea of Zeng and Agrawal, we propose the modified variant of the operators (3) in the following way:

$$D_{n,\alpha}^{*(1/n)}(f, x) = (n + 1) \sum_{k=0}^n Q_{nk}^{(\alpha)}(x) \int_{\frac{k}{n+1}}^{\frac{k+1}{n+1}} f(t) dt, \tag{4}$$

where $\alpha \geq 1$ and

$$Q_{nk}^{(\alpha)}(x) = [J_{n,k}(x)]^\alpha - [J_{n,k+1}(x)]^\alpha, \quad J_{n,k}(x) = \sum_{j=k}^n p_{nj}^{(1/n)}(x).$$

Obviously for $\alpha = 1$, the operators $D_{n,\alpha}^{*(1/n)}(f, x)$ reduces to the operators $D_n^{*(1/n)}(f, x)$.
 Let

$$K_{n,\alpha}(x, t) = \sum_{k=0}^n (n + 1) Q_{nk}^{(\alpha)}(x) \chi_k(t)$$

and

$$R_{n,\alpha}(x, t) = \int_0^t K_{n,\alpha}(x, s) ds,$$

where $\chi_k(t)$ is the characteristic function of the interval $[\frac{k}{n+1}, \frac{k+1}{n+1}]$ with respect to $I = [0, 1]$. By the Lebesgue-Stieltjes integral representations, we have

$$D_{n,\alpha}^{*(1/n)}(f, x) = \int_0^1 f(t) K_{n,\alpha}(x, t) dt = \int_0^1 f(t) d_t R_{n,\alpha}(x, t). \tag{5}$$

The aim of this paper is to establish a direct approximation by means of the Ditzian-Totik modulus of smoothness and a global approximation theorem in terms of second order modulus of continuity. Furthermore, the rate of convergence for some absolutely continuous functions having a derivative equivalent to a bounded function is obtained. The results extend the work of Agrawal and Ispir.

2. Some lemmas

The proof of our results are based on the following lemmas.

LEMMA 1. [4] For $e_i = t^i$, $i = 0, 1, 2$, we have

$$D_n^{*(1/n)}(e_0, x) = 1, \quad D_n^{*(1/n)}(e_1, x) = \frac{2nx + 1}{2(n + 1)},$$

$$D_n^{*(1/n)}(e_2, x) = \frac{3n^3x^2 + 9n^2x - 3n^2x^2 + 3nx + n + 1}{3(n + 1)^3}.$$

By simple applications of Lemma 1, we get

$$D_n^{*(1/n)}(t - x, x) = \frac{1 - 2x}{2(n + 1)}, \tag{6}$$

$$D_n^{*(1/n)}((t - x)^2, x) \leq \frac{2}{n + 1}. \tag{7}$$

LEMMA 2. [4] For $f \in C[0, 1]$, $x \in [0, 1]$, we have

$$\|D_n^{*(1/n)}(f, x)\| \leq \|f\|.$$

LEMMA 3. For $f \in C[0, 1]$, $x \in [0, 1]$, we have

$$\|D_{n,\alpha}^{*(1/n)}(f, x)\| \leq \alpha \|f\|.$$

Proof. Using the inequality $|x^\alpha - y^\alpha| \leq \alpha|x - y|$ with $0 \leq x, y \leq 1$ and $\alpha \geq 1$, we get

$$0 < [J_{n,k}(x)]^\alpha - [J_{n,k+1}(x)]^\alpha \leq \alpha (J_{n,k}(x) - J_{n,k+1}(x)) = \alpha p_{n,k}^{(1/n)}(x).$$

Hence from the definition $D_{n,\alpha}^{*(1/n)}(f, x)$ and Lemma 2, we obtain

$$\|D_{n,\alpha}^{*(1/n)}(f, x)\| \leq \alpha \|D_n^{*(1/n)}(f, x)\| \leq \alpha \|f\|. \quad \square$$

LEMMA 4. (i) For $0 \leq y < x < 1$, there holds

$$R_{n,\alpha}(x, y) = \int_0^y K_{n,\alpha}(x, t) dt \leq \frac{2\alpha}{(n + 1)(x - y)^2}. \tag{8}$$

(ii) For $0 < x < z \leq 1$, there holds

$$1 - R_{n,\alpha}(x, z) = \int_z^1 K_{n,\alpha}(x, t) dt \leq \frac{2\alpha}{(n + 1)(z - x)^2}. \tag{9}$$

Proof. (i) By (5) and (7), we get

$$\begin{aligned}
 R_{n,\alpha}(x,y) &= \int_0^y K_{n,\alpha}(x,t)dt \leq \int_0^y \left(\frac{x-t}{x-y}\right)^2 K_{n,\alpha}(x,t)dt \\
 &\leq \frac{1}{(x-y)^2} \int_0^1 (t-x)^2 K_{n,\alpha}(x,t)dt \\
 &= \frac{1}{(x-y)^2} D_{n,\alpha}^{*(1/n)}((t-x)^2, x) \\
 &\leq \frac{\alpha}{(x-y)^2} D_n^{*(1/n)}((t-x)^2, x) \\
 &\leq \frac{2\alpha}{(n+1)(x-y)^2}.
 \end{aligned}$$

(ii) Using a similar method we can get (9) easily. \square

3. Main results

Let $f(x) \in C[0, 1]$, $t > 0$ and $W^2[0, 1] = \{g \in C[0, 1] : g'' \in C[0, 1]\}$, the appropriate Peetre’s K-functional is defined by

$$K_2(f, t) = \inf_{g \in W^2[0,1]} \{ \|f - g\| + t \|g'\| + t^2 \|g''\| \}.$$

From [17], there exists an absolute constant $C > 0$, such that

$$K_2(f, t) \leq C \omega_2(f, \sqrt{t}), \tag{10}$$

where ω_2 is the second order modulus of continuity of $f \in C[0, 1]$, defined as

$$\omega_2(f, \sqrt{t}) = \sup_{0 < h \leq \sqrt{t}} \sup_{x, x+h, x+2h \in [0,1]} |f(x+2h) - 2f(x+h) + f(x)|.$$

THEOREM 1. For $f \in C[0, 1]$ and $x \in [0, 1]$. Then there exists an absolute constant $C > 0$, such that

$$|D_{n,\alpha}^{*(1/n)}(f, x) - f(x)| \leq C \omega_2 \left(f, \sqrt{\frac{\alpha}{2(n+1)}} \right). \tag{11}$$

Proof. Let $g \in W^2$. By Taylor’s expansion, we get

$$g(t) = g(x) + g'(x)(t-x) + \int_x^t (t-u)g''(u)du.$$

Applying $D_{n,\alpha}^{*(1/n)}(\cdot, x)$ to both sides of the above equation, we have

$$D_{n,\alpha}^{*(1/n)}(g, x) = g(x) + g'(x)D_{n,\alpha}^{*(1/n)}(t-x, x) + D_{n,\alpha}^{*(1/n)}\left(\int_x^t (t-u)g''(u)du, x\right).$$

By Cauchy Schwarz inequality, (7) and Lemma 3, we have

$$\begin{aligned}
 |D_{n,\alpha}^{*(1/n)}(g,x) - g(x)| &\leq |g'(x)| |D_{n,\alpha}^{*(1/n)}(|t-x|,x)| + \left| D_{n,\alpha}^{*(1/n)}\left(\int_x^t (t-u)g''(u)du,x\right) \right| \\
 &\leq \|g'\| |D_{n,\alpha}^{*(1/n)}(|t-x|,x)| + \frac{\|g''\|}{2} |D_{n,\alpha}^{*(1/n)}((t-x)^2,x)| \\
 &\leq \|g'\| |D_{n,\alpha}^{*(1/n)}((t-x)^2,x)|^{1/2} + \frac{\|g''\|}{2} |D_{n,\alpha}^{*(1/n)}((t-x)^2,x)| \\
 &\leq \sqrt{\alpha} \|g'\| |D_n^{*(1/n)}((t-x)^2,x)|^{1/2} + \alpha \frac{\|g''\|}{2} |D_n^{*(1/n)}((t-x)^2,x)| \\
 &\leq \sqrt{\alpha} \|g'\| \sqrt{\frac{2}{n+1}} + \alpha \frac{\|g''\|}{2} \frac{2}{n+1}.
 \end{aligned}$$

Thus

$$\begin{aligned}
 |D_{n,\alpha}^{*(1/n)}(f,x) - f(x)| &\leq |D_{n,\alpha}^{*(1/n)}(f-g,x)| + |f-g| + |D_{n,\alpha}^{*(1/n)}(g,x) - g(x)| \\
 &\leq 2\|f-g\| + \sqrt{\alpha} \|g'\| \sqrt{\frac{2}{n+1}} + \alpha \frac{\|g''\|}{2} \frac{2}{n+1}.
 \end{aligned}$$

Taking the infimum on the right hand side over all $g \in W^2$, we obtain

$$|D_{n,\alpha}^{*(1/n)}(f,x) - f(x)| \leq 2K_2 \left(f, \frac{\alpha}{2(n+1)} \right).$$

By (10), we get (11) immediately. This completes the proof. \square

REMARK 1. When $\alpha = 1$, we have

$$|D_n^{*(1/n)}(f,x) - f(x)| \leq C\omega_2 \left(f, \sqrt{\frac{1}{2(n+1)}} \right),$$

which extend the work of Agrawal et al. [4].

To describe our next result, we recall the definitions of the Ditzian-Totik first order modulus of smoothness and the K-functional [18]. Let $\phi(x) = \sqrt{x(1-x)}$ and $f \in C[0, 1]$, the first order modulus of smoothness is given by

$$\omega_\phi(f,t) = \sup_{0 < h \leq t} \left| f\left(x + \frac{h\phi(x)}{2}\right) - f\left(x - \frac{h\phi(x)}{2}\right) \right|, \quad x \pm \frac{h\phi(x)}{2} \in [0, 1].$$

Further, the corresponding K-functional to $\omega_\phi(f,t)$ is defined by

$$K_\phi(f,t) = \inf_{g \in W_\phi[0,1]} \{ \|f-g\| + t\|\phi g'\| \} (t > 0),$$

where $W_\phi[0, 1] = \{g : g \in AC[0, 1], \|\phi g'\| < \infty\}$ and $AC[0, 1]$ is the class of all absolutely continuous functions on $[0, 1]$. From [18], there exists a constant $C > 0$ such that

$$K_\phi(f,t) \leq C\omega_\phi(f,t). \tag{12}$$

THEOREM 2. For $f \in C[0, 1]$, $x \in (0, 1)$ and $\phi(x) = \sqrt{x(1-x)}$. Then there exists an absolute constant $C > 0$, such that

$$|D_{n,\alpha}^{*(1/n)}(f, x) - f(x)| \leq C\omega_\phi \left(f, \sqrt{\frac{4\alpha}{(n+1)x(1-x)}} \right). \tag{13}$$

Proof. Using the representation

$$g(t) = g(x) + \int_x^t g'(u)du,$$

we get

$$D_{n,\alpha}^{*(1/n)}(g, x) = g(x) + D_{n,\alpha}^{*(1/n)} \left(\int_x^t g'(u)du, x \right).$$

For any $x, t \in (0, 1)$, we find that

$$\left| \int_x^t g'(u)du \right| \leq \|\phi g'\| \left| \int_x^t \frac{1}{\phi(u)} du \right|.$$

But

$$\begin{aligned} \left| \int_x^t \frac{1}{\phi(u)} du \right| &= \left| \int_x^t \frac{1}{\sqrt{u(1-u)}} du \right| \\ &\leq \left| \int_x^t \left(\frac{1}{\sqrt{u}} + \frac{1}{\sqrt{1-u}} \right) du \right| \\ &\leq 2 \left(|\sqrt{t} - \sqrt{x}| + |\sqrt{1-t} - \sqrt{1-x}| \right) \\ &= 2|t-x| \left(\frac{1}{\sqrt{t} + \sqrt{x}} + \frac{1}{\sqrt{1-t} + \sqrt{1-x}} \right) \\ &\leq 2|t-x| \left(\frac{1}{\sqrt{x}} + \frac{1}{\sqrt{1-x}} \right) \leq \frac{2\sqrt{2}|t-x|}{\phi(x)}. \end{aligned}$$

Using Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} |D_{n,\alpha}^{*(1/n)}(g, x) - g(x)| &\leq 2\sqrt{2}\|\phi g'\|\phi^{-1}(x)D_{n,\alpha}^{*(1/n)}(|t-x|, x) \\ &\leq 2\sqrt{2}\|\phi g'\|\phi^{-1}(x) \left(D_{n,\alpha}^{*(1/n)}((t-x)^2, x) \right)^{1/2} \\ &\leq 2\sqrt{2}\sqrt{\alpha}\|\phi g'\|\phi^{-1}(x) \left(D_n^{*(1/n)}((t-x)^2, x) \right)^{1/2} \\ &\leq 4\sqrt{\alpha}\|\phi g'\|\phi^{-1}(x) \sqrt{\frac{1}{n+1}}. \end{aligned}$$

Thus

$$\begin{aligned} |D_{n,\alpha}^{*(1/n)}(f, x) - f(x)| &\leq |D_{n,\alpha}^{*(1/n)}(f-g, x)| + |f-g| + |D_{n,\alpha}^{*(1/n)}(g, x) - g(x)| \\ &\leq 2\|f-g\| + 4\sqrt{\alpha}\|\phi g'\|\phi^{-1}(x) \sqrt{\frac{1}{n+1}}. \end{aligned}$$

Taking the infimum on the right hand side over all $g \in W_\phi[0, 1]$, we obtain

$$|D_{n,\alpha}^{*(1/n)}(f, x) - f(x)| \leq 2K_\phi \left(f, \sqrt{\frac{4\alpha}{(n+1)x(1-x)}} \right).$$

By (12), we get (13) immediately. This completes the proof. \square

REMARK 2. When $\alpha = 1$, we have

$$|D_n^{*(1/n)}(f, x) - f(x)| \leq C\omega_\phi \left(f, \sqrt{\frac{4}{(n+1)x(1-x)}} \right),$$

which extend the work of Agrawal et al. [4].

Lastly, we study the approximation properties of $D_{n,\alpha}^{*(1/n)}(f, x)$ for some absolutely continuous functions $f \in \Phi_{DB}$, which is defined by

$$\Phi_{DB} = \left\{ f \mid f(x) = f(0) + \int_0^x h(t)dt; x \geq 0, h \text{ is bounded on } [0, 1] \right\}.$$

For a bounded function f on $[0, 1]$, we also introduce the following metric form

$$\Omega_x(f, \lambda) = \sup_{t \in [x-\lambda, x+\lambda]} |f(t) - f(x)|, \text{ where } x \in [0, 1] \text{ is fixed, } \lambda \geq 0.$$

It is clear that

- (i) $\Omega_x(f, \lambda)$ is monotone non-decreasing with respect to λ .
- (ii) $\lim_{\lambda \rightarrow 0} \Omega_x(f, \lambda) = 0$, if f is continuous at the point x .

THEOREM 3. Let $f \in \Phi_{DB}$. If $h(x+)$ and $h(x-)$ exist at a fixed point $x \in (0, 1)$, then we have

$$|D_{n,\alpha}^{*(1/n)}(f, x) - f(x)| \leq \alpha(|h(x+)| + |h(x-)|) \sqrt{\frac{2}{n+1}} + \frac{4\alpha + 2x(1-x)}{nx(1-x)} \sum_{k=1}^{[\sqrt{n}]} \Omega_x(\varphi_x, 1/k),$$

where

$$\varphi_x(t) = \begin{cases} h(t) - h(x+), & x < t \leq 1; \\ 0, & t = x; \\ h(t) - h(x-), & 0 \leq t < x. \end{cases}$$

Proof. Let f satisfy the conditions of Theorem 3, by using Bojanic-Cheng's method [19], we have

$$f(t) - f(x) = \int_x^t h(u)du \tag{14}$$

and $h(u)$ can be expressed as

$$\begin{aligned}
 h(u) = & \frac{h(x+) + h(x-)}{2} + \varphi_x(u) + \frac{h(x+) - h(x-)}{2} \text{sign}(u - x) \\
 & + \delta_x(u) \left[h(x) - \frac{h(x+) + h(x-)}{2} \right], \tag{15}
 \end{aligned}$$

where

$$\begin{aligned}
 \delta_x(u) &= \begin{cases} 1, & u = x; \\ 0, & u \neq x. \end{cases} \\
 \text{sign}(x) &= \begin{cases} 1, & x > 0; \\ 0, & x = 0; \\ -1, & x < 0. \end{cases}
 \end{aligned}$$

From (14), (15), and noting $\int_x^t \text{sign}(u - x)du = |t - x|, \int_x^t \delta_x(u)du = 0$, we find that

$$\begin{aligned}
 |D_{n,\alpha}^{*(1/n)}(f, x) - f(x)| &= |D_{n,\alpha}^{*(1/n)}(f(t) - f(x), x)| = \left| D_{n,\alpha}^{*(1/n)} \left(\int_x^t h(u)du, x \right) \right| \\
 &= \left| \frac{h(x+) + h(x-)}{2} D_{n,\alpha}^{*(1/n)}(t - x, x) \right. \\
 &\quad \left. + \frac{h(x+) - h(x-)}{2} D_{n,\alpha}^{*(1/n)}(|t - x|, x) + D_{n,\alpha}^{*(1/n)} \left(\int_x^t \varphi_x(u)du, x \right) \right| \\
 &\leq (|h(x+)| + |h(x-)|) D_{n,\alpha}^{*(1/n)}(|t - x|, x) \\
 &\quad + |D_{n,\alpha}^{*(1/n)} \left(\int_x^t \varphi_x(u)du, x \right)|.
 \end{aligned}$$

By Cauchy Schwarz inequality, (7) and Lemma 3, we have

$$D_{n,\alpha}^{*(1/n)}(|t - x|, x) \leq \alpha D_n^{*(1/n)}(|t - x|, x) \leq \alpha D_n^{*(1/n)}((t - x)^2, x)^{1/2} \leq \alpha \sqrt{\frac{2}{n + 1}}.$$

Thus we have

$$|D_{n,\alpha}^{*(1/n)}(f, x) - f(x)| \leq \alpha (|h(x+)| + |h(x-)|) \sqrt{\frac{2}{n + 1}} + |D_{n,\alpha}^{*(1/n)} \left(\int_x^t \varphi_x(u)du, x \right)|. \tag{16}$$

To complete the proof, we must estimate the term $D_{n,\alpha}^{*(1/n)} \left(\int_x^t \varphi_x(u)du, x \right)$.

From (5), the term $D_{n,\alpha}^{*(1/n)} \left(\int_x^t \varphi_x(u)du, x \right)$ can be stated as

$$\begin{aligned}
 D_{n,\alpha}^{*(1/n)} \left(\int_x^t \varphi_x(u)du, x \right) &= \int_0^1 \left(\int_x^t \varphi_x(u)du \right) K_{n,\alpha}(x, t) dt = \int_0^1 \left(\int_x^t \varphi_x(u)du \right) d_t R_{n,\alpha}(x, t) \\
 &= \int_0^x \left(\int_x^t \varphi_x(u)du \right) d_t R_{n,\alpha}(x, t) + \int_x^1 \left(\int_x^t \varphi_x(u)du \right) d_t R_{n,\alpha}(x, t).
 \end{aligned}$$

Let

$$\Delta_{1n}(f, x) = \int_0^x \left(\int_x^t \varphi_x(u)du \right) d_t R_{n,\alpha}(x, t),$$

$$\Delta_{2n}(f, x) = \int_x^1 \left(\int_x^t \varphi_x(u) du \right) dt R_{n,\alpha}(x, t).$$

Then we have

$$D_{n,\alpha}^{*(1/n)} \left(\int_x^t \varphi_x(u) du, x \right) = \Delta_{1n}(f, x) + \Delta_{2n}(f, x). \tag{17}$$

Using partial integration and noticing $R_{n,\alpha}(x, 0) = 0, \int_x^x \varphi_x(u) du = 0$, we get

$$\begin{aligned} \Delta_{1n}(f, x) &= R_{n,\alpha}(x, t) \int_x^t \varphi_x(u) du \Big|_0^x - \int_0^x R_{n,\alpha}(x, t) \varphi_x(t) dt \\ &= - \int_0^x R_{n,\alpha}(x, t) \varphi_x(t) dt = - \left(\int_0^{x-\frac{x}{\sqrt{n}}} + \int_{x-\frac{x}{\sqrt{n}}}^x \right) R_{n,\alpha}(x, t) \varphi_x(t) dt. \end{aligned}$$

Thus, it follows that

$$|\Delta_{1n}(f, x)| \leq \int_0^{x-\frac{x}{\sqrt{n}}} R_{n,\alpha}(x, t) \Omega_x(\varphi_x, x-t) dt + \int_{x-\frac{x}{\sqrt{n}}}^x R_{n,\alpha}(x, t) \Omega_x(\varphi_x, x-t) dt$$

From Lemma 4 (i) and $0 \leq R_{n,\alpha}(x, t) \leq 1$, we get

$$|\Delta_{1n}(f, x)| \leq \frac{2\alpha}{n+1} \int_0^{x-\frac{x}{\sqrt{n}}} \frac{\Omega_x(\varphi_x, x-t)}{(x-t)^2} dt + \frac{x}{\sqrt{n}} \Omega_x(\varphi_x, x/\sqrt{n}). \tag{18}$$

Putting $t = x - \frac{x}{u}$ for the integral of (18), we get

$$\int_0^{x-\frac{x}{\sqrt{n}}} \frac{\Omega_x(\varphi_x, x-t)}{(x-t)^2} dt = \frac{1}{x} \int_1^{\sqrt{n}} \Omega_x(\varphi_x, x/u) du \leq \frac{2}{x} \sum_{k=1}^{[\sqrt{n}]} \Omega_x(\varphi_x, 1/k). \tag{19}$$

From (17), (18), it follows that

$$\begin{aligned} |\Delta_{1n}(f, x)| &\leq \frac{4\alpha}{(n+1)x} \sum_{k=1}^{[\sqrt{n}]} \Omega_x(\varphi_x, 1/k) + \frac{2x}{n} \sum_{k=1}^{[\sqrt{n}]} \Omega_x(\varphi_x, 1/k) \\ &\leq \frac{4\alpha + 2x^2}{nx} \sum_{k=1}^{[\sqrt{n}]} \Omega_x(\varphi_x, 1/k). \end{aligned} \tag{20}$$

From Lemma 4 (ii), using the same method, we also get

$$|\Delta_{2n}(f, x)| \leq \frac{4\alpha + 2(1-x)^2}{n(1-x)} \sum_{k=1}^{[\sqrt{n}]} \Omega_x(\varphi_x, 1/k). \tag{21}$$

Theorem 3 now follows from (16), (17), (20) and (21). This completes the proof. \square

REMARK 3. If f is a function with derivative of bounded variation, then $f \in \Phi_{DB}$. Thus when $\alpha = 1$, the approximation of $D_n^{*(1/n)}(f, x)$ for functions with derivatives of bounded variation in [5] is a special case of Theorem 3.

REFERENCES

- [1] L. LUPAS AND A. LUPAS, *Polynomials of binomial type and approximation operators*, Stud. Univ. Babeş-Bolyai Math., 1987, **32** (4): 61–69.
- [2] D. MICLAUS, *The revision of some results for Bernstein-Stancu type operators*, Carpathian J. Math., 2012, **28** (2): 289–300.
- [3] V. GUPTA AND T. M. RASSIAS, *Lupas-Durrmeyer operators based on polya distribution*, Banach J. Math. Anal., 2014, **8** (2): 146–155.
- [4] P. N. AGRAWAL, N. ISPIR AND A. KAJLA, *Approximation properties of Lupas-Kantorovich operators based on polya distribution*, Rendiconti del Circolo Matematico di Palermo Series 2, 2016, **65** (2): 185–208.
- [5] N. ISPIR, P. N. AGRAWAL AND A. KAJLA, *Rate of convergence of Lupas Kantorovich operators based on Polya distribution*, Appl. Math. Comput., 2015, **261**: 323–329.
- [6] A. ARAL AND V. GUPTA, *Direct estimates for Lupas-Durrmeyer operators*, Filomat, 2016, **30** (1): 191–199.
- [7] V. GUPTA, A. M. ACU AND D. F. SOFONEA, *Approximation of Baskakov type Polya-Durrmeyer operators*, Appl. Math. Comput., 2017, **294**: 318–331.
- [8] V. GUPTA AND D. SOYBAS, *Convergence of integral operator based on different distributions*, Filomat, 2016, **30** (8): 2277–2287.
- [9] V. GUPTA AND R. P. AGARWAL, *Convergence Estimates in Approximation Theory*, Springer International Publishing, Switzerland, 2014.
- [10] X. M. ZENG AND A. PIRIOU, *On the rate of convergence of two Bernstein-Bézier type operators for bounded variation functions*, J. Approx. Theory, 1998, **95** (3): 369–387.
- [11] X. M. ZENG, *On the rate of convergence of two Bernstein-Bézier type operators for bounded variation functions II*, J. Approx. Theory, 2000, **104** (2): 330–344.
- [12] V. GUPTA AND X. M. ZENG, *Rate of approximation for the Bézier variant of Balazs Kantorovich operators*, Math. Slovaca, 2007, **57** (4): 349–358.
- [13] B. Y. LIAN, *Rate of approximation of bounded variation functions by the Bézier variant of Chlodowsky operators*, J. Math. Inequal., 2013, **7** (4): 647–657.
- [14] X. M. ZENG AND B. Y. LIAN, *An estimate on the convergence of MKZ-Bézier operators*, Comput. Math. Appl., 2008, **56** (12): 3023–3028.
- [15] P. N. AGRAWAL, N. ISPIR AND A. KAJLA, *Approximation properties of Bézier-summation-integral type operators based on Polya-Bernstein functions*, Appl. Math. Comput., 2015, **259**: 533–539.
- [16] T. NEER, A. M. ACU AND P. N. AGRAWAL, *Bézier variant of genuine-Durrmeyer type operators based on Polya distribution*, Carpathian J. Math., 2016, **33** (1): 73–86.
- [17] R. A. DEVORE AND G. G. LORENTZ, *Constructive Approximation*, Springer-Verlag, Berlin, 1993.
- [18] Z. DITZIAN AND V. TOTIK, *Moduli of Smoothness*, Springer, New York, 1987.
- [19] R. BOJANIC AND F. CHENG, *Rate of convergence of Bernstein polynomials for functions with derivatives of bounded variation*, J. Math. Anal. Appl. 1989, **141** (1): 136–151.

(Received August 5, 2017)

Bo-Yong Lian
 Department of Mathematics
 Yang-en University
 Quanzhou 362014, Fujian, China
 e-mail: lianboyong@163.com

Corresponding author:
 Qing-Bo Cai
 School of Mathematics and Computer Science
 Quanzhou Normal University
 Quanzhou 362000, Fujian, China
 e-mail: qbcai@126.com