

IMPROVING SOME INEQUALITIES ASSOCIATED WITH THE EULER-MASCHERONI CONSTANT

JENICA CRINGANU

(Communicated by N. Elezović)

Abstract. The aim of this paper is to improve the results obtained by Chen and Mortici in 2013 about the inequalities for the Euler-Mascheroni constant.

1. Introduction

It is well known that the sequence

$$\gamma_n = 1 + \frac{1}{2} + \dots + \frac{1}{n} - \ln n, \quad n \geq 1,$$

is convergent to a limit denoted $\gamma = 0.5772\dots$ now known as Euler-Mascheroni constant. Many authors have obtained different estimations for $\gamma_n - \gamma$, for example the following increasingly better

$$\frac{1}{2(n+1)} < \gamma_n - \gamma < \frac{1}{2(n-1)}, \quad n \geq 2 \quad ([9]),$$

$$\frac{1}{2(n+1)} < \gamma_n - \gamma < \frac{1}{2n}, \quad n \geq 1 \quad ([12]),$$

$$\frac{1-\gamma}{n} < \gamma_n - \gamma < \frac{1}{2n}, \quad n \geq 1 \quad ([2]),$$

$$\frac{1}{2n+1} < \gamma_n - \gamma < \frac{1}{2n}, \quad n \geq 1 \quad ([6, 7]),$$

$$\frac{1}{2n+\frac{2}{5}} < \gamma_n - \gamma < \frac{1}{2n+\frac{1}{3}}, \quad n \geq 1 \quad ([10]),$$

$$\frac{1}{2n+\frac{2\gamma-1}{1-\gamma}} < \gamma_n - \gamma < \frac{1}{2n+\frac{1}{3}}, \quad n \geq 1 \quad ([1, 10]).$$

Mathematics subject classification (2010): 40A05, 33B15, 11Y60.

Keywords and phrases: Sequence, convergence, Euler-Mascheroni constant.

The convergence of the sequence γ_n to γ is very slow. In 1993, DeTemple [5] studied a modified sequence which converges faster and he proved that

$$\frac{1}{24(n+1)^2} < R_n - \gamma < \frac{1}{24n^2}, \quad n \geq 1,$$

where $R_n = 1 + \frac{1}{2} + \dots + \frac{1}{n} - \ln(n + \frac{1}{2})$.

In 2010, Chen [3] proved that for all integers $n \geq 1$,

$$\frac{1}{24(n+a)^2} \leq R_n - \gamma < \frac{1}{24(n+b)^2},$$

with the best possible constants

$$a = \frac{1}{\sqrt{24[-\gamma + 1 - \ln(\frac{3}{2})]}} - 1 = 0,55106\dots \text{ and } b = \frac{1}{2}.$$

We define the sequence $S_n = \frac{2}{1} + \frac{2}{3} + \dots + \frac{2}{2n-1} - \ln(4n)$, for $n \geq 1$.

Recently, Chen and Mortici [4] obtained the following bounds for $S_n - \gamma$

$$\frac{1}{24(n+a)^2} \leq S_n - \gamma < \frac{1}{24(n+b)^2}, \quad n \geq 1,$$

with the best possible constants

$$a = \frac{1}{\sqrt{24(2 - 2\ln 2 - \gamma)}} - 1 = 0,06858\dots \text{ and } b = 0.$$

In this paper we obtain a better estimation for the left inequality and for the right inequality we remark that $b = 0$ is the best constant using an elementary sequence method.

In 1997, Negoi [8] proved that the sequence $T_n = 1 + \frac{1}{2} + \dots + \frac{1}{n} - \ln(n + \frac{1}{2} + \frac{1}{24n})$ is strictly increasing and convergent to γ . Moreover, he proved that

$$\frac{1}{48(n+1)^3} < \gamma - T_n < \frac{1}{48n^3}, \quad n \geq 1.$$

In 2013, in same paper [4], Chen and Mortici proved that for all integers $n \geq 1$,

$$\frac{1}{48(n+a)^3} \leq \gamma - T_n < \frac{1}{48(n+b)^3},$$

with the best possible constants

$$a = \frac{1}{\sqrt[3]{48[1 - \gamma + \ln(\frac{37}{24})]}} - 1 = 0,27380525\dots \text{ and } b = \frac{83}{360} = 0,23055555\dots$$

The second aim of this paper is to obtain a better estimation for the left inequality and for the right inequality we remark that $b = \frac{83}{360}$ is the best constant also using an elementary sequence method.

2. The main result

THEOREM 1. (i) For every integer $n \geq 1$ we have

$$S_n - \gamma < \frac{1}{24n^2};$$

(ii) For every $a > 0$ there exists $n_a \in N$ such that

$$\frac{1}{24(n+a)^2} < S_n - \gamma \text{ for all } n \geq n_a.$$

Proof. We define the sequence

$$a_n = S_n - \gamma - \frac{1}{24(n+a)^2} = \frac{2}{1} + \frac{2}{3} + \dots + \frac{2}{2n-1} - \ln(4n) - \gamma - \frac{1}{24(n+a)^2}, \text{ for } a \geq 0,$$

and so $a_{n+1} - a_n = f(n)$, where

$$f(n) = \frac{2}{2n+1} - \ln(4n+4) + \ln(4n) - \frac{1}{24(n+a+1)^2} + \frac{1}{24(n+a)^2}.$$

The derivative of function f is equal to

$$f'(n) = \frac{P(n)}{12n(n+1)(2n+1)^2(n+a)^3(n+a+1)^3},$$

where

$$\begin{aligned} P(n) = & 48an^5 + (168a^2 + 120a - 7)n^4 + 2(120a^3 + 168a^2 + 45a - 7)n^3 + 180a^4 \\ & + 360a^3 + 201a^2 + 15a - 8)n^2 + (72a^5 + 180a^4 + 144a^3 + 33a^2 - 3a - 1)n \\ & + 12a^6 + 36a^5 + 36a^4 + 12a^3. \end{aligned}$$

(i) If $a = 0$ then $P(n) = -7n^4 - 14n^3 - 8n^2 - n < 0$, for all $n \geq 1$,

and then f is strictly decreasing. We have $f(\infty) = 0$ and then $f(n) > 0$ for all $n \geq 1$, so that $(a_n)_{n \geq 1}$ is strictly increasing. Since (a_n) converges to zero it results that $a_n < 0$ for all $n \geq 1$, so that

$$S_n - \gamma < \frac{1}{24n^2}, \text{ for all } n \geq 1.$$

(ii) If $a > 0$ then there exists $n_a \in N$ such that $P(n) > 0$ for all $n \geq n_a$ and

then f is strictly increasing on $[n_a, \infty)$. Since $f(\infty) = 0$ it results that $f(n) < 0$ for all $n \geq n_a$, so that $(a_n)_{n \geq n_a}$ is strictly decreasing. The sequence (a_n) converges to zero and then it results that $a_n > 0$ for all $n \geq n_a$, so that

$$\frac{1}{24(n+a)^2} < S_n - \gamma, \text{ for all } n \geq n_a. \quad \square$$

Now we find the constant n_a in some particular cases. For example, if $a = 0,06 = \frac{3}{50} < \frac{1}{\sqrt{24(2-2\ln 2-\gamma)}} - 1 = 0,06858\dots$, then

$$P(n) = \frac{72}{25}n^5 + \frac{503}{625}n^4 - \frac{22933}{3125}n^3 - \frac{491899}{78125}n^2 - \frac{40144813}{39062500}n + \frac{12059037}{3906250000} > 0,$$

for all $n \geq 2$, and so

$$\frac{1}{24(n+\frac{3}{50})^2} < S_n - \gamma < \frac{1}{24n^2}, \text{ for all } n \geq 2.$$

Now, if $a = 0,01 = \frac{1}{100}$, then

$$P(n) = \frac{12}{25}n^5 - \frac{7229}{1250}n^4 - \frac{163327}{12500}n^3 - \frac{39147691}{5000000}n^2 - \frac{1283192741}{12500000000}n + \frac{3090903}{2500000000000} > 0,$$

for all $n \geq 15$, and so

$$\frac{1}{24(n+\frac{1}{100})^2} < S_n - \gamma < \frac{1}{24n^2}, \text{ for all } n \geq 15.$$

Let us remark that a direct calculus show that these inequalities hold and for $n \in \{9, 10, 11, 12, 13, 14\}$, and then

$$\frac{1}{24(n+\frac{1}{100})^2} < S_n - \gamma < \frac{1}{24n^2}, \text{ for all } n \geq 9.$$

THEOREM 2. (i) For every $0 < a \leq \frac{83}{360}$ we have

$$\frac{1}{48(n+a)^3} < \gamma - T_n, \text{ for all } n \geq 1.$$

(ii) For every $a > \frac{83}{360}$ there exists $n_a \in N$ such that

$$\gamma - T_n < \frac{1}{48(n+a)^3}, \text{ for all } n \geq n_a.$$

Consequently $a = \frac{83}{360}$ is the best constant for the inequality of (i).

Proof. The sequence

$$a_n = \gamma - T_n - \frac{1}{48(n+a)^3} = \gamma - 1 - \frac{1}{2} - \dots - \frac{1}{n} + \ln(n + \frac{1}{2} + \frac{1}{24n}) - \frac{1}{48(n+a)^3}, \text{ for } a > 0,$$

converges to zero and it is strictly increasing for $a \geq \frac{83}{360}$ and strictly decreasing for $a < \frac{83}{360}$. For this, we have $a_{n+1} - a_n = f(n)$, where

$$f(n) = -\frac{1}{n+1} + \ln(n + \frac{3}{2} + \frac{1}{24(n+1)}) - \ln(n + \frac{1}{2} + \frac{1}{24n}) - \frac{1}{48(n+a+1)^3} + \frac{1}{48(n+a)^3}.$$

The derivative of function f is equal to

$$f'(n) = \frac{P(n)}{16n(n+1)^2(24n^2+12n+1)(24n^2+60n+37)(n+a)^4(n+a+1)^4},$$

where

$$\begin{aligned} P(n) = & 32(360a - 83)n^9 + (57600a^2 + 47872a - 15963)n^8 + 32(3960a^3 + 7576a^2 \\ & + 2180a - 1183)n^7 + 4(40320a^4 + 121216a^3 + 99856a^2 + 7240a - 12737)n^6 \\ & + 2(64512a^5 + 269920a^4 + 360768a^3 + 160656a^2 - 14062a - 20275)n^5 \\ & + 4(16128a^6 + 91840a^5 + 170840a^4 + 131384a^3 + 31541a^2 - 9197a - 4807)n^4 \\ & + (18432a^7 + 151424a^6 + 376832a^5 + 408320a^4 + 193708a^3 + 19306a^2 \\ & - 15720a - 5211)n^3 + 2(1152a^8 + 17024a^7 + 58656a^6 + 85344a^5 + 58832a^4 \\ & + 16892a^3 - 180a^2 - 1378a - 363)n^2 + (3104a^8 + 17152a^7 + 35200a^6 \\ & + 33728a^5 + 14944a^4 + 2220a^3 - 222a^2 - 148a - 37)n + 592(a^8 + 4a^7 + 6a^6 \\ & + 4a^5 + a^4). \end{aligned}$$

(i) If $a = \frac{83}{360}$ then

$$\begin{aligned} P(n) = & -\frac{27961}{15}n^8 - \frac{29697679}{4050}n^7 - \frac{4043881267}{243000}n^6 - \frac{8530365029807}{437400000}n^5 \\ & - \frac{324974432017513}{26244000000}n^4 - \frac{113729719684675837}{28343520000000}n^3 \\ & - \frac{20572043271868317373}{40814668800000000}n^2 + \frac{126831706407483304897}{8815968460800000000}n \\ & + \frac{67628382210434547877}{17631936921600000000} < 0, \end{aligned}$$

for all $n \geq 1$, and then f is strictly decreasing. Since $f(\infty) = 0$ it results that $f(n) > 0$ for all $n \geq 1$, so that $(a_n)_{n \geq 1}$ is strictly increasing. Since (a_n) converges to zero it results that $a_n < 0$ for all $n \geq 1$, so that

$$\gamma - T_n < \frac{1}{48(n + \frac{83}{360})^3}, \text{ for all } n \geq 1.$$

If $a < \frac{83}{360}$ then $\frac{1}{48(n + \frac{83}{360})^3} < \frac{1}{48(n+a)^3}$ and so

$$\gamma - T_n < \frac{1}{48(n+a)^3}, \text{ for all } n \geq 1.$$

(ii) If $a > \frac{83}{360}$ then there exists $n_a \in N$ such that $P(n) > 0$ for all $n \geq n_a$ and then f is strictly increasing on $[n_a, \infty)$. Since $f(\infty) = 0$ it results that $f(n) < 0$ for all $n \geq n_a$,

so that $(a_n)_{n \geq n_a}$ is strictly decreasing. By convergence to zero of (a_n) it results that $a_n > 0$ for all $n \geq n_a$, so that

$$\frac{1}{48(n+a)^3} < \gamma - T_n, \text{ for all } n \geq n_a. \quad \square$$

Finally we find the constant n_a in some particular cases. For example, if $a = 0,25 = \frac{1}{4} \in (\frac{83}{360}, \frac{1}{\sqrt[3]{48[1-\gamma+\ln(\frac{37}{24})]} - 1})$, then

$$\begin{aligned} P(n) = & 224n^9 - 395n^8 - 3284n^7 - 10538n^6 - \frac{55961}{4}n^5 - \frac{74275}{8}n^4 - \frac{93011}{32}n^3 \\ & - \frac{64655}{256}n^2 + \frac{97921}{2048}n + \frac{23125}{4096} > 0, \end{aligned}$$

for all $n \geq 6$, and so

$$\frac{1}{48(n+\frac{1}{4})^3} < \gamma - T_n < \frac{1}{48(n+\frac{83}{360})^3}, \text{ for all } n \geq 6.$$

Let us remark that a direct calculus show that these inequalities hold and for $n \in \{4, 5\}$, and then

$$\frac{1}{48(n+\frac{1}{4})^3} < \gamma - T_n < \frac{1}{48(n+\frac{83}{360})^3}, \text{ for all } n \geq 4.$$

If $a = 0,24 = \frac{6}{25} \in (\frac{83}{360}, \frac{1}{4})$, then

$$\begin{aligned} P(n) = & \frac{544}{5}n^9 - \frac{28899}{25}n^8 - \frac{16867936}{3125}n^7 - \frac{1074447604}{78125}n^6 - \frac{165272252126}{9765625}n^5 \\ & - \frac{2674071306628}{244140625}n^4 - \frac{21419721292923}{6103515625}n^3 - \frac{59638519190286}{152587890625}n^2 \\ & - \frac{4481000881739}{152587890625}n + \frac{708554863872}{152587890625} > 0, \end{aligned}$$

for all $n \geq 15$, and so

$$\frac{1}{48(n+\frac{6}{25})^3} < \gamma - T_n < \frac{1}{48(n+\frac{83}{360})^3}, \text{ for all } n \geq 15.$$

Moreover, we remark that a direct calculus show that these inequalities hold and for $n \in \{8, 9, 10, 11, 12, 13, 14\}$, and then

$$\frac{1}{48(n+\frac{6}{25})^3} < \gamma - T_n < \frac{1}{48(n+\frac{83}{360})^3}, \text{ for all } n \geq 8.$$

REFERENCES

- [1] H. ALZER, *Inequalities for the gamma and polygamma functions*, Abh. Math. Sem. Univ. Hamburg **68** (1998) 363–372.

- [2] G. D. ANDERSON, R. W. BARNARD, K. C. RICHARDS, M. K. VAMANAMURTHY, M. VUORINEN, *Inequalities for zero-balanced hypergeometric functions*, Trans. Amer. Math. Soc. **345** (1995) 1713–1723.
- [3] C. P. CHEN, *Inequalities for the Euler-Mascheroni constant*, Applied Mathematics Letters **23** (2010) 161–164.
- [4] C. P. CHEN, C. MORTICI, *Limits and inequalities associated with the Euler-Mascheroni constant*, Applied Mathematics and Computation **219** (2013) 9755–9761.
- [5] D. W. DETEMPLE, *A quicker convergence to Euler's constant*, Amer. Math. Monthly **100** (5) (1993) 468–470.
- [6] C. MORTICI, A. VERNESCU, *An improvement of the convergence speed of the sequence $(\gamma_n)_{n \geq 1}$ converging to Euler's constant*, An. Stiint. Univ. "Ovidius" Constanta **13** (1) (2005) 97–100.
- [7] C. MORTICI, A. VERNESCU, *Some new facts in discrete asymptotic analysis*, Math. Balkanica (NS) **21** (Fasc. 3-4) (2007) 301–308.
- [8] T. NEGOI, *A faster convergence to the constant of Euler*, Gazeta Matematica Seria A **15** (1997) 111–113 (in Romanian).
- [9] S. R. TIMS, J. A. TYRREL, *Approximate evaluation of Euler's constant*, Math. Gaz., **55** (1971) 65–67.
- [10] L. TOTH, *Problem E3432*, Amer. Math. Monthly, **98** (3) (1991) 264.
- [11] L. TOTH AND J. BUKOR, *On the alternating series $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$* , J. Math. Anal. Appl. **282** (2003) 21–25.
- [12] R. M. YOUNG, *Euler's constant*, Math. Gaz. **75** (472) (1991) 187–190.

(Received June 29, 2018)

Jenica Cringanu
 "Dunarea de Jos" University of Galati
 Department of Mathematics and Computer Science
 Domneasca No 111, Galati, Romania
 e-mail: jcringanu@ugal.ro