

SOME GENERAL GRADIENT ESTIMATES FOR TWO NONLINEAR PARABOLIC EQUATIONS ALONG RICCI FLOW

WEN WANG*, HUI ZHOU, RULONG XIE AND SONGTING YIN

(Communicated by J. Pečarić)

Abstract. In this paper, by maximum principle and cutoff function, we investigate gradient estimates for positive solutions to two nonlinear parabolic equations along the Ricci flow. As applications, the related Harnack inequalities for positive solutions to the nonlinear parabolic equations along the Ricci flow are derived. These results can be regarded as generalizations of the results of Li-Yau, J. Y. Li, Hamilton and Li-Xu to a more general nonlinear parabolic equation along the Ricci flow. Our results also improve the estimates of S. P. Liu, J. Sun and Y. Y. Yang to the nonlinear parabolic equation along the Ricci flow.

1. Introduction

Beginning with the pioneering work of Li and Yau [14], gradient estimates are also known as differential Harnack inequalities, which have tremendous impact in geometric analysis, as shown for example in [14, 15, 16]. Indeed, they have very important applications in singularity analysis, especially for the Perelman’s breaking work [22, 23] on the Poincaré conjecture.

We first simply introduce research progress associated with this article.

Let (M^n, g) be a complete Riemannian manifold. Li and Yau [14] established a famous gradient estimate for positive solutions to the following heat equation

$$u_t = \Delta u \quad (1.1)$$

on (M^n, g) , which is described as

THEOREM A. (Li-Yau [14]) Let (M^n, g) be a complete Riemannian manifold. Suppose that on the ball $B_R \times (0, T)$ ($B_R := \{(x, t) | d(x, x_0, t) \leq R\}$), $Ricci(B_R) \geq -K$. Then for any $\alpha > 1$,

$$\sup_{B_{2R}} \left(\frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} \right) \leq \frac{C\alpha^2}{R^2} \left(\frac{\alpha^2}{\alpha^2 - 1} + \sqrt{KR} \right) + \frac{n\alpha^2 K}{\alpha - 1} + \frac{n\alpha^2}{2t}. \quad (1.2)$$

Mathematics subject classification (2010): 58J35, 35K05, 53C21.

Keywords and phrases: Gradient estimate, nonlinear parabolic equation, heat equation, Ricci flow, Harnack inequality.

* Corresponding author.

In general, on a complete Riemannian manifold, if $Ricci(M) \geq -k$, by letting $R \rightarrow \infty$ in (1.2), one has

$$\frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} \leq \frac{n\alpha^2 k}{2(\alpha-1)} + \frac{n\alpha^2}{2t}. \quad (1.3)$$

In 1991, Li [15] generalized Li and Yau's estimates to the nonlinear parabolic equation

$$\left(\Delta - \frac{\partial}{\partial t} \right) u(x, t) + h(x, t) u^\alpha(x, t) = 0 \quad (1.4)$$

on (M^n, g) . Hamilton in [8] generalized the constant α of Li and Yau's result to the function $\alpha(t) = e^{2Kt}$. Sun [27] also obtained a gradient estimate with different coefficients. Li and Xu in [17] further promoted Li and Yau's result, and found two new functions $\alpha(t)$. Recently, the first author and Zhang in [28] further generalized Li and Xu's results to the nonlinear parabolic equation (1.4). Related results can be referred to [6, 11, 33].

Motivated by [1, 8, 17, 27], in this paper, we investigate two nonlinear parabolic equations

$$\partial_t u(x, t) = \Delta u(x, t) + h(x, t) u^l(x, t) \quad (1.5)$$

and

$$\partial_t u(x, t) = \Delta u(x, t) + a u(x, t) \log u(x, t) \quad (1.6)$$

along the Ricci flow, where function $h(x, t) \geq 0$ is defined on $M^n \times [0, T]$, which is C^2 in the first variable and C^1 in the second variable, and T is a positive constant and $l, a \in \mathbb{R}$, respectively.

Assume M^n is an n -dimensional manifold without boundary, and let $(M^n, g(t))_{t \in [0, T]}$ be an n -dimensional complete manifold with metric $g(t)$ evolving by the Ricci flow

$$\frac{\partial g(t)}{\partial t} = -2R_{ij}, \quad (x, t) \in M^n \times [0, T]. \quad (1.7)$$

Recently, there are a number of gradient estimates on manifolds along the Ricci flow, see for the example [9, 10] and others, because the Ricci flow is a powerful tool in analyzing the structure of manifolds.

In 2008, Kuang and Zhang [11] proved a gradient estimate for positive solutions to the conjugate heat equation along the Ricci flow on a closed manifold. In 2009, Liu [18] derived a gradient estimate for positive solutions to the heat equation along the Ricci flow. Afterwards, Sun [26] generalized Liu's results to a general geometric flow. In 2010, Bailesteanu, Cao and Pulemotov [2] established some gradient estimates for positive solutions to the heat equation along the Ricci flow. In 2016, Li and Zhu [19] generalized J. Y. Li's [15] estimates along the Ricci flow. Recently, Cao and Zhu [4] derived some Aronson and Bénilan estimates for porous medium equation

$$u_t = \Delta u^m, \quad m > 1$$

along the Ricci flow. Li, Bai and Zhang [13] studied fast diffusion equation

$$u_t = \Delta u^m, \quad 0 < m < 1$$

along the Ricci flow. Zhao and Fang [32] generalized Yang's result [30] to the Ricci flow.

This paper is organized as follows: We prove gradient estimates for the equation (1.5) in Section 2 and gradient estimates for the equation (1.6) in Section 3. We derive related Harnack inequalities in Section 4. As a special case, we deduce gradient estimates and Harnack inequalities to the heat equation in Section 5. Detailed calculation of some specific functions $\alpha(t)$, $\varphi(t)$ and $\gamma(t)$ are given in Section 6.

2. Gradient estimates to the equation (1.5)

In this section, we will derive some new gradient estimates for positive solutions to equation (1.5) along the Ricci flow.

2.1. Main results

Firstly, we introduce three C^1 functions $\alpha(t)$, $\varphi(t)$ and $\gamma(t) : (0, +\infty) \rightarrow (0, +\infty)$. Suppose that three C^1 functions $\alpha(t)$, $\varphi(t)$ and $\gamma(t)$ satisfy the following conditions:

$$(C1) \quad \alpha(t) > 1.$$

$$(C2) \quad \alpha(t) \text{ and } \varphi(t) \text{ satisfy the following system}$$

$$\begin{cases} \frac{2\varphi}{n} - 2\alpha K \geq (\frac{2\varphi}{n} - \alpha')\frac{1}{\alpha}, \\ \frac{2\varphi}{n} - \alpha' > 0, \\ \frac{\varphi^2}{n} + \alpha\varphi' \geq 0. \end{cases}$$

$$(C3) \quad \gamma(t) \text{ satisfies}$$

$$\frac{\gamma'}{\gamma} - (\frac{2\varphi}{n} - \alpha')\frac{1}{\alpha} \leq 0.$$

$$(C4) \quad \gamma(t) \text{ is non-decreasing, and } \alpha(t) \text{ is also non-decreasing or is bounded uniformly.}$$

$$\text{Here } \alpha' = \frac{d\alpha}{dt}, \varphi' = \frac{d\varphi}{dt} \text{ and } \gamma' = \frac{d\gamma}{dt}.$$

We state our results as follows.

THEOREM 2.1. *Let $(M^n, g(t))_{t \in [0, T]}$ be a complete solution to the Ricci flow (1.7). Suppose that there exist three functions $\alpha(t)$, $\varphi(t)$ and $\gamma(t)$ satisfy conditions (C1), (C2), (C3) and (C4).*

Given $x_0 \in M^n$ and $R > 0$, let u be a positive solution of the equation (1.5) in the cube $B_{2R,T} := \{(x, t) | d(x, x_0, t) \leq 2R, 0 \leq t \leq T\}$. Assume that $|\text{Ric}(x, t)| \leq K$ for some $K > 0$ on $B_{2R,T}$. Let $h(x, t)$ be a function defined on $M^n \times [0, T]$ which is C^1 in t and C^2 in x , satisfying $|\nabla h|^2 \leq \delta_2 h$ and $\Delta h \geq -\delta_3$ on $B_{2R,T}$ for some positive constants δ_2 and δ_3 .

(1) $l \leq 1$. If $\frac{\gamma\alpha^4}{\alpha-1} \leq C_1$ for some constant C_1 , then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} \\ & \leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) + \frac{Cn^2}{R^2\gamma} + n^{\frac{3}{2}}\alpha^2 K \\ & \quad + \alpha \sqrt{n\bar{u}_1\delta_3} + \frac{n\alpha^2[\alpha(3-2l)-1]}{\alpha-1} \bar{u}_1\delta_1 + \sqrt{\frac{2-l}{2}}\alpha^{\frac{3}{2}}\sqrt{n\bar{u}_1\delta_2} + \alpha\varphi, \end{aligned}$$

where $C = C(n, C_1)$ is a constant.

If $\frac{\gamma}{\alpha-1} \leq C_2$ for some constant C_2 , then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} \\ & \leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) + \frac{Cn^2\alpha^4}{R^2\gamma} + n^{\frac{3}{2}}\alpha^2 K \\ & \quad + \alpha \sqrt{n\bar{u}_1\delta_3} + \frac{n\alpha^2[\alpha 3 - 2l - 1]}{\alpha-1} \bar{u}_1\delta_1 + \sqrt{\frac{2-l}{2}}\alpha^{\frac{3}{2}}\sqrt{n\bar{u}_1\delta_2} + \alpha\varphi, \end{aligned}$$

where $C = C(n, C_2)$ is a constant and set

$$\bar{u}_1 := \max_{B_{2R,T}} u^{l-1}, \quad \delta_1 := \max_{B_{2R,T}} h(x, t).$$

(2) $l > 1$. If $\frac{\gamma\alpha^4}{\alpha-1} \leq C_1$ for some constant C_1 , then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} \\ & \leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) + \frac{Cn^2}{R^2\gamma} + n^{\frac{3}{2}}\alpha^2 K + n\alpha^2(l-1)\delta_1\bar{u}_2 \\ & \quad + \alpha \sqrt{\frac{n(l\alpha-1)\bar{u}_2\delta_2}{l-1}} + \alpha^{\frac{3}{2}}\sqrt{n(l-1)\delta_1\varphi} + \alpha^{\frac{3}{2}}\sqrt{n\delta_3\bar{u}_2} + \alpha\varphi, \end{aligned}$$

where $C = C(n, C_1)$ is a constant.

If $\frac{\gamma}{\alpha-1} \leq C_2$ for some constant C_2 , then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} \\ & \leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) + \frac{Cn^2\alpha^4}{R^2\gamma} + n^{\frac{3}{2}}\alpha^2 K + n\alpha^2(l-1)\delta_1\bar{u}_2 \\ & \quad + \alpha \sqrt{\frac{n(l\alpha-1)\bar{u}_2\delta_2}{l-1}} + \alpha^{\frac{3}{2}}\sqrt{n(l-1)\delta_1\varphi} + \alpha^{\frac{3}{2}}\sqrt{n\delta_3\bar{u}_2} + \alpha\varphi, \end{aligned}$$

where $C = C(n, C_2)$ is a constant and set

$$\bar{u}_2 := \max_{B_{2R,T}} u^{l-1}, \quad \delta_1 := \max_{B_{2R,T}} h(x,t).$$

Let us list some examples to illustrate that Theorem 2.1 holds for special cases and see appendix in Section 6 for detailed calculation process.

COROLLARY 2.1. Suppose that $(M^n, g(t))_{t \in [0, T]}$ satisfies the hypotheses of Theorem 2.1. Then the following special estimates are valid.

1. Li-Yau type:

$$\alpha(t) = \text{constant}, \quad \varphi(t) = \frac{\alpha n}{t} + \frac{nK\alpha^2}{\alpha - 1}, \quad \gamma(t) = t^\theta \quad \text{with } 0 < \theta \leq 2.$$

If $l \leq 1$, then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x,t) u^{l-1} \\ & \leq C\alpha^2 \left[\frac{1}{R^2} (1 + \sqrt{KR}) + \frac{\alpha^2}{\alpha - 1} \frac{1}{R^2} + K \right] + \alpha \varphi \\ & \quad + n^{\frac{3}{2}} \alpha^2 K + \alpha \sqrt{n\bar{u}_1 \delta_3} + \frac{n\alpha^2 [\alpha(3 - 2l) - 1]}{\alpha - 1} \bar{u}_1 \delta_1 + \sqrt{\frac{2-l}{2}} \alpha^{\frac{3}{2}} \sqrt{n\bar{u}_1 \delta_2}. \end{aligned}$$

If $l > 1$, then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x,t) u^{l-1} \\ & \leq C\alpha^2 \left[\frac{1}{R^2} (1 + \sqrt{KR}) + \frac{\alpha^2}{\alpha - 1} \frac{1}{R^2} + K \right] + \alpha \varphi \\ & \quad + n^{\frac{3}{2}} \alpha^2 K + n\alpha^2(l-1)\delta_1\bar{u}_2 + \alpha \sqrt{\frac{n(l\alpha-1)\bar{u}_2\delta_2}{l-1}} \\ & \quad + \alpha^{\frac{3}{2}} \sqrt{n(l-1)\delta_1\varphi} + \alpha^{\frac{3}{2}} \sqrt{n\delta_3\bar{u}_2}. \end{aligned}$$

2. Hamilton type:

$$\alpha(t) = e^{2Kt}, \quad \varphi(t) = \frac{n}{t} e^{4Kt}, \quad \gamma(t) = t e^{2Kt}.$$

If $l \leq 1$, then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x,t) u^{l-1} \\ & \leq C\alpha^2 \left[\frac{1}{R^2} (1 + \sqrt{KR}) + K \right] + \frac{C\alpha^4}{R^2 t e^{2Kt}} + \alpha \varphi \end{aligned}$$

$$+ n^{\frac{3}{2}} \alpha^2 K + \alpha \sqrt{n\bar{u}_1 \delta_3} + \frac{n\alpha^2[\alpha(3-2l)-1]}{\alpha-1} \bar{u}_1 \delta_1 + \sqrt{\frac{2-l}{2}} \alpha \sqrt{n\bar{u}_1 \delta_2}.$$

If $l > 1$, then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} \\ & \leq C \alpha^2 \left[\frac{1}{R^2} (1 + \sqrt{K}R) + K \right] + \frac{C \alpha^4}{R^2 t e^{2Kt}} + \alpha \varphi \\ & \quad + n^{\frac{3}{2}} \alpha^2 K + n\alpha^2(l-1)\delta_1 \bar{u}_2 + \alpha \sqrt{\frac{n(l\alpha-1)\bar{u}_2 \delta_2}{l-1}} \\ & \quad + \alpha^{\frac{3}{2}} \sqrt{n(l-1)\delta_1 \varphi} + \alpha^{\frac{3}{2}} \sqrt{n\delta_3 \bar{u}_2}. \end{aligned}$$

3. Li-Xu type:

$$\begin{aligned} \alpha(t) &= 1 + \frac{\sinh(Kt) \cosh(Kt) - Kt}{\sinh^2(Kt)}, \quad \varphi(t) = 2nK[1 + \coth(Kt)], \\ \gamma(t) &= \tanh(Kt). \end{aligned}$$

If $l \leq 1$, then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} \\ & \leq C \left[\frac{1}{R^2} (1 + \sqrt{K}R) + K \right] + \frac{C}{R^2 \tanh(Kt)} + \alpha \varphi \\ & \quad + n^{\frac{3}{2}} \alpha^2 K + \alpha \sqrt{n\bar{u}_1 \delta_3} + \frac{n\alpha^2[\alpha(3-2l)-1]}{\alpha-1} \bar{u}_1 \delta_1 + \sqrt{\frac{2-l}{2}} \alpha \sqrt{n\bar{u}_1 \delta_2}. \end{aligned}$$

If $l > 1$, then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} \\ & \leq C \alpha^2 \left[\frac{1}{R^2} (1 + \sqrt{K}R) + K \right] + \frac{C}{R^2 \tanh(Kt)} + \alpha \varphi \\ & \quad + n^{\frac{3}{2}} \alpha^2 K + n\alpha^2(l-1)\delta_1 \bar{u}_2 + \alpha \sqrt{\frac{n(l\alpha-1)\bar{u}_2 \delta_2}{l-1}} \\ & \quad + \alpha^{\frac{3}{2}} \sqrt{n(l-1)\delta_1 \varphi} + \alpha^{\frac{3}{2}} \sqrt{n\delta_3 \bar{u}_2}, \end{aligned}$$

where $\alpha(t)$ is bounded uniformly.

4. Linear Li-Xu type:

$$\alpha(t) = 1 + 2Kt, \varphi(t) = \frac{n}{t} + nK(1 + 2Kt + \mu Kt), \gamma(t) = Kt \quad \text{with} \quad \mu \geq \frac{1}{4}.$$

If $l \leq 1$, then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} \\ & \leq C\alpha^2 \left[\frac{1}{R^2} (1 + \sqrt{KR}) + K \right] + \frac{C\alpha^4}{R^2 K t} + \alpha \varphi \\ & \quad + n^{\frac{3}{2}} \alpha^2 K + \alpha \sqrt{n \bar{u}_1 \delta_3} + \frac{n \alpha^2 [\alpha(3-2l)-1]}{\alpha-1} \bar{u}_1 \delta_1 + \sqrt{\frac{2-l}{2}} \alpha \sqrt{n \bar{u}_1 \delta_2}. \end{aligned}$$

If $l > 1$, then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} \\ & \leq C\alpha^2 \left[\frac{1}{R^2} (1 + \sqrt{KR}) + K \right] + \frac{C\alpha^4}{R^2 K t} + \alpha \varphi \\ & \quad + n^{\frac{3}{2}} \alpha^2 K + n \alpha^2 (l-1) \delta_1 \bar{u}_2 + \alpha \sqrt{\frac{n(l\alpha-1)\bar{u}_2\delta_2}{l-1}} \\ & \quad + \alpha^{\frac{3}{2}} \sqrt{n(l-1)\delta_1\varphi} + \alpha^{\frac{3}{2}} \sqrt{n\delta_3\bar{u}_2}. \end{aligned}$$

REMARK 2.1. The above results can be regard as some generalizations of the cases of Li-Yau [14], J. Y. Li [15], Hamilton [8] and Li-Xu [17] to the Ricci flow. Our results also generalize the estimates of S. P. Liu [18] and J. Sun [26] to the nonlinear parabolic equation along the Ricci flow. If $l < 0$ and c is a constant, there is an analog of (1.5) (see [31]), namely

$$\Delta u + cu^l = 0.$$

So our result also generalize Yang's result in [31].

The local estimates in Theorem 2.1 imply global estimates.

COROLLARY 2.2. Let $(M^n, g(t))_{t \in [0, T]}$ be a complete solution to the Ricci flow (1.7). Assume that $|\text{Ric}(x, t)| \leq K$ for some $K > 0$ and all $(x, t) \in M^n \times [0, T]$. Let $u(x, t)$ be a positive solution to equation (1.5) on $M^n \times [0, T]$. Let $h(x, t)$ be a function defined on $M^n \times [0, T]$ which is C^1 in t and C^2 in x , satisfying $|\nabla h|^2 \leq \delta_2 h$ and $\Delta h \geq -\delta_3$ on $M^n \times [0, T]$ for some positive constants δ_2 and δ_3 .

If $l \leq 1$ and for $(x, t) \in M^n \times (0, T]$, then

$$\begin{aligned} & \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} \\ & \leq \alpha \varphi + C\alpha \left[\alpha K + \sqrt{\bar{u}_1 \delta_3} + \frac{\alpha[\alpha(3-2l)-1]}{\alpha-1} \bar{u}_1 \delta_1 + \sqrt{\frac{2-l}{2}} \sqrt{\bar{u}_1 \delta_2} \right], \end{aligned}$$

where C is a positive constant depending only on n and set

$$\bar{u}_1 := \max_{M^n \times [0, T]} u^{l-1}, \quad \delta_1 := \max_{M^n \times [0, T]} h(x, t).$$

If $l > 1$ and for $(x, t) \in M^n \times (0, T]$, then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha h(x, t) u^{l-1} &\leq \alpha \varphi \\ + C\alpha \left[\alpha K + (l-1)\alpha \bar{u}_2 \delta_1 + \sqrt{\frac{(l\alpha-1)\bar{u}_2 \delta_2}{l-1}} + \alpha^{\frac{1}{2}} \sqrt{(l-1)\delta_1 \varphi} + \alpha^{\frac{1}{2}} \sqrt{\bar{u}_2 \delta_3} \right], \end{aligned}$$

where C is a positive constant depending only on n and set

$$\bar{u}_1 := \max_{M^n \times [0, T]} u^{l-1}, \quad \delta_1 := \max_{M^n \times [0, T]} h(x, t).$$

We can derive a gradient estimate for an any positive solution to the following nonlinear parabolic equation along the Ricci flow on a closed manifold without any curvature conditions. The method of the proof is inspired by Hamilton [10], Shi [24] and Liu [18].

THEOREM 2.2. *Let $(M^n, g(x, t))_{t \in [0, T]}$ be a solution to the Ricci flow (1.7) on a closed manifold. If u is a positive solution to equation*

$$\partial_t u = \Delta u + h(t) u^l,$$

where $h(t)$ is a C^1 function and $h(t) \leq 0$. Then for $l \geq 1$, we have

$$|\nabla u(x, t)|^2 \leq \frac{1}{2t} \left(\max_{x \in M^n} u^2(x, 0) - u^2(x, t) \right) \quad \text{for } (x, t) \in M^n \times [0, T]. \quad (2.1)$$

2.2. Auxilliary lemma

To prove main results, we need a lemma.

Let $f = \log u$. Then

$$f_t = \Delta f + |\nabla f|^2 + h u^{l-1}. \quad (2.2)$$

Let $F = |\nabla f|^2 - \alpha f_t + \alpha h u^{l-1} - \alpha \varphi$, where $\alpha = \alpha(t) > 1$ and $\varphi = \varphi(t) > 0$.

LEMMA 2.1. *Suppose that $(M^n, g(t))_{t \in [0, T]}$ satisfies the hypotheses of Theorem 2.1. We also assume that $\alpha(t) > 1$ and $\varphi(t) > 0$ satisfy the following system*

$$\left\{ \begin{array}{l} \frac{2\varphi}{n} - 2\alpha K \geq (\frac{2\varphi}{n} - \alpha') \frac{1}{\alpha}, \\ \frac{2\varphi}{n} - \alpha' > 0, \\ \frac{\varphi^2}{n} + \alpha \varphi' \geq 0, \end{array} \right. \quad (2.3)$$

and $\alpha(t)$ is non-decreasing. Then

$$(\Delta - \partial_t)F \geq |f_{ij} + \frac{\varphi}{n} \delta_{ij}|^2 + (\frac{2\varphi}{n} - \alpha') \frac{1}{\alpha} F - \alpha^2 n^2 K^2 - 2\nabla f \nabla F$$

$$\begin{aligned}
& + 2h(\alpha - 1)(l-1)u^{l-1}|\nabla f|^2 + \alpha(l-1)^2hu^{l-1}|\nabla f|^2 \\
& + 2[(\alpha - 1) + \alpha(l-1)]u^{l-1}\nabla f \cdot \nabla h \\
& + \alpha(l-1)hu^{l-1}\Delta f + \alpha u^{l-1}\Delta h.
\end{aligned} \tag{2.4}$$

Proof. By directly computing, we have

$$\begin{aligned}
\Delta F &= \Delta|\nabla f|^2 - \alpha\Delta(f_t) + \alpha\Delta(hu^{l-1}) \\
&= 2|f_{ij}|^2 + 2f_j f_{iij} + 2R_{ij} f_i f_j - \alpha\Delta(f_t) + \alpha h\Delta(u^{l-1}) \\
&\quad + \alpha u^{l-1}\Delta h + 2\alpha\nabla h\nabla u^{l-1} \\
&= 2\left(|f_{ij}|^2 + \alpha R_{ij} f_{ij}\right) + 2f_j f_{iij} + 2R_{ij} f_i f_j - \alpha(\Delta f)_t \\
&\quad + \alpha h\Delta(u^{l-1}) + \alpha u^{l-1}\Delta h + 2\alpha\nabla h\nabla u^{l-1},
\end{aligned}$$

where we have used Bochner's formula and

$$\Delta(f_t) = (\Delta f)_t - 2R_{ij} f_{ij}.$$

Applying Young's inequality

$$R_{ij} f_{ij} \leq |R_{ij} f_{ij}| \leq \frac{\alpha}{2} |R_{ij}|^2 + \frac{1}{2\alpha} |f_{ij}|^2,$$

we conclude for $|R_{ij}| \leq K$,

$$\begin{aligned}
\Delta F &\geq |f_{ij}|^2 - \alpha^2|R_{ij}|^2 + 2f_j f_{iij} + 2R_{ij} f_i f_j - \alpha(\Delta f)_t \\
&\quad + \alpha h\Delta(u^{l-1}) + \alpha u^{l-1}\Delta h + 2\alpha\nabla h\nabla u^{l-1} \\
&\geq |f_{ij}|^2 - \alpha^2 n^2 K^2 + 2f_j f_{iij} + 2R_{ij} f_i f_j - \alpha(\Delta f)_t \\
&\quad + \alpha h\Delta(u^{l-1}) + \alpha u^{l-1}\Delta h + 2\alpha\nabla h\nabla u^{l-1}.
\end{aligned} \tag{2.5}$$

On the other hand, we infer

$$\begin{aligned}
\partial_t F &= (|\nabla f|^2)_t - \alpha f_{tt} - \alpha' f_t + \alpha' h u^{l-1} + \alpha h (\Delta u^{l-1})_t \\
&\quad + \alpha u^{l-1} h_t - \alpha \varphi' - \alpha' \varphi \\
&= 2\nabla f \nabla(f_t) + 2R_{ij} f_i f_j - \alpha f_{tt} - \alpha' f_t + \alpha' h u^{l-1} + \alpha u^{l-1} h_t \\
&\quad + \alpha h (\Delta u^{l-1})_t - \alpha \varphi' - \alpha' \varphi,
\end{aligned} \tag{2.6}$$

where we used the fact that

$$(|\nabla f|^2)_t = 2\nabla f \cdot \nabla(f_t) + 2\text{Ric}(\nabla f, \nabla f). \tag{2.7}$$

We follow from (2.5) and (2.6),

$$\begin{aligned}
(\Delta - \partial_t)F &\geq |f_{ij}|^2 - \alpha^2 n^2 K^2 + 2\nabla f \nabla(\Delta f) - \alpha(\Delta f)_t + \alpha h\Delta(u^{l-1}) \\
&\quad + \alpha u^{l-1}\Delta h + 2\alpha\nabla h\nabla u^{l-1} - 2\nabla f \nabla(f_t) + \alpha f_{tt} + \alpha' f_t
\end{aligned}$$

$$\begin{aligned}
& -\alpha' h u^{l-1} - \alpha h (u^{l-1})_t - \alpha u^{l-1} h_t + \alpha \varphi' + \alpha' \varphi \\
&= |f_{ij}|^2 - \alpha^2 n^2 K^2 + 2\nabla f \nabla (\Delta f) - \alpha (f_t - |\nabla f|^2 - h u^{l-1})_t \\
&\quad + \alpha h \Delta (u^{l-1}) + \alpha u^{l-1} \Delta h + 2\alpha \nabla h \nabla u^{l-1} - 2\nabla f \nabla (f_t) + \alpha f_{it} \\
&\quad + \alpha' f_t - \alpha' h u^{l-1} - \alpha h (u^{l-1})_t - \alpha u^{l-1} h_t + \alpha \varphi' + \alpha' \varphi \\
&= |f_{ij}|^2 - \alpha^2 n^2 K^2 + 2\nabla f \nabla (\Delta f) + \alpha (|\nabla f|^2)_t + \alpha h \Delta (u^{l-1}) \\
&\quad + \alpha u^{l-1} \Delta h + 2\alpha \nabla h \nabla u^{l-1} - 2\nabla f \nabla (f_t) + \alpha' f_t \\
&\quad - \alpha' h u^{l-1} + \alpha \varphi' + \alpha' \varphi.
\end{aligned}$$

By using the formula (2.7) and (2.2), we have

$$\begin{aligned}
(\Delta - \partial_t)F &\geq |f_{ij}|^2 - \alpha^2 n^2 K^2 + 2\nabla f \nabla (\Delta f) + 2\alpha \nabla f \nabla (f_t) \\
&\quad + 2\alpha R_{ij} f_i f_j + \alpha h \Delta (u^{l-1}) + \alpha u^{l-1} \Delta h + 2\alpha \nabla h \nabla u^{l-1} \\
&\quad - 2\nabla f \nabla (f_t) + \alpha' f_t - \alpha' h u^{l-1} + \alpha \varphi' + \alpha' \varphi \\
&= |f_{ij}|^2 + 2\alpha R_{ij} f_i f_j - \alpha^2 n^2 K^2 - 2\nabla f \nabla F \\
&\quad + 2(\alpha - 1) \nabla f \nabla (h u^{l-1}) + \alpha h \Delta (u^{l-1}) + \alpha u^{l-1} \Delta h \\
&\quad + 2\alpha \nabla h \nabla u^{l-1} + \alpha' f_t - \alpha' h u^{l-1} + \alpha \varphi' + \alpha' \varphi. \tag{2.8}
\end{aligned}$$

Applying the following two equations

$$\begin{aligned}
\nabla(u^{l-1}) &= (l-1)u^{l-1}\nabla f, \\
\Delta(u^{l-1}) &= (l-1)^2 u^{l-1} |\nabla f|^2 + (l-1)u^{l-1} \Delta f,
\end{aligned}$$

to (2.8), we have

$$\begin{aligned}
(\Delta - \partial_t)F &\geq |f_{ij}|^2 + 2\alpha R_{ij} f_i f_j - \alpha^2 n^2 K^2 + 2\nabla f \nabla F + \alpha u^{l-1} \Delta h \\
&\quad + 2h(\alpha - 1)(l-1)u^{l-1} |\nabla f|^2 + 2[(\alpha - 1) + \alpha(l-1)]u^{l-1} \nabla f \cdot \nabla h \\
&\quad + h\alpha(l-1)^2 u^{l-1} |\nabla f|^2 + h\alpha(l-1)u^{l-1} \Delta f \\
&\quad + \alpha' f_t - \alpha' h u^{l-1} + \alpha \varphi' + \alpha' \varphi. \tag{2.9}
\end{aligned}$$

Further applying unit matrix $(\delta_{ij})_{n \times n}$ and (2.9), we derive

$$\begin{aligned}
(\Delta - \partial_t)F &\geq |f_{ij}|^2 + \frac{\varphi}{n} \delta_{ij} - 2\alpha K |\nabla f|^2 - \alpha^2 n^2 K^2 + 2\nabla f \nabla F \\
&\quad + 2h(\alpha - 1)(l-1)u^{l-1} |\nabla f|^2 + 2[(\alpha - 1) + \alpha(l-1)]u^{l-1} \nabla f \cdot \nabla h \\
&\quad + h\alpha(l-1)^2 u^{l-1} |\nabla f|^2 + h\alpha(l-1)u^{l-1} \Delta f + \alpha u^{l-1} \Delta h \\
&\quad + \alpha' f_t - \alpha' h u^{l-1} + \alpha \varphi' + \alpha' \varphi - \frac{\varphi^2}{n} - 2\frac{\varphi}{n} \Delta f.
\end{aligned}$$

Applying (2.2), we have

$$(\Delta - \partial_t)F \geq |f_{ij}|^2 + \left(\frac{2\varphi}{n} - 2\alpha K\right) |\nabla f|^2 - \left(\frac{2\varphi}{n} - \alpha'\right) f_t$$

$$\begin{aligned}
& + \left(\frac{2\varphi}{n} - \alpha' \right) h u^{l-1} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{\alpha\varphi}{\alpha} - \alpha^2 n^2 K^2 - 2\nabla f \nabla F \\
& + 2h(\alpha-1)(l-1)u^{l-1}|\nabla f|^2 + 2[(\alpha-1)+\alpha(l-1)]u^{l-1}\nabla f \cdot \nabla h \\
& + h\alpha(l-1)^2u^{l-1}|\nabla f|^2 + h\alpha(l-1)u^{l-1}\Delta f + \alpha u^{l-1}\Delta h \\
& + \alpha\varphi' + \alpha'\varphi - \frac{\varphi^2}{n} + \left(\frac{2\varphi}{n} - \alpha' \right) \frac{\alpha\varphi}{\alpha}.
\end{aligned} \tag{2.10}$$

Therefore, (2.4) is derived from (2.3) and (2.10). The proof is complete.

2.3. Proof of Theorem 2.1 and 2.2

In this section, we will prove the Theorems 2.1 and 2.2.

Proof of Theorem 2.1. Let $G = \gamma(t)F$ and $\gamma(t) > 0$ be non-decreasing. Then

$$\begin{aligned}
(\Delta - \partial_t)G &= \gamma(\Delta - \partial_t)F - \gamma'F \\
&\geq \gamma|f_{ij} + \frac{\varphi}{n}\delta_{ij}|^2 + \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha}G - \gamma\alpha^2 n^2 K^2 - 2\nabla f \nabla G \\
&\quad + 2h\gamma(\alpha-1)(l-1)u^{l-1}|\nabla f|^2 + 2[(\alpha-1)+\alpha(l-1)]\gamma u^{l-1}\nabla f \cdot \nabla h \\
&\quad + h\gamma\alpha(l-1)^2u^{l-1}|\nabla f|^2 + h\gamma\alpha(l-1)u^{l-1}\Delta f + \alpha\gamma u^{l-1}\Delta h - \gamma'F \\
&= \gamma|f_{ij} + \frac{\varphi}{n}g_{ij}|^2 + \left[\left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma} \right] G - \gamma\alpha^2 n^2 K^2 - 2\nabla f \nabla G \\
&\quad + 2h\gamma(\alpha-1)(l-1)u^{l-1}|\nabla f|^2 + 2[(\alpha-1)+\alpha(l-1)]\gamma u^{l-1}\nabla f \cdot \nabla h \\
&\quad + \gamma\alpha(l-1)^2hu^{l-1}|\nabla f|^2 + \gamma\alpha(l-1)hu^{l-1}\Delta f + \alpha\gamma u^{l-1}\Delta h. \tag{2.11}
\end{aligned}$$

Now let $\varphi(r)$ be a C^2 function on $[0, \infty)$ such that

$$\varphi(r) = \begin{cases} 1, & \text{if } r \in [0, 1], \\ 0, & \text{if } r \in [2, \infty), \end{cases}$$

and

$$0 \leq \varphi(r) \leq 1, \quad \varphi'(r) \leq 0, \quad \varphi''(r) \leq 0, \quad \frac{|\varphi'(r)|}{\varphi(r)} \leq C,$$

where C is an absolute constant. Let define by

$$\phi(x, t) = \varphi(d(x, x_0, t)) = \varphi\left(\frac{d(x, x_0, t)}{R}\right) = \varphi\left(\frac{\rho(x, t)}{R}\right),$$

where $\rho(x, t) = d(x, x_0, t)$. By using the maximum principle, the argument of Calabi [2] allows us to suppose that the function $\phi(x, t)$ with support in $B_{2R, T}$, is C^2 at the maximum point. By utilizing the Laplacian comparison theorem, we deduce that

$$\frac{|\nabla\phi|^2}{\phi} \leq \frac{C}{R^2}, \quad -\Delta\phi \leq \frac{C}{R^2}(1 + \sqrt{KR}), \tag{2.12}$$

For any $0 \leq T_1 \leq T$, let $H = \phi G$ and (x_1, t_1) be the point in B_{2R, T_1} at which H attains its maximum value. We can suppose that the value is positive, because otherwise the proof is trivial. Then at the point (x_1, t_1) , we infer

$$\left. \begin{aligned} 0 &= \nabla(\phi G) = G\nabla\phi + \phi\nabla G, \\ \Delta(\phi G) &\leq 0, \\ \partial_t(\phi G) &\geq 0. \end{aligned} \right\} \quad (2.13)$$

By the evolution formula of the geodesic length along the Ricci flow [6], we calculate

$$\begin{aligned} \phi_t G &= -G\phi' \left(\frac{\rho}{R} \right) \frac{1}{R} \frac{d\rho}{dt} = G\phi' \left(\frac{\rho}{R} \right) \int_{\gamma_1} \text{Ric}(S, S) ds \\ &\leq G\phi' \left(\frac{\rho}{R} \right) \frac{1}{R} K\rho \leq G\phi' \left(\frac{\rho}{R} \right) K_2 \leq G\sqrt{C}K, \end{aligned}$$

where γ_1 is the geodesic connecting x and x_0 along the metric $g(t_1)$, S is the unite tangent vector to γ_1 , and ds is the element of the arc length.

All the following computations are at the point (x_1, t_1) . It is not difficult to find that

$$\begin{aligned} |f_{ij} + \frac{\varphi}{n}\delta_{ij}|^2 &\geq \frac{1}{n} \left(\text{tr}|f_{ij} + \frac{\varphi}{n}\delta_{ij}| \right)^2 \\ &= \frac{1}{n} (\Delta f + \varphi)^2 \\ &= \frac{1}{n} \left[-\frac{1}{\alpha} F - \frac{1}{\alpha} (\alpha - 1) |\nabla f|^2 \right]^2 \\ &= \frac{1}{\alpha^2 n} \left[\frac{G}{\gamma} + (\alpha - 1) |\nabla f|^2 \right]^2, \end{aligned} \quad (2.14)$$

and

$$\begin{aligned} \Delta f &= f_t - |\nabla f|^2 - hu^{l-1} \\ &= -\frac{F}{\alpha} - \frac{\alpha - 1}{\alpha} |\nabla f|^2 - \varphi < 0. \end{aligned} \quad (2.15)$$

To obtain main results, two cases will be shown.

Case 1 $l \leq 1$.

From (2.15), we have $\Delta f \leq 0$. Then by substituting it into (2.11), we obtain

$$\begin{aligned} (\Delta - \partial_t)G &= \gamma(\Delta - \partial_t)F - \gamma'F \\ &\geq \gamma|f_{ij} + \frac{\varphi}{n}\delta_{ij}|^2 + \left[\left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma} \right] G - \gamma\alpha^2 n^2 K^2 - 2\nabla f \nabla G \\ &\quad + 2h\gamma(\alpha - 1)(l - 1)u^{l-1}|\nabla f|^2 + \alpha\gamma u^{l-1}\Delta h \\ &\quad + 2[(\alpha - 1) + \alpha(l - 1)]\gamma u^{l-1}\nabla f \cdot \nabla h, \end{aligned}$$

where we drop one term $h\gamma\alpha(l-1)^2u^{l-1}|\nabla f|^2$. Using (2.14), we infer

$$\begin{aligned}
 0 &\geq (\Delta - \partial_t)(\phi G) \\
 &= G\left(\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi}\right) + \phi(\Delta - \partial_t)G - G\phi_t \\
 &\geq G\left(\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi}\right) + \frac{\phi\gamma}{\alpha^2 n} \left[\frac{G}{\gamma} + (\alpha-1)|\nabla f|^2\right]^2 \\
 &\quad + \left[(\frac{2\phi}{n} - \alpha')\frac{1}{\alpha} - \frac{\gamma'}{\gamma}\right]\phi G - \gamma\phi\alpha^2 n^2 K^2 - 2\phi\nabla f\nabla G \\
 &\quad + 2h\phi\gamma(\alpha-1)(l-1)u^{l-1}|\nabla f|^2 + \phi\alpha\gamma u^{l-1}\Delta h \\
 &\quad + 2[(\alpha-1) + \alpha(l-1)]\phi\gamma u^{l-1}\nabla f \cdot \nabla h - G\sqrt{C}K. \tag{2.16}
 \end{aligned}$$

Multiply ϕ to inequality (2.16), we have

$$\begin{aligned}
 0 &\geq \phi G \left[\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi} + \left(\frac{2\phi}{n} - \alpha'\right)\frac{\phi}{\alpha} - \frac{\gamma'}{\gamma}\phi \right] + \frac{\phi^2\gamma}{\alpha^2 n} \left[\frac{G}{\gamma} + (\alpha-1)|\nabla f|^2\right]^2 \\
 &\quad - \gamma\phi^2\alpha^2 n^2 K^2 - 2\phi^2\nabla f\nabla G + 2h\phi^2\gamma(\alpha-1)(l-1)u^{l-1}|\nabla f|^2 \\
 &\quad + 2[(\alpha-1) + \alpha(l-1)]\phi^2\gamma u^{l-1}\nabla f \cdot \nabla h + \phi^2\alpha\gamma u^{l-1}\Delta h - \phi G\sqrt{C}K \\
 &= \phi G \left[\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi} + \left(\frac{2\phi}{n} - \alpha'\right)\frac{1}{\alpha} - \frac{\gamma'}{\gamma} \right] + \frac{\phi^2 G^2}{\alpha^2 n \gamma} + \frac{\phi^2(\alpha-1)^2\gamma}{n\alpha^2} |\nabla f|^4 \\
 &\quad + \frac{2\phi^2(\alpha-1)}{n\alpha^2} G|\nabla f|^2 - \gamma\phi^2\alpha^2 n^2 K^2 + 2\phi G\nabla\phi\nabla f \\
 &\quad + 2h\phi^2\gamma(\alpha-1)(l-1)u^{l-1}|\nabla f|^2 + \phi^2\alpha\gamma u^{l-1}\Delta h \\
 &\quad + 2[(\alpha-1) + \alpha(l-1)]\phi^2\gamma u^{l-1}\nabla f \cdot \nabla h - \phi G\sqrt{C}K.
 \end{aligned}$$

Using the Cauchy inequality

$$\nabla f \cdot \nabla h \geq -|\nabla f||\nabla h| \geq -h|\nabla f|^2 - \frac{|\nabla h|^2}{4h},$$

$$\nabla f \cdot \nabla h \leq h|\nabla f|^2 + \frac{|\nabla h|^2}{4h},$$

we conclude

$$\begin{aligned}
 0 &\geq \phi G \left[\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi} + \left(\frac{2\phi}{n} - \alpha'\right)\frac{\phi}{\alpha} - \frac{\gamma'}{\gamma}\phi \right] + \frac{\phi^2 G^2}{\alpha^2 n \gamma} + \frac{\phi^2(\alpha-1)^2\gamma}{n\alpha^2} |\nabla f|^4 \\
 &\quad + \frac{2\phi^2(\alpha-1)}{n\alpha^2} G|\nabla f|^2 - \gamma\phi^2\alpha^2 n^2 K^2 + 2\phi G\nabla\phi\nabla f \\
 &\quad - 2h\phi^2\gamma(\alpha-1)(l-1)u^{l-1}|\nabla f|^2 - 2[(\alpha-1) + \alpha(1-l)]\phi^2\gamma u^{l-1}h|\nabla f|^2 \\
 &\quad - \frac{1}{2}[(\alpha-1) + \alpha(1-l)]\phi^2\gamma u^{l-1}\frac{|\nabla h|^2}{h} + \phi^2\alpha\gamma u^{l-1}\Delta h - \phi G\sqrt{C}K
 \end{aligned}$$

$$\begin{aligned}
&\geq \phi G \left[\Delta \phi - 2 \frac{|\nabla \phi|^2}{\phi} + \left(\frac{2\varphi}{n} - \alpha' \right) \frac{\phi}{\alpha} - \frac{\gamma'}{\gamma} \phi \right] + \frac{\phi^2 G^2}{\alpha^2 n \gamma} + \frac{\phi^2 (\alpha - 1)^2 \gamma}{n \alpha^2} |\nabla f|^4 \\
&\quad + \frac{2\phi^2 (\alpha - 1)}{n \alpha^2} G |\nabla f|^2 - \gamma \phi^2 \alpha^2 n^2 K^2 + 2\phi G \nabla \phi \nabla f \\
&\quad - 2[\alpha(3 - 2\alpha) - 1] \phi^2 \gamma u^{l-1} h |\nabla f|^2 - \frac{1}{2} [(\alpha - 1) + \alpha(1 - l)] \phi^2 \gamma u^{l-1} \frac{|\nabla h|^2}{h} \\
&\quad + \phi^2 \alpha \gamma u^{l-1} \Delta h - \phi G \sqrt{C} K,
\end{aligned} \tag{2.17}$$

where we used the fact that $(\alpha - 1)(1 - l) + (\alpha - 1) + \alpha(1 - l) \leq \alpha(3 - 2l) - 1$. Further using the inequality $Ax^2 + Bx \geq -\frac{B^2}{4A}$ with $A > 0$, we have

$$\frac{2\phi^2 (\alpha - 1)}{n \alpha^2} G |\nabla f|^2 + 2\phi G \nabla \phi \nabla f \geq -\frac{n \alpha^2}{2(\alpha - 1)} \frac{|\nabla \phi|^2}{\phi} \phi G,$$

and

$$\begin{aligned}
&\frac{\phi^2 (\alpha - 1)^2 \gamma}{n \alpha^2} |\nabla f|^4 - 2[\alpha(3 - 2l) - 1] \phi^2 \gamma u^{l-1} h |\nabla f|^2 \\
&\geq -\frac{n \alpha^2 [\alpha(3 - 2l) - 1]^2}{(\alpha - 1)^2} \gamma \phi^2 u^{2(l-1)} h^2 \\
&\geq -\frac{n \alpha^2 [\alpha(3 - 2\alpha) - 1]^2}{(\alpha - 1)^2} \gamma \phi^2 \bar{u}_1^2 \delta_1^2.
\end{aligned}$$

Substituting above two inequalities into (2.17), we deduce that

$$\begin{aligned}
0 &\geq \phi G \left[\Delta \phi - 2 \frac{|\nabla \phi|^2}{\phi} + \left(\frac{2\varphi}{n} - \alpha' \right) \frac{\phi}{\alpha} - \frac{\gamma'}{\gamma} \phi - \frac{n \alpha^2}{2(\alpha - 1)} \frac{|\nabla \phi|^2}{\phi} - \sqrt{C} K \right] \\
&\quad + \frac{\phi^2 G^2}{\alpha^2 n \gamma} - \gamma \phi^2 \alpha^2 n^2 K^2 - \frac{n \alpha^2 [\alpha(3 - 2l) - 1]^2}{(\alpha - 1)^2} \gamma \phi^2 \bar{u}_1^2 \delta_1^2 \\
&\quad - \frac{1}{2} [(\alpha - 1) + \alpha(1 - l)] \phi^2 \gamma \bar{u}_1 \delta_2 - \phi^2 \alpha \gamma \bar{u}_1 \delta_3.
\end{aligned}$$

Applying (2.12), we infer

$$\begin{aligned}
0 &\geq \left[-\frac{C}{R^2} (1 + \sqrt{k} R) - \frac{2C}{R^2} + \left(\frac{2\varphi}{n} - \alpha' \right) \frac{\phi}{\alpha} - \frac{\gamma'}{\gamma} \phi \right. \\
&\quad \left. - \frac{n \alpha^2}{2(\alpha - 1)} \frac{C}{R^2} - \sqrt{C} K \right] \phi G + \frac{\phi^2 G^2}{\alpha^2 n \gamma} - \gamma \phi^2 \alpha^2 n^2 K^2 \\
&\quad - \frac{n \alpha^2 [\alpha(3 - 2l) - 1]^2}{(\alpha - 1)^2} \gamma \phi^2 \bar{u}_1^2 \delta_1^2 \\
&\quad - \frac{1}{2} [(\alpha - 1) + \alpha(1 - l)] \phi^2 \gamma \bar{u}_1 \delta_2 - \phi^2 \alpha \gamma \bar{u}_1 \delta_3.
\end{aligned} \tag{2.18}$$

For the inequality $Ax^2 - Bx \leq C$, one has $x \leq \frac{B}{A} + \left(\frac{C}{A}\right)^{\frac{1}{2}}$, where $A, B, C > 0$. By using this inequality to (2.18) and then we arrive at

$$\begin{aligned} \phi G(x, T_1) &\leq (\phi G)(x_1, t_1) \\ &\leq \left\{ n\gamma\alpha^2 \left[\frac{C}{R^2} (1 + \sqrt{KR}) + \frac{n\alpha^2}{2(\alpha-1)} \frac{C}{R^2} + \sqrt{CK} \right] \right. \\ &\quad + n\gamma\alpha^2 \left[\frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \frac{\alpha'}{\alpha} \right) \frac{1}{\alpha} \right] + n^{\frac{3}{2}}\gamma\alpha^2\phi K \\ &\quad + \frac{n\alpha^2[\alpha(3-2l)-1]}{\alpha-1} \phi\bar{u}_1\delta_1 + \alpha\phi\gamma\sqrt{n\bar{u}_1\delta_3} \\ &\quad \left. + \sqrt{\frac{[(\alpha-1)+\alpha(1-l)]}{2}} \alpha\phi\gamma\sqrt{n\bar{u}_1\delta_2} \right\} (x_1, t_1). \end{aligned}$$

If γ is nondecreasing which satisfies the system

$$\begin{cases} \frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} \leq 0, \\ \frac{\gamma\alpha^4}{\alpha-1} \leq C_1. \end{cases} \quad (2.19)$$

Recall that $\alpha(t)$ and $\gamma(t)$ are non-decreasing and $t_1 < T_1$. Hence, we have

$$\begin{aligned} \phi G(x, T_1) &\leq (\phi G)(x_1, t_1) \\ &\leq n\gamma(T_1)\alpha^2(T_1) \left[\frac{C}{R^2} (1 + \sqrt{KR}) + K \right] + \frac{n^2C}{R^2} \\ &\quad + n^{\frac{3}{2}}\gamma(T_1)\alpha^2(T_1)\phi K + \phi\alpha(T_1)\gamma(T_1)\sqrt{n\bar{u}_1\delta_3} \\ &\quad + \frac{n\alpha^2(T_1)[\alpha(T_1)(3-2l)-1]}{\alpha(T_1)-1} \phi\gamma(T_1)\bar{u}_1\delta_1 \\ &\quad + \sqrt{\frac{[(\alpha(T_1)-1)+\alpha(T_1)(1-l)]}{2}} \alpha(T_1)\phi\gamma(T_1)\sqrt{n\bar{u}_1\delta_2}. \end{aligned}$$

Hence, we have for $\phi \equiv 1$ on $B_{R,T}$,

$$\begin{aligned} F(x, T_1) &\leq n\alpha^2(T_1) \left[\frac{C}{R^2} (1 + \sqrt{KR}) + CK \right] + \frac{n^2C}{R^2\gamma(T_1)} \\ &\quad + n^{\frac{3}{2}}\alpha^2(T_1)K + \alpha(T_1)\sqrt{n\bar{u}_1\delta_3} \\ &\quad + \frac{n\alpha^2(T_1)[\alpha(T_1)(3-2l)-1]}{\alpha(T_1)-1} \bar{u}_1\delta_1 \\ &\quad + \sqrt{\frac{[(\alpha(T_1)-1)+\alpha(T_1)(1-l)]}{2}} \alpha(T_1)\sqrt{n\bar{u}_1\delta_2}. \end{aligned}$$

If γ is nondecreasing which satisfies the system

$$\begin{cases} \frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \alpha'\right) \frac{1}{\alpha} \leq 0, \\ \frac{\gamma}{\alpha - 1} \leq C_2. \end{cases} \quad (2.20)$$

Recall that $\alpha(t)$ and $\gamma(t)$ are non-decreasing and $t_1 < T_1$. Hence, we have

$$\begin{aligned} \phi G(x, T_1) &\leq (\phi G)(x_1, t_1) \\ &\leq n\gamma(T_1)\alpha^2(T_1) \left[\frac{C}{R^2} \left(1 + \sqrt{KR} \right) + \frac{n\alpha^2}{\alpha - 1} \frac{C}{R^2} + CK \right] \\ &\quad + n^{\frac{3}{2}}\gamma(T_1)\alpha^2(T_1)\phi K + \phi\alpha(T_1)\gamma(T_1)\sqrt{n\bar{u}_1\delta_3} \\ &\quad + \frac{n\alpha^2(T_1)[\alpha(T_1)(3-2l)-1]}{\alpha(T_1)-1}\phi\gamma(T_1)\bar{u}_1\delta_1 \\ &\quad + \sqrt{\frac{[(\alpha(T_1)-1)+\alpha(T_1)(1-l)]}{2}}\alpha(T_1)\phi\gamma(T_1)\sqrt{n\bar{u}_1\delta_2}. \end{aligned}$$

Hence, we have for $\phi \equiv 1$ on $B_{R,T}$,

$$\begin{aligned} F(x, T_1) &\leq n\alpha^2(T_1) \left[\frac{C}{R^2} \left(1 + \sqrt{KR} \right) + K \right] + \frac{n^2C\alpha^4}{R^2\gamma(T)} \\ &\quad + n^{\frac{3}{2}}\gamma(T_1)\alpha^2(T_1)K + \alpha(T_1)\sqrt{n\bar{u}_1\delta_3} \\ &\quad + \frac{n\alpha^2(T_1)[\alpha(T_1)(3-2l)-1]}{\alpha(T_1)-1}\bar{u}_1\delta_1 \\ &\quad + \sqrt{\frac{[(\alpha(T_1)-1)+\alpha(T_1)(1-l)]}{2}}\alpha(T_1)\sqrt{n\bar{u}_1\delta_2}. \end{aligned}$$

Because T_1 is arbitrary in $0 < T_1 < T$ and $\alpha - 1 + \alpha(1-l) \leq \alpha(2-l)$. Thus the conclusion is valid.

Case 2 $l > 1$.

Substituting (2.15) into (2.11), we have

$$\begin{aligned} (\Delta - \partial_t)G &\geq \gamma|f_{ij}|^2 + \left[\left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma} \right] G - \gamma\alpha^2 n^2 K^2 \\ &\quad - 2\nabla f \nabla G + 2(l\alpha - 1)\gamma u^{l-1} \nabla f \cdot \nabla h - h(l-1)u^{l-1}G \\ &\quad - h\gamma(l-1)u^{l-1}\alpha\varphi + h\gamma(l-1)(l\alpha - 1)u^{l-1}|\nabla f|^2 + \alpha\gamma u^{l-1}\Delta h. \end{aligned}$$

Using (2.14), we infer

$$\begin{aligned} 0 &\geq (\Delta - \partial_t)(\phi G) \\ &= G \left(\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi} \right) + \phi(\Delta - \partial_t)G - \gamma G\phi_t \end{aligned}$$

$$\begin{aligned}
&\geq G \left(\Delta \phi - 2 \frac{|\nabla \phi|^2}{\phi} \right) + \frac{\phi \gamma}{\alpha^2 n} \left[\frac{G}{\gamma} + (\alpha - 1) |\nabla f|^2 \right]^2 \\
&\quad + \left[\left(\frac{2\phi}{2} - \alpha' \right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma} \right] \phi G - \gamma \phi \alpha^2 n^2 K^2 - 2 \phi \nabla f \nabla G \\
&\quad + 2(l\alpha - 1) \phi \gamma u^{l-1} \nabla f \cdot \nabla h - h(l-1) u^{l-1} G - h \gamma(l-1) u^{l-1} \phi \alpha \varphi \\
&\quad + \phi \alpha \gamma u^{l-1} \Delta h + h \phi \gamma(l-1) (l\alpha - 1) u^{l-1} |\nabla f|^2 - G \sqrt{C} K. \tag{2.21}
\end{aligned}$$

Multiply ϕ to (2.21), and we have

$$\begin{aligned}
0 &\geq \phi G \left[\Delta \phi - 2 \frac{|\nabla \phi|^2}{\phi} + \left(\frac{2\phi}{n} - \alpha' \right) \frac{\phi}{\alpha} - \frac{\gamma'}{\gamma} \phi \right] + \frac{\phi^2 \gamma}{\alpha^2 n} \left[\frac{G}{\gamma} + (\alpha - 1) |\nabla f|^2 \right]^2 \\
&\quad - \gamma \phi^2 \alpha^2 n^2 K^2 - 2 \phi^2 \nabla f \nabla G + 2(l\alpha - 1) \phi^2 \gamma u^{l-1} \nabla f \cdot \nabla h \\
&\quad + \phi^2 \alpha \gamma u^{l-1} \Delta h - h(l-1) u^{l-1} \phi G - h \gamma(l-1) u^{l-1} \phi^2 \alpha \varphi \\
&\quad + h \phi^2 \gamma(l-1) (l\alpha - 1) u^{l-1} |\nabla f|^2 - \phi G \sqrt{C} K \\
&\geq \phi G \left[\Delta \phi - 2 \frac{|\nabla \phi|^2}{\phi} + \left(\frac{2\phi}{n} - \alpha' \right) \frac{\phi}{\alpha} - \frac{\gamma'}{\gamma} \phi \right] + \frac{\phi^2 G^2}{\alpha^2 n \gamma} \\
&\quad + \frac{2\phi^2(\alpha-1)}{n\alpha^2} G |\nabla f|^2 - \gamma \phi^2 \alpha^2 n^2 K^2 + 2 \phi G \nabla \phi \nabla f \\
&\quad + 2(l\alpha - 1) \phi^2 \gamma u^{l-1} \nabla f \cdot \nabla h + \phi^2 \alpha \gamma u^{l-1} \Delta h \\
&\quad - h(l-1) u^{l-1} \phi G - h \gamma(l-1) u^{l-1} \phi^2 \alpha \varphi \\
&\quad + h \phi^2 \gamma(l-1) (l\alpha - 1) u^{l-1} |\nabla f|^2 - \phi G \sqrt{C} K, \tag{2.22}
\end{aligned}$$

where we dropped the term $\frac{\phi^2(\alpha-1)^2 \gamma}{n\alpha^2} |\nabla f|^4$.

Further using the inequality $Ax^2 + Bx \geq -\frac{B^2}{4A}$ with $A > 0$, we have

$$\frac{2\phi^2(\alpha-1)}{n\alpha^2} G |\nabla f|^2 + 2 \phi G \nabla \phi \nabla f \geq -\frac{n\alpha^2}{2(\alpha-1)} \frac{|\nabla \phi|^2}{\phi} \phi G,$$

and

$$\begin{aligned}
&h \phi^2 \gamma(l-1) (l\alpha - 1) u^{l-1} |\nabla f|^2 + 2(l\alpha - 1) \phi^2 \gamma u^{l-1} \nabla f \cdot \nabla h \\
&\geq h \phi^2 \gamma(l-1) (l\alpha - 1) u^{l-1} |\nabla f|^2 - 2(l\alpha - 1) \phi^2 \gamma u^{l-1} |\nabla f| \cdot |\nabla h| \\
&\geq -\frac{l\alpha - 1}{l-1} \gamma \phi^2 u^{l-1} \frac{|\nabla h|^2}{h} \\
&\geq -\frac{l\alpha - 1}{l-1} \gamma \phi^2 \bar{u}_2 \delta_2.
\end{aligned}$$

Substituting above two inequalities into (2.22), we deduce that

$$0 \geq \phi G \left[\Delta \phi - 2 \frac{|\nabla \phi|^2}{\phi} + \left(\frac{2\phi}{n} - \alpha' \right) \frac{\phi}{\alpha} - \frac{\gamma'}{\gamma} \phi - \frac{n\alpha^2}{2(\alpha-1)} \frac{|\nabla \phi|^2}{\phi} \right]$$

$$\begin{aligned} & -\delta_1(l-1)\bar{u}_2 - \sqrt{CK} \Big] + \frac{\phi^2 G^2}{\alpha^2 n \gamma} - \gamma \phi^2 \alpha^2 n^2 K^2 \\ & - \frac{l\alpha-1}{l-1} \gamma \phi^2 \bar{u}_2 \delta_2 - \gamma(l-1) \bar{u}_2 \phi^2 \delta_1 \alpha \varphi - \phi^2 \alpha \gamma \bar{u}_2 \delta_3. \end{aligned}$$

Applying (2.12), we infer

$$\begin{aligned} 0 \geqslant & \left[-\frac{C}{R^2}(1+\sqrt{k}R) - \frac{2C}{R^2} + \left(\frac{2\phi}{n} - \alpha' \right) \frac{\phi}{\alpha} - \frac{\gamma'}{\gamma} \phi - \frac{n\alpha^2}{2(\alpha-1)} \frac{C}{R^2} \right. \\ & \left. - \delta_1(l-1)\bar{u}_2 - \sqrt{CK} \right] \phi G + \frac{\phi^2 G^2}{\alpha^2 n \gamma} - \gamma \phi^2 \alpha^2 n^2 K^2 \\ & - \frac{l\alpha-1}{l-1} \gamma \phi^2 \bar{u}_2 \delta_2 - \gamma(l-1) \bar{u}_2 \phi^2 \delta_1 \alpha \varphi - \phi^2 \alpha \gamma \bar{u}_2 \delta_3. \end{aligned} \quad (2.23)$$

For the inequality $Ax^2 - 2Bx \leq C$, one has $x \leq \frac{2B}{A} + \left(\frac{C}{A} \right)^{\frac{1}{2}}$, where $A, B, C > 0$. By using this inequality to (2.23) and then we arrive at

$$\begin{aligned} \phi G(x, T_1) & \leq (\phi G)(x_1, t_1) \\ & \leq \left\{ n\gamma\alpha^2 \left[\frac{C}{R^2}(1+\sqrt{KR}) + \frac{n\alpha^2}{2(\alpha-1)} \frac{C}{R^2} + \delta_1(l-1)\bar{u}_2 + \sqrt{CK} \right] \right. \\ & \quad + n\gamma\alpha^2 \left[\frac{\gamma'}{\gamma} - \left(\frac{2\phi}{n} - \frac{\alpha'}{\alpha} \right) \frac{1}{\alpha} \right] + n^{\frac{3}{2}} \gamma \phi \alpha^2 K + \alpha \phi \gamma \sqrt{\frac{n(l\alpha-1)\bar{u}_2 \delta_2}{l-1}} \\ & \quad \left. + \alpha^{\frac{3}{2}} \gamma \phi \sqrt{n(l-1)\delta_1 \varphi} + \alpha^{\frac{3}{2}} \gamma \phi \sqrt{n\delta_3 \bar{u}_2} \right\} (x_1, t_1). \end{aligned}$$

If γ is nondecreasing which satisfies the system

$$\begin{cases} \frac{\gamma'}{\gamma} - \left(\frac{2\phi}{n} - \alpha' \right) \frac{1}{\alpha} \leq 0, \\ \frac{\gamma\alpha^4}{\alpha-1} \leq C_1. \end{cases} \quad (2.24)$$

Recall that $\alpha(t)$ and $\gamma(t)$ are non-decreasing and $t_1 < T_1$. Hence, we have

$$\begin{aligned} \phi G(x, T_1) & \leq (\phi G)(x_1, t_1) \\ & \leq n\gamma(T_1)\alpha^2(T_1) \left[\frac{C}{R^2} \left(1 + \sqrt{KR} \right) + K \right] + \frac{n^2 C}{R^2} \\ & \quad + n\gamma(T_1)\alpha^2(T_1)(l-1)\delta_1 \bar{u}_2 + n^{\frac{3}{2}} \gamma \phi(T_1)\alpha^2(T_1)K \\ & \quad + \alpha(T_1)\phi\gamma(T_1)\sqrt{\frac{n(l\alpha(T_1)-1)\bar{u}_2 \delta_2}{l-1}} \\ & \quad + \alpha^{\frac{3}{2}}(T_1)\gamma(T_1)\phi\sqrt{n(l-1)\delta_1 \varphi} + \alpha^{\frac{3}{2}}(T_1)\gamma(T_1)\phi\sqrt{n\delta_3 \bar{u}_2}. \end{aligned}$$

Hence, we have for $\phi \equiv 1$ on $B_{R,T}$,

$$F(x, T_1) \leq n\alpha^2(T_1) \left[\frac{C}{R^2} \left(1 + \sqrt{KR} \right) + K \right] + \frac{n^2 C}{R^2}$$

$$\begin{aligned}
& +n\alpha^2(T_1)(l-1)\delta_1\bar{u}_2+n^{\frac{3}{2}}\alpha^2(T_1)K \\
& +\alpha(T_1)\sqrt{\frac{n(l\alpha(T_1)-1)\bar{u}_2\delta_2}{l-1}} \\
& +\alpha^{\frac{3}{2}}(T_1)\sqrt{n(l-1)\delta_1\varphi}+\alpha^{\frac{3}{2}}(T_1)\sqrt{n\delta_3\bar{u}_2}.
\end{aligned}$$

If γ is nondecreasing which satisfies the system

$$\begin{cases} \frac{\gamma'}{\gamma}-\left(\frac{2\varphi}{n}-\alpha'\right)\frac{1}{\alpha}\leqslant 0, \\ \frac{\gamma}{\alpha-1}\leqslant C_2. \end{cases} \quad (2.25)$$

Recall that $\alpha(t)$ and $\gamma(t)$ are non-decreasing and $t_1 < T_1$. Hence, we have

$$\begin{aligned}
\phi G(x, T_1) &\leqslant (\phi G)(x_1, t_1) \\
&\leqslant n\gamma(T_1)\alpha^2(T_1)\left[\frac{C}{R^2}\left(1+\sqrt{KR}\right)+\frac{Cn\alpha^2(T)}{R^2}+CK\right] \\
&\quad +n\gamma(T_1)\alpha^2(T_1)(l-1)\delta_1\bar{u}_2+n^{\frac{3}{2}}\gamma(T_1)\phi\alpha^2(T_1)K \\
&\quad +\alpha(T_1)\phi\gamma(T_1)\sqrt{\frac{n(l\alpha(T_1)-1)\bar{u}_2\delta_2}{l-1}} \\
&\quad +\alpha^{\frac{3}{2}}(T_1)\gamma(T_1)\phi\sqrt{n(l-1)\delta_1\varphi}+\alpha^{\frac{3}{2}}(T_1)\gamma(T_1)\phi\sqrt{n\delta_3\bar{u}_2}.
\end{aligned}$$

Hence, we have for $\phi \equiv 1$ on $B_{R,T}$,

$$\begin{aligned}
F(x, T_1) &\leqslant n\alpha^2(T_1)\left[\frac{C}{R^2}\left(1+\sqrt{KR}\right)+\frac{Cn\alpha^2(T)}{R^2}+CK\right] \\
&\quad +n\alpha^2(T_1)(l-1)\delta_1\bar{u}_2+n^{\frac{3}{2}}\alpha^2(T_1)K \\
&\quad +\alpha(T_1)\sqrt{\frac{n(l\alpha(T_1)-1)\bar{u}_2\delta_2}{l-1}} \\
&\quad +\alpha^{\frac{3}{2}}(T_1)\sqrt{n(l-1)\delta_1\varphi}+\alpha^{\frac{3}{2}}(T_1)\sqrt{n\delta_3\bar{u}_2}.
\end{aligned}$$

Because T_1 is arbitrary in $0 < T_1 < T$, the conclusion is valid.

Proof of Theorem 2.2. Since $u_t = \Delta u + h(t)u^l$, we have

$$\begin{aligned}
\partial_t(|\nabla u|^2) &= 2\text{Ric}(\nabla u, \nabla u) + 2 < \nabla u, \nabla(u_t) > \\
&= 2\text{Ric}(\nabla u, \nabla u) + 2 < \nabla u, \nabla(\Delta u) > + 2 < \nabla u, \nabla(h(t)u^l) >.
\end{aligned}$$

Applying the Bochner's formula, the above equation becomes

$$\partial_t(|\nabla u|^2) = \Delta(|\nabla u|^2) - 2|\nabla^2 u|^2 + 2 < \nabla u, \nabla(h(t)u^l) >. \quad (2.26)$$

Besides,

$$\partial_t(u^2) = \Delta(u^2) - 2|\nabla u|^2 + 2h(t)u^{l+1}. \quad (2.27)$$

Let $\bar{F} = t|\nabla u|^2 + Xu^2$, where X decided later. Then combining (2.26) with (2.27), we obtain

$$\begin{aligned}\partial_t \bar{F} &= |\nabla u|^2 + t[\Delta(|\nabla u|^2) - 2|\nabla^2 u|^2 + 2 < \nabla u, \nabla(h(t)u^l) >] \\ &\quad + X[\Delta(u^2) - 2|\nabla u|^2 + 2h(t)u^{l+1}] \\ &= |\nabla u|^2 + t[\Delta(|\nabla u|^2) - 2|\nabla^2 u|^2 + 2h(t)(l-1)u^{l-2}|\nabla u|^2] \\ &\quad + X[\Delta(u^2) - 2|\nabla u|^2 + 2h(t)u^{l+1}] \\ &\leq \Delta \bar{F} + (1-2X)|\nabla u|^2.\end{aligned}\tag{2.28}$$

Selecting $X = \frac{1}{2}$ and using the maximum principle, we infer

$$\bar{F}(x, t) \leq \max_{x \in M^n} \bar{F}(x, 0) = \frac{1}{2} \max_{x \in M^n} u^2(x, 0),$$

which implies the theorem is valid.

3. Gradient estimates for the equation (1.6)

Recall that (M^n, g) is called a gradient Ricci soliton if there is a smooth function f on M^n such that for some constant $c \in \mathbb{R}$, which satisfies

$$Rc = cg + D^2f,\tag{3.1}$$

where D^2f is the Hessian of f . Letting $u = e^f$, after some computation applying (3.1) as done in [21], we get

$$\Delta u + 2cu \log u = (A_0 - nc)u \quad \text{in } M^n$$

for some constant A_0 , where n is the dimension of M^n . In [21], Ma proved a local gradient estimate of positive solutions to the equation

$$\Delta u + au \log u + bu = 0 \quad \text{in } M^n,$$

where $a > 0$ and $b \in \mathbb{R}$ are constants for complete noncompact manifolds with a fixed metric and curvature locally bounded below. In [30], Yang generalized Ma's result and derived a local gradient estimates for positive solutions to the equation

$$u_t = \Delta u + au \log u + bu \quad \text{in } M^n \times (0, T],$$

where $a, b \in \mathbb{R}$ are constants for complete noncompact manifolds with a fixed metric and curvature locally bounded below. Replacing u by $e^{b/a}u$, the above equation becomes

$$u_t = \Delta u + au \log u.\tag{3.2}$$

One can find in [29, 30] some related results for equation (3.2) on manifolds.

In this section, we consider the nonlinear parabolic equation (1.6) along the Ricci flow.

3.1. Main results

Our main results state as follows.

THEOREM 3.1. *Let $(M^n, g(t))_{t \in [0, T]}$ be a complete solution to the Ricci flow (1.7). Assume that $|\text{Ric}(x, t)| \leq K$ for some $K > 0$ and all $t \in [0, T]$. Suppose that there exist three functions $\alpha(t)$, $\varphi(t)$ and $\gamma(t)$ which satisfy the following conditions (C1), (C2), (C3) and (C4).*

Given $x_0 \in M$ and $R > 0$, let u be a positive solution of the nonlinear parabolic equation

$$\partial_t u = \Delta u + a u \log u$$

in the cube $B_{2R, T} := \{(x, t) | d(x, x_0, t) \leq 2R, 0 \leq t \leq T\}$, where a is a constant.

(1) *For $a \leq 0$. If $\frac{\gamma\alpha^4}{\alpha-1} \leq C_1$ for some constant C_1 , then*

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) \\ &+ \frac{Cn^2}{R^2\gamma} + n^{\frac{3}{2}}\alpha^2 K + n|a|\alpha^2 + \alpha\varphi, \end{aligned}$$

where $C = C(n, C_1)$ is a constant.

If $\frac{\gamma}{\alpha-1} \leq C_2$ for some constant C_2 , then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) \\ &+ \frac{Cn^2\alpha^4}{R^2\gamma} + n^{\frac{3}{2}}\alpha^2 K + n|a|\alpha^2 + \alpha\varphi, \end{aligned}$$

where $C = C(n, C_2)$ is a constant.

(2) *For $a > 0$. If $\frac{\gamma\alpha^4}{\alpha-1} \leq C_1$ for some constant C_1 , then*

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + a + K \right) + \frac{n^2 C}{R^2\gamma} \\ &+ n^{\frac{3}{2}}\alpha^2 K + \alpha\varphi, \end{aligned}$$

where $C = C(n, C_1)$ is a constant.

If $\frac{\gamma}{\alpha-1} \leq C_2$ for some constant C_2 , then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + a + K \right) + \frac{n^2 C \alpha^4}{R^2\gamma} \\ &+ n^{\frac{3}{2}}\alpha^2 K + \alpha\varphi, \end{aligned}$$

where $C = C(n, C_2)$ is a constant.

COROLLARY 3.1. *Let $(M^n, g(t))_{t \in [0, T]}$ be a complete solution to the Ricci flow (1.7). Assume that $|\text{Ric}(x, t)| \leq K$ for some $K > 0$ and all $t \in [0, T]$. Given $x_0 \in M$ and $R > 0$, let u be a positive solution of the nonlinear parabolic equation (1.6) in the cube $B_{2R, T} := \{(x, t) | d(x, x_0, t) \leq 2R, 0 \leq t \leq T\}$. Then the following special estimates are valid.*

1. *Li-Yau type:*

$$\alpha(t) = \text{constant}, \quad \varphi(t) = \frac{\alpha n}{t} + \frac{nK\alpha^2}{\alpha - 1}, \quad \gamma(t) = t^\theta \quad \text{with} \quad 0 < \theta \leq 2.$$

If $a \leq 0$, then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + \frac{\alpha^2}{\alpha - 1} \frac{1}{R^2} + K \right) \\ &\quad + n^{\frac{3}{2}} \alpha^2 K + n|a|\alpha^2 + \alpha\varphi. \end{aligned}$$

If $a > 0$, then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + \frac{\alpha^2}{\alpha - 1} \frac{1}{R^2} + a + K \right) \\ &\quad + n^{\frac{3}{2}} \alpha^2 K + \alpha\varphi. \end{aligned}$$

2. *Hamilton type:*

$$\alpha(t) = e^{2Kt}, \quad \varphi(t) = \frac{n}{t} e^{4Kt}, \quad \gamma(t) = t e^{2Kt}.$$

If $a \leq 0$, then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) \\ &\quad + \frac{C\alpha^4}{R^2 t e^{2Kt}} + \alpha\varphi + n^{\frac{3}{2}} \alpha^2 K + n|a|\alpha^2 + \alpha\varphi. \end{aligned}$$

If $a > 0$, then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + a + K \right) \\ &\quad + \frac{C\alpha^4}{R^2 t e^{2Kt}} + n^{\frac{3}{2}} \alpha^2 K + \alpha\varphi. \end{aligned}$$

3. *Li-Xu type:*

$$\alpha(t) = 1 + \frac{\sinh(Kt) \cosh(Kt) - Kt}{\sinh^2(Kt)}, \quad \varphi(t) = 2nK[1 + \coth(Kt)],$$

$$\gamma(t) = \tanh(Kt).$$

If $a \leq 0$, then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) \\ &+ \frac{C}{R^2 \tanh(Kt)} + n^{\frac{3}{2}} \alpha^2 K + n|a|\alpha^2 + \alpha\varphi. \end{aligned}$$

If $a > 0$, then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + a + K \right) \\ &+ \frac{C}{R^2 \tanh(Kt)} + n^{\frac{3}{2}} \alpha^2 K + \alpha\varphi. \end{aligned}$$

4. Linear Li-Xu type:

$$\alpha(t) = 1 + 2Kt, \varphi(t) = \frac{n}{t} + nK(1 + 2Kt + \mu Kt), \gamma(t) = Kt \quad \text{with } \mu \geq \frac{1}{4}.$$

If $a \leq 0$, then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) \\ &+ \frac{C\alpha^4}{R^2 Kt} + n^{\frac{3}{2}} \alpha^2 K + n|a|\alpha^2 + \alpha\varphi. \end{aligned}$$

If $a > 0$, then

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u &\leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + a + K \right) \\ &+ \frac{C\alpha^4}{R^2 Kt} + n^{\frac{3}{2}} \alpha^2 K + \alpha\varphi. \end{aligned}$$

The local estimates above imply global estimates.

COROLLARY 3.2. Let $(M^n, g(0))$ be a complete noncompact Riemannian manifold without boundary, and assume $g(t)$ evolves by Ricci flow in such a way that $|\text{Ric}| \leq K$ for $t \in [0, T]$. Let $u(x, t)$ be a positive solution to the equation (1.6). If $l \in \mathbb{R}$ and for $(x, t) \in M^n \times (0, T]$, then

$$\frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} + \alpha a \log u \leq C\alpha^2(K + |a|) + \alpha\varphi.$$

REMARK 3.1. The above results may be regarded as generalizations the gradient estimates of Yang [30] to the Ricci flow.

3.2. Auxiliary lemma

To prove the Theorem 3.1, the following lemma is needed.

Let $f = \log u$. Then

$$(\Delta - \partial_t)f = -|\nabla f|^2 - af. \quad (3.3)$$

Let $F = |\nabla f|^2 - \alpha f_t + \alpha a f - \alpha \varphi$, where $\alpha = \alpha(t)$ and $\varphi = \varphi(t)$. Then

$$\begin{aligned} \Delta f &= f_t - af - |\nabla f|^2 \\ &= -\frac{F}{\alpha} - \left(\frac{\alpha-1}{\alpha}\right)|\nabla f|^2 - \varphi. \end{aligned} \quad (3.4)$$

LEMMA 3.1. *We assume that $\alpha(t) > 1$ and $\varphi(t) > 0$ satisfy the system (2.3). Then*

$$\begin{aligned} (\Delta - \partial_t)F &\geq |f_{ij} + \frac{\varphi}{n} \delta_{ij}|^2 + \left(\frac{2\varphi}{n} - \alpha'\right) \frac{1}{\alpha} F - \alpha^2 n^2 K^2 - 2\nabla f \nabla F \\ &\quad + 2a(\alpha-1)|\nabla f|^2 + a\alpha\Delta f. \end{aligned} \quad (3.5)$$

Proof. A computation is shown that

$$\begin{aligned} \Delta F &= \Delta|\nabla f|^2 - \alpha\Delta(f_t) + \alpha a\Delta f \\ &= 2|f_{ij}|^2 + 2f_j f_{iij} + 2R_{ij} f_i f_j - \alpha\Delta(f_t) + \alpha a\Delta f \\ &= 2(|f_{ij}|^2 + \alpha R_{ij} f_{ij}) + 2f_j f_{iij} + 2R_{ij} f_i f_j - \alpha(\Delta f)_t + \alpha a\Delta f \\ &\geq 2|f_{ij}|^2 - 2\alpha|R_{ij}||f_{ij}| + 2f_j f_{iij} + 2R_{ij} f_i f_j - \alpha(\Delta f)_t + \alpha a\Delta f \\ &\geq 2|f_{ij}|^2 - (\alpha^2|R_{ij}|^2 + |f_{ij}|^2) + 2f_j f_{iij} + 2R_{ij} f_i f_j - \alpha(\Delta f)_t + \alpha a\Delta f \\ &\geq |f_{ij}|^2 - \alpha^2|R_{ij}|^2 + 2f_j f_{iij} + 2R_{ij} f_i f_j - \alpha(\Delta f)_t + \alpha a\Delta f \\ &\geq |f_{ij}|^2 - \alpha^2 n^2 K^2 + 2f_j f_{iij} + 2R_{ij} f_i f_j - \alpha(\Delta f)_t + \alpha a\Delta f, \end{aligned} \quad (3.6)$$

and

$$\begin{aligned} \partial_t F &= (|\nabla f|^2)_t - \alpha f_{tt} - \alpha' f_t + \alpha' a f + \alpha a f_t - \alpha \varphi' - \alpha' \varphi \\ &= 2\nabla f \nabla(f_t) + 2R_{ij} f_i f_j - \alpha f_{tt} - \alpha' f_t + \alpha' a f \\ &\quad + \alpha a f_t - \alpha \varphi' - \alpha' \varphi. \end{aligned} \quad (3.7)$$

We follow that from (3.6) and (3.7)

$$\begin{aligned} (\Delta - \partial_t)F &\geq |f_{ij}|^2 - \alpha^2 n^2 K^2 + 2\nabla f \nabla(\Delta f) - \alpha(\Delta f)_t + \alpha a \Delta d - 2\nabla f \nabla(f_t) \\ &\quad + \alpha f_{tt} + \alpha' f_t - \alpha' a f - \alpha a f_t + \alpha \varphi' + \alpha' \varphi \\ &= |f_{ij}|^2 - \alpha^2 n^2 K^2 + 2\nabla f \nabla(\Delta f) - \alpha(f_t - |\nabla f|^2 - af)_t \\ &\quad + \alpha a \Delta f - 2\nabla f \nabla(f_t) + \alpha f_{tt} + \alpha' f_t \\ &\quad - \alpha' a f - \alpha c f_t + \alpha \varphi' + \alpha' \varphi \end{aligned}$$

$$\begin{aligned}
&= |f_{ij}|^2 - \alpha^2 n^2 K^2 + 2\nabla f \nabla (\Delta f) + \alpha(|\nabla f|^2)_t + \alpha a \Delta f \\
&\quad - 2\nabla f \nabla (f_t) + \alpha' f_t - \alpha' a f + \alpha \varphi' + \alpha' \varphi \\
&= |f_{ij}|^2 - \alpha^2 n^2 K^2 + 2\nabla f \nabla (\Delta f) + 2\alpha \nabla f \nabla (f_t) + 2\alpha R_{ij} f_i f_j \\
&\quad + \alpha a \Delta f - 2\nabla f \nabla (f_t) + \alpha' f_t - \alpha' a f + \alpha \varphi' + \alpha' \varphi \\
&= |f_{ij}|^2 + 2\alpha R_{ij} f_i f_j - \alpha^2 n^2 K^2 + 2\nabla f \nabla F + 2a(\alpha - 1)(l - 1)|\nabla f|^2 \\
&\quad + \alpha a \Delta f + \alpha' f_t - \alpha' a f + \alpha \varphi' + \alpha' \varphi. \tag{3.8}
\end{aligned}$$

Further, by utilizing the unit matrix $(\delta_{ij})_{n \times n}$ and (3.8), we obtain

$$\begin{aligned}
(\Delta - \partial_t)F &\geq |f_{ij} + \frac{\varphi}{n} \delta_{ij}|^2 - 2\alpha K |\nabla f|^2 - \alpha^2 n^2 K^2 + 2\nabla f \nabla F \\
&\quad + 2a(\alpha - 1) |\nabla f|^2 + a\alpha \Delta f + \alpha' f_t - \alpha' a f + \alpha \varphi' + \alpha' \varphi \\
&\quad - \frac{\varphi^2}{n} - 2\frac{\varphi}{n} \Delta f \\
&= |f_{ij} + \frac{\varphi}{n} \delta_{ij}|^2 + \left(\frac{2\varphi}{n} - 2\alpha K\right) |\nabla f|^2 - \left(\frac{2\varphi}{n} - \alpha'\right) f_t \\
&\quad + \left(\frac{2\varphi}{n} - \alpha'\right) c u^{l-1} - \alpha^2 n^2 K^2 - 2\nabla f \nabla F + 2a(\alpha - 1) |\nabla f|^2 \\
&\quad + a\alpha \Delta f + \alpha \varphi' + \alpha' \varphi - \frac{\varphi^2}{n} + \left(\frac{2\varphi}{n} - \alpha'\right) \frac{\alpha \varphi}{\alpha}. \tag{3.9}
\end{aligned}$$

3.3. The proof of Theorem

In this section, we will prove Theorem 3.1.

Proof of Theorem 3.1. Let $G = \gamma(t)F$ and $\gamma(t) > 0$ be non-decreasing. Then

$$\begin{aligned}
(\Delta - \partial_t)G &= \gamma(\Delta - \partial_t)F - \gamma' F \\
&\geq \gamma |f_{ij} + \frac{\varphi}{n} g_{ij}|^2 + \left(\frac{2\varphi}{n} - \alpha'\right) \frac{1}{\alpha} G - \gamma \alpha^2 n^2 K^2 - 2\nabla f \nabla G \\
&\quad + 2a\gamma(\alpha - 1) |\nabla f|^2 + a\gamma \alpha \Delta f - \gamma' F \\
&= \gamma |f_{ij} + \frac{\varphi}{n} g_{ij}|^2 + \left[\left(\frac{2\varphi}{n} - \alpha'\right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma}\right] G - \gamma \alpha^2 n^2 K^2 \\
&\quad - 2\nabla f \nabla G + 2a\gamma(\alpha - 1) |\nabla f|^2 + a\gamma \alpha \Delta f. \tag{3.10}
\end{aligned}$$

Now, let $\varphi(r)$ be a C^2 function on $[0, \infty)$ such that

$$\varphi(r) = \begin{cases} 1, & \text{if } r \in [0, 1], \\ 0, & \text{if } r \in [2, \infty), \end{cases}$$

and

$$0 \leq \varphi(r) \leq 1, \quad \varphi'(r) \leq 0, \quad \varphi''(r) \leq 0, \quad \frac{|\varphi'(r)|}{\varphi(r)} \leq C,$$

where C is an absolute constant. Define by

$$\phi(x, t) = \varphi(d(x, x_0, t)) = \varphi\left(\frac{d(x, x_0, t)}{R}\right) = \varphi\left(\frac{\rho(x, t)}{R}\right),$$

where $\rho(x, t) = d(x, x_0, t)$. By using maximum principle, the argument of Calabi [2] allows us to suppose that the function $\phi(x, t)$ with support in $B_{2R, T}$, is C^2 at the maximum point. By utilizing the Laplacian theorem, we deduce that

$$\frac{|\nabla \phi|^2}{\phi} \leq \frac{C}{R^2}, \quad -\Delta \phi \leq \frac{C}{R^2}(1 + \sqrt{K}R). \quad (3.11)$$

For any $0 \leq T_1 \leq T$, let $H = \phi G$ and (x_1, t_1) be the point in B_{2R, T_1} at which H attains its maximum value. We can suppose that H is positive, because otherwise the proof is trivial. Then at the point (x_1, t_1) , we infer

$$\left. \begin{aligned} 0 &= \nabla(\phi G) = G\nabla\phi + \phi\nabla G, \\ \Delta(\phi G) &\leq 0, \\ \partial_t(\phi G) &\geq 0. \end{aligned} \right\} \quad (3.12)$$

By the evolution formula of the geodesic length along the Ricci flow [6], we calculate

$$\begin{aligned} \phi_t G &= -G\phi' \left(\frac{\rho}{R}\right) \frac{1}{R} \frac{d\rho}{dt} = G\phi' \left(\frac{\rho}{R}\right) \int_{\gamma_1} \text{Ric}(S, S) ds \\ &\leq G\phi' \left(\frac{\rho}{R}\right) \frac{1}{R} K\rho \leq G\phi' \left(\frac{\rho}{R}\right) K_2 \leq G\sqrt{C}K, \end{aligned}$$

where γ_1 is the geodesic connecting x and x_0 along the metric $g(t_1)$, S is the unite tangent vector to γ_1 , and ds is the element of the arc length.

All the following computations are at the point (x_1, t_1) . Since

$$\begin{aligned} |f_{ij} + \frac{\varphi}{n}\delta_{ij}|^2 &\geq \frac{1}{n} \left(\text{tr}|f_{ij} + \frac{\varphi}{n}\delta_{ij}| \right)^2 \\ &= \frac{1}{n} (\Delta f + \varphi)^2 \\ &= \frac{1}{n} \left[-\frac{1}{\alpha}F - \frac{1}{\alpha}(\alpha-1)|\nabla f|^2 \right]^2 \\ &= \frac{1}{\alpha^2 n} \left[\frac{G}{\gamma} + (\alpha-1)|\nabla f|^2 \right]^2, \end{aligned} \quad (3.13)$$

and

$$\begin{aligned} \Delta f &= f_t - |\nabla f|^2 - af \\ &= -\frac{F}{\alpha} - \frac{\alpha-1}{\alpha}|\nabla f|^2 - \varphi < 0. \end{aligned} \quad (3.14)$$

Case 1 $a \leq 0$. Combining (3.14) with (3.10), we have

$$\begin{aligned} (\Delta - \partial_t)G &\geq \gamma|f_{ij}| + \frac{\varphi}{n}\delta_{ij}|^2 + \left[\left(\frac{2\varphi}{n} - \alpha'\right)\frac{1}{\alpha} - \frac{\gamma'}{\gamma}\right]G - \gamma\alpha^2 n^2 K^2 \\ &\quad - 2\nabla f \nabla G + 2a\gamma(\alpha - 1)|\nabla f|^2. \end{aligned}$$

Using (3.12) and (3.13), we infer

$$\begin{aligned} 0 &\geq (\Delta - \partial_t)(\phi G) \\ &= G\left(\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi}\right) + \phi(\Delta - \partial_t)G - \gamma G\phi_t \\ &\geq G\left(\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi}\right) + \frac{\phi\gamma}{\alpha^2 n}\left[\frac{G}{\gamma} + (\alpha - 1)|\nabla f|^2\right]^2 \\ &\quad + \left[\left(\frac{2\varphi}{2} - \alpha'\right)\frac{1}{\alpha} - \frac{\gamma'}{\gamma}\right]\phi G - \gamma\phi\alpha^2 n^2 K^2 - 2\phi\nabla f \nabla G \\ &\quad + 2a\phi\gamma(\alpha - 1)|\nabla f|^2 - G\sqrt{CK}. \end{aligned} \tag{3.15}$$

Multiply ϕ to (3.15), and we have

$$\begin{aligned} 0 &\geq \phi G\left[\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi} + \left(\frac{2\varphi}{n} - \alpha'\right)\frac{1}{\alpha} - \frac{\gamma'}{\gamma}\right] + \frac{\phi^2\gamma}{\alpha^2 n}\left[\frac{G}{\gamma} + (\alpha - 1)|\nabla f|^2\right]^2 \\ &\quad - \gamma\phi^2\alpha^2 n^2 K^2 - 2\phi^2\nabla f \nabla G + 2a\phi^2\gamma(\alpha - 1)|\nabla f|^2 - \phi G\sqrt{CK} \\ &\geq \phi G\left[\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi} + \left(\frac{2\varphi}{n} - \alpha'\right)\frac{1}{\alpha} - \frac{\gamma'}{\gamma}\right] + \frac{\phi^2 G^2}{\alpha^2 n \gamma} + \frac{\phi^2(\alpha - 1)^2 \gamma}{n \alpha^2} |\nabla f|^4 \\ &\quad + \frac{2\phi^2(\alpha - 1)}{n \alpha^2} G |\nabla f|^2 - \gamma\phi^2\alpha^2 n^2 K^2 + 2\phi G \nabla \phi \nabla f \\ &\quad + 2a\phi^2\gamma(\alpha - 1)|\nabla f|^2 - \phi G\sqrt{CK}. \end{aligned} \tag{3.16}$$

Using the fact

$$\frac{2\phi^2(\alpha - 1)}{n \alpha^2} G |\nabla f|^2 + 2\phi G \nabla \phi \nabla f \geq -\frac{n \alpha^2}{2(\alpha - 1)} \frac{|\nabla\phi|^2}{\phi} \phi G,$$

and

$$\frac{\phi^2(\alpha - 1)^2 \gamma}{n \alpha^2} |\nabla f|^4 + 2a\phi^2\gamma(\alpha - 1)|\nabla f|^2 \geq -na^2\alpha^2\gamma\phi^2,$$

to (3.16), we deduce that

$$\begin{aligned} 0 &\geq \phi G\left[\Delta\phi - 2\frac{|\nabla\phi|^2}{\phi} + \left(\frac{2\varphi}{n} - \alpha'\right)\frac{1}{\alpha} - \frac{\gamma'}{\gamma} - \frac{n \alpha^2}{2(\alpha - 1)} \frac{|\nabla\phi|^2}{\phi} - \sqrt{CK}\right] \\ &\quad + \frac{\phi^2 G^2}{\alpha^2 n \gamma} - \gamma\phi^2\alpha^2 n^2 K^2 - na^2\alpha^2\gamma\phi^2 \\ &\geq \left[-\frac{C}{R^2}(1 + \sqrt{k}R) - \frac{2C}{R^2} + \left(\frac{2\varphi}{n} - \alpha'\right)\frac{1}{\alpha} - \frac{\gamma'}{\gamma} - \frac{n \alpha^2}{2(\alpha - 1)} \frac{C}{R^2} - \sqrt{CK}\right] \phi G \end{aligned}$$

$$+\frac{\phi^2 G^2}{\alpha^2 n \gamma} - \gamma \phi^2 \alpha^2 n^2 K^2 - n a^2 \alpha^2 \gamma \phi^2.$$

For the inequality $Ax^2 - 2Bx \leq C$, one has $x \leq \frac{2B}{A} + \left(\frac{C}{A}\right)^{\frac{1}{2}}$, where $A, B, C > 0$. Hence, we infer

$$\begin{aligned} \phi G(x, T_1) &\leq (\phi G)(x_1, t_1) \\ &\leq \left\{ n \gamma \alpha^2 \left[\frac{C}{R^2} (1 + \sqrt{KR}) + \frac{n \alpha^2}{2(\alpha - 1)} \frac{C}{R^2} + \sqrt{CK} \right] \right. \\ &\quad + n \gamma \alpha^2 \left[\frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} \right] \\ &\quad \left. + n^{\frac{3}{2}} \gamma \alpha^2 \phi K + n a \alpha^2 \gamma \phi \right\} (x_1, t_1). \end{aligned}$$

If γ is nondecreasing which satisfies the system

$$\begin{cases} \frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} \leq 0, \\ \frac{\gamma \alpha^4}{\alpha - 1} \leq C_1. \end{cases}$$

Recall that $\alpha(t)$ and $\gamma(t)$ are non-decreasing and $t_1 < T_1$. Hence, we have

$$\begin{aligned} \phi G(x, T_1) &\leq (\phi G)(x_1, t_1) \\ &\leq n \gamma(T_1) \alpha^2(T_1) \left[\frac{C}{R^2} (1 + \sqrt{KR}) + K \right] + \frac{n^2 C}{R^2} \\ &\quad + n^{\frac{3}{2}} \gamma(T_1) \alpha^2(T_1) K + n a \alpha^2(T_1) \gamma(T_1). \end{aligned}$$

Hence, we have for $\phi \equiv 1$ on $B_{R,T}$,

$$\begin{aligned} F(x, T_1) &\leq n \alpha^2(T_1) \left[\frac{C}{R^2} (1 + \sqrt{KR}) + CK \right] + \frac{n^2 C}{R^2 \gamma(T_1)} \\ &\quad + n^{\frac{3}{2}} \alpha^2(T_1) K + n a \alpha^2(T_1). \end{aligned}$$

If γ is nondecreasing which satisfies the system

$$\begin{cases} \frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} \leq 0, \\ \frac{\gamma}{\alpha - 1} \leq C_2. \end{cases}$$

Recall that $\alpha(t)$ and $\gamma(t)$ are non-decreasing and $t_1 < T_1$. Hence, we have

$$\phi G(x, T_1) \leq (\phi G)(x_1, t_1)$$

$$\leq n\gamma(T_1)\alpha^2(T_1) \left[\frac{C}{R^2} \left(1 + \sqrt{KR} \right) + \frac{Cn\alpha^4}{R^2} + CK \right] \\ + n^{\frac{3}{2}}\gamma(T_1)\alpha^2(T_1)K + na\alpha^2(T_1)\gamma(T_1).$$

Hence, we have for $\phi \equiv 1$ on $B_{R,T}$,

$$F(x, T_1) \leq n\alpha^2(T_1) \left[\frac{C}{R^2} \left(1 + \sqrt{KR} \right) + K \right] + \frac{n^2 C \alpha^4}{R^2 \gamma(T_1)} \\ + n^{\frac{3}{2}} \alpha^2(T_1)K + na\alpha^2(T_1).$$

Because T_1 is arbitrary in $0 < T_1 < T$, the conclusion is valid.

Case 2 $a \geq 0$. It is not difficult to find $\Delta f \leq -\frac{F}{\alpha}$ from (3.14). Then, we have from (3.10)

$$(\Delta - \partial_t)G \geq \gamma|f_{ij}| + \frac{\varphi}{n}\delta_{ij}|^2 + \left[\left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma} \right] G \\ - \gamma\alpha^2 n^2 K^2 2\nabla f \nabla G - aG.$$

Using (3.13) and (3.13), we infer

$$0 \geq (\Delta - \partial_t)(\phi G) \\ = G \left(\Delta\phi - 2 \frac{|\nabla\phi|^2}{\phi} \right) + \phi(\Delta - \partial_t)G - \gamma G\phi_t \\ \geq G \left(\Delta\phi - 2 \frac{|\nabla\phi|^2}{\phi} \right) + \frac{\phi\gamma}{\alpha^2 n} \left[\frac{G}{\gamma} + (\alpha - 1)|\nabla f|^2 \right]^2 \\ + \left[\left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma} \right] \phi G - \gamma\phi\alpha^2 n^2 K^2 - 2\phi\nabla f \nabla G \\ - a\phi G - G\sqrt{CK}.$$

Multiply ϕ , and we have

$$0 \geq \phi G \left[\Delta\phi - 2 \frac{|\nabla\phi|^2}{\phi} + \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma} \right] + \frac{\phi^2\gamma}{\alpha^2 n} \left[\frac{G}{\gamma} + (\alpha - 1)|\nabla f|^2 \right]^2 \\ - \gamma\phi^2\alpha^2 n^2 K^2 - 2\phi^2\nabla f \nabla G - a\phi^2 G - \phi G\sqrt{CK} \\ \geq \phi G \left[\Delta\phi - 2 \frac{|\nabla\phi|^2}{\phi} + \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma} \right] + \frac{\phi^2 G^2}{\alpha^2 n \gamma} + \frac{2\phi^2(\alpha - 1)}{n\alpha^2} G |\nabla f|^2 \\ - \gamma\phi^2\alpha^2 n^2 K^2 + 2\phi G \nabla \phi \nabla f - a\phi^2 G - \phi G\sqrt{CK}, \tag{3.17}$$

where we dropped the term $\frac{\phi^2(\alpha - 1)^2\gamma}{n\alpha^2} |\nabla f|^4$. We use the fact

$$\frac{2\phi^2(\alpha - 1)}{n\alpha^2} G |\nabla f|^2 + 2\phi G \nabla \phi \nabla f \geq -\frac{n\alpha^2}{2(\alpha - 1)} \frac{|\nabla\phi|^2}{\phi} \phi G,$$

to (3.17), we deduce that

$$\begin{aligned} 0 &\geq \phi G \left[\Delta\phi - 2 \frac{|\nabla\phi|^2}{\phi} + \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma} - \frac{n\alpha^2}{2(\alpha-1)} \frac{|\nabla\phi|^2}{\phi} - a\phi - \sqrt{CK} \right] \\ &\quad + \frac{\phi^2 G^2}{\alpha^2 n \gamma} - \gamma \phi^2 \alpha^2 n^2 K^2 \\ &\geq \left[-\frac{C}{R^2} (1 + \sqrt{k}R) - \frac{2C}{R^2} + \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} - \frac{\gamma'}{\gamma} - \frac{n\alpha^2}{2(\alpha-1)} \frac{C}{R^2} - \sqrt{CK} \right] \phi G \\ &\quad + \frac{\phi^2 G^2}{\alpha^2 n \gamma} - \gamma \phi^2 \alpha^2 n^2 K^2. \end{aligned}$$

For the inequality $Ax^2 - 2Bx \leq C$, one has $x \leq \frac{2B}{A} + \left(\frac{C}{A}\right)^{\frac{1}{2}}$, where $A, B, C > 0$.

$$\begin{aligned} \phi G(x, T_1) &\leq (\phi G)(x_1, t_1) \\ &\leq \left\{ n\gamma\alpha^2 \left[\frac{C}{R^2} (1 + \sqrt{KR}) + \frac{n\alpha^2}{2(\alpha-1)} \frac{C}{R^2} + a\phi + \sqrt{CK} \right] \right. \\ &\quad \left. + n\gamma\alpha^2 \left[\frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \frac{\alpha'}{\alpha} \right) \frac{1}{\alpha} \right] + n^{\frac{3}{2}} \gamma \alpha^2 \phi K \right\} (x_1, t_1). \end{aligned}$$

If γ is nondecreasing which satisfies the system

$$\begin{cases} \frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} \leq 0, \\ \frac{\gamma \alpha^4}{\alpha - 1} \leq C_1. \end{cases}$$

Recall that $\alpha(t)$ and $\gamma(t)$ are non-decreasing and $t_1 < T_1$. Hence, we have

$$\begin{aligned} \phi G(x, T_1) &\leq (\phi G)(x_1, t_1) \\ &\leq n\gamma(T_1)\alpha^2(T_1) \left[\frac{C}{R^2} (1 + \sqrt{KR}) + a\phi + K \right] + \frac{n^2 C}{R^2} \\ &\quad + n^{\frac{3}{2}} \gamma(T_1) \alpha^2(T_1) K. \end{aligned}$$

Hence, we have for $\phi \equiv 1$ on $B_{R,T}$,

$$\begin{aligned} \sup_{B_R} F(x, T_1) &\leq n\alpha^2(T_1) \left[\frac{C}{R^2} (1 + \sqrt{KR}) + a + CK \right] + \frac{n^2 C}{R^2 \gamma(T_1)} \\ &\quad + n^{\frac{3}{2}} \alpha^2(T_1) K. \end{aligned}$$

If γ is nondecreasing which satisfies the system

$$\begin{cases} \frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} \leq 0, \\ \frac{\gamma}{\alpha - 1} \leq C_2. \end{cases}$$

Recall that $\alpha(t)$ and $\gamma(t)$ are non-decreasing and $t_1 < T_1$. Hence, we have

$$\begin{aligned}\phi G(x, T_1) &\leq (\phi G)(x_1, t_1) \\ &\leq n\gamma(T_1)\alpha^2(T_1) \left[\frac{C}{R^2} \left(1 + \sqrt{KR} \right) + \frac{Cn\alpha^4}{R^2} + a\phi + CK \right] \\ &\quad + n^{\frac{3}{2}}\gamma(T_1)\alpha^2(T_1)K.\end{aligned}$$

Hence, we have for $\phi \equiv 1$ on $B_{R,T}$,

$$\begin{aligned}F(x, T_1) &\leq n\alpha^2(T_1) \left[\frac{C}{R^2} \left(1 + \sqrt{KR} \right) + a + K \right] + \frac{n^2 C \alpha^4}{R^2 \gamma(T_1)} \\ &\quad + n^{\frac{3}{2}} \alpha^2(T_1)K.\end{aligned}$$

Because T_1 is arbitrary in $0 < T_1 < T$, the conclusion is valid. This proof is complete.

4. Harnack inequalities

In this section, as application of main theorems, some Harnack inequalities are derived.

THEOREM 4.1. *Let $(M^n, g(x, t))_{t \in [0, T]}$ be a complete solution to the Ricci flow (1.7). Suppose that $|\text{Ric}| \leq K$ for some $K > 0$, and all $(x, t) \in M^n \times [0, T]$. Assume that $u(x, t)$ is a positive solution to (1.6). Let $h(x, t)$ be a function defined on $M^n \times [0, T]$ which is C^1 in t and C^2 in x , satisfying $|\nabla h|^2 \leq \delta_2 h$ and $\Delta h \geq -\delta_3$ on $M^n \times [0, T]$ for some positive constants δ_2 and δ_3 . Then for all $(x_1, t_1) \in M^n \times (0, T)$ and $(x_2, t_2) \in M^n \times (0, T)$ such that $t_1 < t_2$, we have*

$$u(x_2, t_2) \leq \begin{cases} u(x_1, t_1) \times \exp(\Gamma(t_1, t_2, \delta_1, \delta_2, \delta_3, \bar{u}_1)), & l \leq 1, \\ u(x_1, t_1) \times \exp(\Lambda(t_1, t_2, \delta_1, \delta_2, \delta_3, \bar{u}_1)), & l > 1, \end{cases}$$

where $\gamma(s)$ is a smooth curve connecting x_1 and x_2 with $\gamma(1) = x_1$ and $\gamma(0) = x_2$, and

$$\begin{aligned}&\Gamma(t_1, t_2, \delta_1, \delta_2, \delta_3, \bar{u}_1) \\ &= \int_0^1 \frac{|\gamma'(s)|^4}{2(t_2 - t_1)^2} ds + \int_{t_1}^{t_2} \frac{\alpha^2(t)}{32} dt + \int_{t_1}^{t_2} [\varphi + C\alpha^2(K + \mu_1) + \delta_1 \bar{u}_1] dt, \\ &\Lambda(t_1, t_2, \delta_1, \delta_2, \delta_3, \bar{u}_1) \\ &= \int_0^1 \frac{|\gamma'(s)|^4}{2(t_2 - t_1)^2} ds + \int_{t_1}^{t_2} \frac{\alpha^2(t)}{32} dt + \int_{t_1}^{t_2} [\varphi + C\alpha^2(K + \mu_2)] dt \\ &\quad + \int_{t_1}^{t_2} \left[\delta \sqrt{\frac{l\alpha - 1}{l-1}} \sqrt{\bar{u}_2 \delta_2} + \alpha^{\frac{3}{2}} \sqrt{n(l-1)\varphi \delta_1} \right] dt.\end{aligned}$$

Proof. Firstly, the estimate in Corollary 2.2 can be written as

$$\begin{aligned} & \frac{|\nabla u(x,t)|^2}{u^2(x,t)} - \alpha(t) \frac{u_t(x,t)}{u(x,t)} + \alpha(t) h(x,t) u^{l-1}(x,t) \\ & \leq \begin{cases} \alpha\varphi + C\alpha^2(K + \mu_1), & l \leq 1, \\ \alpha\varphi + C\alpha^2(K + \mu_2) + \delta \sqrt{\frac{l\alpha-1}{l-1}} \sqrt{\bar{u}_2\delta_2} \\ \quad + \alpha^{\frac{3}{2}} \sqrt{n(l-1)\varphi\delta_1}, & l > 1, \end{cases} \end{aligned} \quad (4.1)$$

where $\mu_1 = \sqrt{\bar{u}_1\delta_3} + \frac{\alpha(3-2l)-1}{\alpha-1}\bar{u}_1\delta_1 + \sqrt{(2-l)\bar{u}_1\delta_2}$ and $\mu_2 = (l-1)\bar{u}_2\delta_1 + \sqrt{\bar{u}_2\delta_3}$.

Now we only prove the conclusion for $l \leq 1$.

Define $l(s) = \log u(\gamma(s), (1-s)t_2 + st_1)$. Obviously, we infer that $l(0) = \log u(x_2, t_2)$ and $l(1) = \log u(x_1, t_1)$. Direct calculation shows

$$\begin{aligned} \frac{\partial l(s)}{\partial s} &= (t_2 - t_1) \left(\frac{\nabla u}{u} \frac{\gamma'(s)}{t_2 - t_1} - \frac{u_t}{u} \right) \\ &\leq (t_2 - t_1) \left[\frac{\nabla u}{u} \frac{\gamma'(s)}{t_2 - t_1} - \frac{1}{\alpha(t)} \frac{|\nabla u|^2}{u^2} - h(x,t) u^{l-1} + \varphi + C\alpha(K + \mu_1) \right] \\ &\leq \frac{\alpha(t)}{4} \frac{|\gamma'(s)|^2}{t_2 - t_1} + (t_2 - t_1)[\varphi + C\alpha(K + \mu) + \delta_1\bar{u}_1]. \end{aligned}$$

Integrating the above inequality over $\gamma(s)$, we obtain

$$\begin{aligned} \log \frac{u(x_1, t_1)}{u(x_2, t_2)} &= \int_0^1 \frac{\partial l(s)}{\partial s} ds \\ &\leq \int_0^1 \left[\frac{\alpha(t)}{4} \frac{|\gamma'(s)|^2}{t_2 - t_1} + (t_2 - t_1)[\varphi + C\alpha(K + \mu_1) + \delta_1\bar{u}_1] \right] ds \\ &\leq \int_0^1 \frac{|\gamma'(s)|^4}{2(t_2 - t_1)^2} ds + \int_{t_1}^{t_2} \frac{\alpha^2(t)}{32} dt \\ &\quad + \int_{t_1}^{t_2} [\varphi + C\alpha(K + \mu) + \delta_1\bar{u}_1] dt. \end{aligned}$$

The proof is complete.

We also derive an Harnack inequality for the equation (1.6). The proof is similar to Theorem 4.1, so we omit it.

THEOREM 4.2. *Let $(M^n, g(x,t))_{t \in [0,T]}$ be a complete solution to the Ricci flow (1.7). Suppose that $|\text{Ric}| \leq K$ for some $K > 0$, and all $(x,t) \in M^n \times [0,T]$. Assume that $u(x,t)$ is a positive solution to (1.6). Then for all $(x_1, t_1) \in M^n \times (0, T)$ and $(x_2, t_2) \in M^n \times (0, T)$ such that $t_1 < t_2$, we have*

$$u(x_2, t_2) \leq u(x_1, t_1)$$

$$\times \exp \left(\int_0^1 \frac{|\gamma'(s)|^4}{2(t_2 - t_1)^2} ds + \int_{t_1}^{t_2} \frac{\alpha^2(t)}{32} dt + \int_{t_1}^{t_2} [\varphi + C\alpha(K + |a|) + |a \log N|] dt \right),$$

where $N = \max_{M^n \times [0,T]} u$.

5. Application to heat equation

According to Theorem 2.1 and Theorem 3.1, we derive corresponding gradient estimates and Harnack inequalities to the heat equation along the Ricci flow.

THEOREM 5.1. *Let $(M^n, g(t))_{t \in [0, T]}$ be a complete solution to the Ricci flow (1.7). Assume that $|\text{Ric}(x, t)| \leq K$ for some $K > 0$ and all $t \in [0, T]$. Suppose that there exist three functions $\alpha(t)$, $\varphi(t)$ and $\gamma(t)$ satisfy the conditions (C1), (C2), (C3) and (C4).*

Given $x_0 \in M$ and $R > 0$, let $u(x, t)$ be a positive solution of the heat equation

$$u_t = \Delta u, \quad (5.1)$$

in the cube $B_{2R, T} := \{(x, t) | d(x, x_0, t) \leq 2R, 0 \leq t \leq T\}$, where c is a constant.

If $\frac{\gamma\alpha^4}{\alpha-1} \leq C_1$ for some constant C_1 , then

$$\frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} \leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) + \frac{Cn^2}{R^2\gamma} + \alpha\varphi.$$

Here $C = C(n, C_2)$ is a constant.

If $\frac{\gamma}{\alpha-1} \leq C_2$ for some constant C_2 , then

$$\frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} \leq C\alpha^2 \left(\frac{1}{R^2} + \frac{\sqrt{K}}{R} + K \right) + \frac{Cn^2\alpha^4}{R^2\gamma} + \alpha\varphi.$$

Here $C = C(n, C_2)$ is a constant.

COROLLARY 5.1. *Let $(M^n, g(t))_{t \in [0, T]}$ be a complete solution to the Ricci flow (1.7). Assume that $|\text{Ric}(x, t)| \leq K$ for some $K > 0$ and all $t \in [0, T]$. Given $x_0 \in M$ and $R > 0$, let $u(x, t)$ be a positive solution of the heat equation (5.1) in the cube $B_{2R, T} := \{(x, t) | d(x, x_0, t) \leq 2R, 0 \leq t \leq T\}$. Then the following special estimates are valid.*

1. *Li-Yau type:*

$$\alpha(t) = \text{constant}, \quad \varphi(t) = \frac{n}{t} + \frac{nK\alpha^2}{\alpha-1}, \quad \gamma(t) = t^\theta \quad \text{with} \quad 0 < \theta \leq 2.$$

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} &\leq C\alpha^2 \left[\frac{1}{R^2} (1 + \sqrt{K}R) + \frac{\alpha^2}{\alpha-1} \frac{1}{R^2} + K \right] \\ &\quad + \alpha\varphi + n^{\frac{3}{2}}\alpha^2 K. \end{aligned}$$

2. *Hamilton type:*

$$\alpha(t) = e^{2Kt}, \quad \varphi(t) = \frac{n}{t} e^{4Kt}, \quad \gamma(t) = t e^{2Kt}.$$

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} &\leq C\alpha^2 \left[\frac{1}{R^2}(1 + \sqrt{KR}) + K \right] + \frac{C\alpha^4}{R^2 t e^{2Kt}} \\ &\quad + \alpha\varphi + n^{\frac{3}{2}}\alpha^2 K. \end{aligned}$$

3. Li-Xu type:

$$\begin{aligned} \alpha(t) &= 1 + \frac{\sinh(Kt)\cosh(Kt) - Kt}{\sinh^2(Kt)}, \quad \varphi(t) = 2nK[1 + \coth(Kt)], \\ \gamma(t) &= \tanh(Kt). \end{aligned}$$

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} &\leq C\alpha^2 \left[\frac{1}{R^2}(1 + \sqrt{KR}) + K \right] + \frac{C}{R^2 \tanh(Kt)} \\ &\quad + \alpha\varphi + n^{\frac{3}{2}}\alpha^2 K. \end{aligned}$$

4. Linear Li-Xu type:

$$\alpha(t) = 1 + 2Kt, \quad \varphi(t) = \frac{n}{t} + nK(1 + 2Kt + \mu Kt), \quad \gamma(t) = Kt \quad \text{with} \quad \mu \geq \frac{1}{4}.$$

$$\begin{aligned} \frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} &\leq C\alpha^2 \left[\frac{1}{R^2}(1 + \sqrt{KR}) + K \right] + \frac{C\alpha^4}{R^2 Kt} \\ &\quad + \alpha\varphi + n^{\frac{3}{2}}\alpha^2 K. \end{aligned}$$

Let $R \rightarrow \infty$, a global estimate is derived.

COROLLARY 5.2. *Let $(M^n, g(t))_{t \in [0, T]}$ be a complete solution to the Ricci flow (1.7). Assume that $|\text{Ric}(x, t)| \leq K$ for some $K > 0$ and all $t \in [0, T]$. Suppose that there exist three functions $\alpha(t)$, $\varphi(t)$ and $\gamma(t)$ which satisfy the following conditions (C1), (C2), (C3) and (C4).*

Given $x_0 \in M$ and $R > 0$, let $u(x, t)$ be a positive solution of the heat equation (5.2) in the cube $M^n \times [0, T]$. Then

$$\frac{|\nabla u|^2}{u^2} - \alpha \frac{u_t}{u} \leq C\alpha^2 K + \alpha\varphi,$$

where $C = C(n, C_1, C_2)$ is a constant.

Using Theorem 4.1, we derive a Harnack inequality.

COROLLARY 5.3. (Harnack Inequality) *Let $(M^n, g(t))_{t \in [0, T]}$ be a complete solution to the Ricci flow (1.7). Suppose that $|\text{Ric}| \leq K$ for some $K > 0$, and all $(x, t) \in M^n \times [0, T]$. Assume that $u(x, t)$ is a positive solution to (5.1). Then for all $(x_1, t_1) \in M^n \times (0, T)$ and $(x_2, t_2) \in M^n \times (0, T)$ such that $t_1 < t_2$, we have*

$$u(x_2, t_2) \leq u(x_1, t_1)$$

$$\times \exp \left(\int_0^1 \frac{|\gamma'(s)|^4}{2(t_2-t_1)^2} ds + \int_{t_1}^{t_2} \frac{\alpha^2(t)}{32} dt + \int_{t_1}^{t_2} [\varphi + C\alpha K] dt \right),$$

where $C = C(n)$ and $\gamma(s)$ is a smooth curve connecting x_1 and x_2 with $\gamma(1) = x_1$ and $\gamma(0) = x_2$.

6. Appendix

We will check some special functions $\alpha(t) > 1$, $\varphi(t) > 0$ and $\gamma(t) > 0$ satisfy the following two systems

$$\begin{cases} \frac{2\varphi}{n} - 2\alpha K \geq (\frac{2\varphi}{n} - \alpha') \frac{1}{\alpha}, \\ \frac{2\varphi}{n} - \alpha' > 0, \\ \frac{\varphi^2}{n} + \alpha\varphi' \geq 0, \end{cases} \quad (6.1)$$

and

$$\begin{cases} \frac{\gamma'}{\gamma} - (\frac{2\varphi}{n} - \alpha') \frac{1}{\alpha} \leq 0, \\ \frac{\gamma\alpha^4}{\alpha-1} \leq C, \text{ or } \frac{\gamma}{\alpha-1} \leq C. \end{cases} \quad (6.2)$$

Besides, $\alpha(t)$ and $\gamma(t)$ are non-decreasing.

- (1) Let $\alpha(t) = 1 + 2Kt$, $\varphi(t) = \frac{n}{t} + nK(1 + 2Kt + \mu Kt)$ ($\mu \geq \frac{1}{4}$) and $\gamma(t) = Kt$.

One can has

$$\begin{aligned} \text{(i)} \quad & \frac{2\varphi}{n} - \alpha' = \frac{2}{t} + 2K(1 + 2Kt + \mu Kt) - 2K > 0, \\ \text{(ii)} \quad & \frac{\varphi^2}{n} + \alpha\varphi' = \frac{n}{t^2} + nK^2(1 + 2Kt + \mu Kt)^2 + \frac{2nK}{t}(1 + 2Kt + \mu Kt) \\ & + (1 + 2Kt)(-\frac{n}{t^2} + 2nK^2 + n\mu K^2) \\ & = nK^2(1 + 2Kt + \mu Kt)^2 + \frac{2nK}{t}(2Kt + \mu Kt) \\ & + (1 + 2Kt)(2nK^2 + n\mu K^2) > 0, \\ \text{(iii)} \quad & \frac{2\varphi}{n} - 2\alpha K - (\frac{2\varphi}{n} - \alpha') \frac{1}{\alpha} \\ & = \frac{2}{t} + 2K(1 + 2Kt + \mu Kt) - 2K(1 + 2Kt) \\ & - \left[\frac{2}{t} + 2K(1 + 2Kt + \mu Kt) - 2K \right] \cdot \frac{1}{1 + 2Kt} \\ & = \frac{4Kt(\mu K^2 t^2 - Kt + 1)}{t(1 + 2Kt)} \geq 0, \quad \text{for } \mu \geq \frac{1}{4}. \end{aligned}$$

Hence, $\alpha(t) = 1 + 2Kt$, $\varphi(t) = \frac{n}{t} + nK(1 + 2Kt + \mu Kt)$ ($0 < \mu \leq \frac{1}{4}$) satisfy system (6.1).

On the other hand, one has

$$\begin{aligned} & \frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} \\ &= \frac{1}{t} - \left(\frac{2}{t} + 2K(1 + 2Kt + \mu Kt) - 2K \right) \frac{1}{1 + 2Kt} \\ &= \frac{1}{t(1 + 2Kt)} [-(4K^2 + 2K\mu)t^2 + 2Kt - 1] \\ &= \frac{1}{t(1 + 2Kt)} [-(3K^2 + 2K\mu)t^2 - (Kt - 1)^2] \\ &\leq 0, \quad \text{for } t \geq 0, \end{aligned}$$

and $\frac{\gamma}{\alpha-1} = \frac{1}{2}$. So, (6.2) is also satisfied.

- (2) $\alpha(t) = e^{2Kt}$, $\varphi(t) = \frac{n}{t}e^{4Kt}$ and $\gamma(t) = te^{2Kt}$, where ($0 < Kt \leq 1$). Direct calculation gives

$$\begin{aligned} \text{(i)} \quad & \frac{2\varphi}{n} - \alpha' = \frac{2}{t}e^{2Kt}(e^{2Kt} - Kt) > 0, \\ \text{(ii)} \quad & \frac{\varphi^2}{n} + \alpha\varphi' = \frac{n}{t^2}e^{6Kt}(e^{2Kt} - 1 + 4Kt) > 0, \\ \text{(iii)} \quad & \frac{2\varphi}{n} - 2\alpha K - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} = \frac{2}{t}e^{4Kt} - 2Ke^{2Kt} - \frac{2}{t}e^{2Kt} + 2K \\ &= (e^{2Kt} - 1)\left(\frac{2}{t}e^{2Kt} - 2K\right) \geq 0. \end{aligned}$$

Hence, $\alpha(t) = e^{2Kt}$ and $\varphi(t) = \frac{n}{t}e^{4Kt}$ satisfy system (6.1).

Besides, we have

$$\begin{aligned} & \frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} \\ &= \frac{1+2Kt}{t} - \left(\frac{2}{t}e^{2Kt} - 2K \right) \\ &= \frac{1}{t}(1+4Kt-2e^{2Kt}) \\ &\leq 0, \quad \text{for } t \geq 0, \end{aligned}$$

and as $t \rightarrow 0^+$, $\frac{\gamma}{\alpha-1} = \frac{te^{2Kt}}{e^{2Kt}-1} \rightarrow \frac{1}{2K}$. This implies $\frac{\gamma}{\alpha-1} \leq C$. So, (6.2) is also satisfied.

- (3) $\alpha(t) = 1 + \frac{\sinh(Kt)\cosh(Kt)-Kt}{\sinh^2(Kt)}$, $\varphi(t) = 2nK[1 + \coth(Kt)]$ and $\gamma(t) = \tanh(Kt)$. Direct calculation gives

$$\text{(i)} \quad \frac{2\varphi}{n} - \alpha' = 4K[1 + \coth(Kt)] - 2K + 2K\coth^2(Kt) - \frac{2K^2t}{\sinh^2(Kt)} \coth(Kt),$$

$$\begin{aligned}
&= 2K + 2K(1+\alpha) \coth(Kt) > 0, \\
\text{(ii)} \quad &\alpha \left(\frac{2\varphi}{n} - 2\alpha K \right) - \left(\frac{2\varphi}{n} - \alpha' \right) \\
&= 4K\alpha[1 + \coth(Kt)] - 2K\alpha^2 - [2K + 2K(1+\alpha) \coth(Kt)] \\
&= 2K\alpha \left[1 + \coth(Kt) + \frac{Kt}{\sinh^2(Kt)} \right] - [2K + 2K(1+\alpha) \coth(Kt)] \\
&= 2K(\alpha-1) \frac{Kt}{\sinh^2(Kt)} > 0, \\
\text{(iii)} \quad &\frac{\varphi^2}{n} + \alpha\varphi' = \frac{2nK^2}{\sinh^2(Kt)} \left[2(1 + \coth(Kt))^2 \sinh^2(Kt) - \alpha \right] \\
&= \frac{2nK^2}{\sinh^2(Kt)} \left[2e^{2Kt} - 1 - \frac{e^{4Kt} - 1 - 4Kte^{2Kt}}{(e^{2Kt} - 1)^2} \right] \\
&= \frac{4nK^2 e^{2Kt}}{(e^{2Kt} - 1)^2 \sinh^2(Kt)} \left[e^{4Kt} - 3e^{2Kt} + 2 + 4Kt \right].
\end{aligned}$$

Let $f(x) = e^{4x} - 3e^{2x} + 2 + 4x$ with $x \leq 0$. Obviously, $f(0) = 0$ and

$$f'(x) = 4e^{4x} - 6e^{2x} + 4 > 0.$$

Then we get $f(x) > 0$ for $x > 0$. Hence, we have

$$\begin{aligned}
&\left(\frac{2\varphi}{n} - \alpha' \right) \varphi + \alpha\varphi' + \alpha'\varphi - \frac{\varphi^2}{n} \\
&= \frac{4nK^2 e^{2Kt}}{(e^{2Kt} - 1)^2 \sinh^2(Kt)} \left[e^{4Kt} - 3e^{2Kt} + 2 + 4Kt \right] > 0.
\end{aligned}$$

Hence, $\alpha(t) = 1 + \frac{\sinh(Kt)\cosh(Kt)-Kt}{\sinh^2(Kt)}$ and $\varphi(t) = 2nK[1 + \coth(Kt)]$ satisfy system (6.1).

On the other hand, as $t \rightarrow 0$, we have $\frac{\gamma\alpha^4}{\alpha-1} \rightarrow 2$; $\frac{\gamma\alpha^4}{\alpha-1} \rightarrow 1$ for $t \rightarrow \infty$. These imply $\frac{\gamma\alpha^4}{\alpha-1} \leq C$, here C is a universal constant.

Besides, we have

$$\begin{aligned}
&\frac{\gamma}{\gamma} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} \\
&= \frac{1}{\alpha} \left[\frac{K\alpha}{\sinh(Kt)\cosh(Kt)} - 2K - 2K(1+\alpha) \coth(Kt) \right] \\
&= \frac{1}{\alpha} \left[\frac{K}{\sinh(Kt)\cosh(Kt)} [\alpha - 2(1+\alpha) \cosh^2(Kt)] - 2K \right] \\
&= \frac{1}{\alpha} \left[\frac{K}{\sinh(Kt)} [\alpha(1 - 2\cosh(Kt)) - 2\cosh(Kt)] - 2K \right] \\
&\leq 0, \quad \text{for } t \geq 0.
\end{aligned}$$

So, (6.2) is also satisfied.

- (4) $\alpha(t) = \text{constant}$, $\varphi(t) = \frac{\alpha n}{t} + \frac{nK\alpha^2}{\alpha-1}$ and $\gamma(t) = t^\theta$ with $0 < \theta \leq 2$. Direct calculation gives

$$\begin{aligned} \text{(i)} \quad & \frac{2\varphi}{n} - \alpha' = \frac{2}{n} \left[\frac{\alpha n}{t} + \frac{nK\alpha^2}{\alpha-1} \right] > 0, \\ \text{(ii)} \quad & \frac{\varphi^2}{n} + \alpha\varphi' = \frac{n\alpha^2}{t^2} + \frac{n^2 K^2 \alpha^4}{n(\alpha-1)^2} + \frac{2nK\alpha^2}{(\alpha-1)t} - \frac{n\alpha^2}{t^2} > 0, \\ \text{(iii)} \quad & \left(\frac{2\varphi}{n} - 2\alpha K \right) - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} \\ &= \frac{2\varphi}{n\alpha} (\alpha-1) - 2K\alpha \\ &\geq \frac{2}{n\alpha} (\alpha-1) \frac{nK\alpha^2}{\alpha-1} - 2K\alpha = 0. \end{aligned}$$

Hence, $\alpha(t) = \text{constant}$, and $\varphi(t) = \frac{\alpha n}{t} + \frac{nK\alpha^2}{\alpha-1}$ which satisfy system (6.1).

On the other hand, we have

$$\begin{aligned} \frac{\gamma'}{\gamma} - \left(\frac{2\varphi}{n} - \alpha' \right) \frac{1}{\alpha} &= \frac{\theta}{t} - \frac{2}{t} - \frac{2K\alpha}{\alpha-1} \\ &\leq 0, \quad \text{for } t \geq 0 \quad \text{and } 0 < \theta \leq 2. \end{aligned}$$

So, (6.2) is also satisfied.

Acknowledgement. We are grateful to Professor Jiayu Li for his encouragement. We also thank Professor Qi S Zhang for introduction of this problem in the summer course. The authors would like to thank the referee for valuable comments and suggestions. This work was supported by the University outstanding young talent support project of Anhui province in 2017 (gxyq2017048, GXYQ2017070), the Universities Natural Science Foundation of Anhui Province (KJ2017A937, KJ2016A506, KJ2017A454, KJ2019A0712, KJ2019A0713), and the Natural Science Foundation of Anhui Province (1908085QA04, 1708085MA16, 1608085QA12).

REFERENCES

- [1] ABIMBOLA ABOLARINWA, *Gradient estimates for a nonlinear elliptic equation on complete noncompact Riemannian manifold*, Journal of Mathematical Inequalities 12(2), (2018), 391–402.
- [2] M. BAILESTEANU, X. D. CAO, A. PULEMOTOV, *Gradient estimates for the heat equation under the Ricci flow*, J. Funct. Anal., 258 (2010), 3517–3542.
- [3] E. CALABI, *An extension of E. Hopf's maximum principle with an application to Riemannian geometry*, Duke Math. J. 25 (1958), 45–56, MR 19,1056e Zbl 0079.11801.
- [4] H. CAO, M. ZHU, *Aronson-Bénilan estimates for the porous medium equation under the Ricci flow*, Journal De Mathématiques Pures Et Appliquées, 2015, 104(4):90–94.
- [5] X. D. CAO, B. F. LJUNGBERG, B. W. LIU, *Differential Harnack estimates for a nonlinear heat equation*, J. Funct. Anal., (2103): 1–19.
- [6] D. G. CHEN, C. W. XIONG, *Gradient estimates for doubly nonlinear diffusion equations*, Nonlinear Anal. 112 (2015), 156–164.

- [7] B. CHOW AND D. KNOPF, *The Ricci flow: An introduction, mathematical Surveys and Monographs 110*, American Society, Providence, RI, 2004, MR 2005e:53101 Zbl 1086.53085.
- [8] R. S. HAMILTON, *A matrix Harnack estimates for the heat equation*, Comm. Anal. Geom., 1 (1993), 113–126.
- [9] R. S. HAMILTON, *Three manifolds with positive Ricci curvature*, J. Differential Geom. 17:2 (1982): 255–306, MR 84a:53050 Zbl 0504.53034.
- [10] R. S. HAMILTON, *The formation of singularities in the Ricci flow*, pp. 7–136 in *Surveys in differential geometry, II* (Cambridge, MA, 1993), edited by S. T. Yau, International, Cambridge, MA, 1995., MR 97e:53075 Zbl 0867.53030.
- [11] G. Y. HUANG, Z. J. HUANG, H. Z. LI, *Gradient estimates and differential Harnack inequalities for a nonlinear parabolic equation on Riemannian manifolds*, Ann Glob Anal Geom, (2013) 43: 209–232.
- [12] S. KUANG, Q. S. ZHANG, *A gradient estimate for all positive solutions of the conjugate heat equation under Ricci flow*, J. Funct. Anal. 255:4 (2008), 1008–1023, MR 2433960 Zbl 1146.58017.
- [13] H. LI, H. BAI, G. ZHANG, *Hamilton’s gradient estimates for fast diffusion equation under the Ricci flow*, J. Math. Anal. Appl., 444 (2016) 1372–1379.
- [14] P. LI, S. T. YAU, *On the parabolic kernel of the Schrödinger operator*, Acta Math., 156 (1986), 153–201, MR 87f:58156.
- [15] J. Y. LI, *Gradient estimates and Harnack inequalities for nonlinear parabolic and nonlinear elliptic equations on Riemannian manifolds*, J. Funct. Anal., 100 (1991), 233–256.
- [16] J. Y. LI, *Gradient estimate for the heat kernel of a complete Riemannian manifold and its applications*, J. Funct. Anal., 97:2 (1991), 293–310, MR 92f:58174 Zbl 0724.58064.
- [17] J. LI AND X. XU, *Differential Harnack inequalities on Riemannian manifolds I: Linear heat equation*, Adv. in Math., 226 (2011), 4456–4491.
- [18] S. P. LIU, *Gradient estimates for solutions of the heat equation under flow*, Pacific J. of Math., 243 (1), (2009), 165–179., MR 2010g:53122 Zbl 1180.58017.
- [19] Y. LI, X. R. ZHU, *Harnack estimates for a heat-type equation under the Ricci flow*, J. Differential Equations 260 (2016), 3270–3301.
- [20] P. LU, L. NI, J. L. VÁZQUEZ AND C. VILLANI, *Local Aronson-Bénilan estimates and entropy formulae for porous medium and fast diffusion equations on manifolds*, J. Math. Pures Appl. 91 (2009), 1–19.
- [21] L. MA, *Gradient estimate for a sample elliptic equation on non-compact Riemannian manifolds*, J. Funct. Anal., 241 (2006), 374–382, MR 2264255(2007e:53034).
- [22] G. PERELMAN, *The entropy formula for the Ricci flow and its geometric applications*, preprint, 2002, arXiv math.DG/0211159.
- [23] G. PERELMAN, *Ricci flow with surgery on three manifolds*, preprint, 2003, arXiv math.DG/0203109.
- [24] W. X. SHI, *Deforming the metric on complete Riemannian manifolds*, J. Differential Geom., 30: 1 (1989), 223–301, MR 90i:58202 Zbl 6076.53044.
- [25] B. H. SONG, *The Harnack Estimate for a Nonlinear Parabolic Equation under the Ricci Flow*, Acta Mathematica Sinica, English Series 27(10),(2011): 1935C1940 DOI: 10.1007/s10114-011-0074-z.
- [26] J. SUN, *Gradient estimates for positive solutions of the heat equation under geometric flow*, Pacific J. Math. 253, 489–510 (2011).
- [27] H. J. SUN, *Higher Eigenvalue Estimates on Riemannian Manifolds with Ricci Curvature Bounded Below*, Acta Math. Sinica (Chin. Ser.) 49 (2006), 3, 539–548.
- [28] W. WANG, P. ZHANG, *Some Gradient Estimates and Harnack Inequalities for Nonlinear Parabolic Equations on Riemannian Manifolds*, Mathematische Nachrichten, (2017) 290,11-12,1905–1917.
- [29] J. Y. WU, *Li-Yau type estimates for a nonlinear parabolic equation on complete manifolds*, J. Math. Anal. Appl., 369 (2010), 400–407.
- [30] Y. Y. YANG, *Gradient estimate for a nonlinear parabolic equation on Riemannian manifold*, Proc. Amer. Math. Soc. 136 (2008),4095–4102.
- [31] Y. Y. YANG, *Gradient estimates for the equation $\Delta u + cu^\alpha = 0$ on Riemannian manifolds*, Acta. Math. Sin. 26(B), 1177–1182, (2010).

- [32] L. ZHAO, S. FANG, *Gradient estimates for a nonlinear liichenrowicz equation under generalized geometric flow on complete noncompact manifolds*, Pacific J. Math. 285 (1) (2016):243–256.
- [33] X. B. ZHU, *Gradient estimates and Liouville theorems for nonlinear parabolic equations on noncompact Riemannian manifolds*, Nonlinear Analysis, 74 (2011), 5141–5146.

(Received April 14, 2019)

Wen Wang

School of Mathematics and Statistics

Hefei Normal University

Hefei 230601, P.R.China

School of mathematical Science

University of Science and Technology of China

Hefei 230026, China

e-mail: wwen2014@mail.ustc.edu.cn

Hui Zhou

School of Mathematics and Statistics

Hefei Normal University

Hefei 230601, P.R.China

School of mathematical Science

University of Science and Technology of China

Hefei 230026, China

e-mail: zhouchui0309@126.com

Rulong Xie

School of Mathematics and Statistics

Chaochu University

Chaochu, 238000, P. R. China

e-mail: xierl@ustc.edu.cn

Songting Yin

Department of Mathematics and Computer Science

Tongling University

Tongling, 244000, P. R. China

e-mail: yst419@163.com