

NEW GENERALIZED RIEMANN—LOUVILLE FRACTIONAL INTEGRAL INEQUALITIES FOR CONVEX FUNCTIONS

PSHTIWAN OTHMAN MOHAMMED

(Communicated by J. Pečarić)

Abstract. In the literature, the right-side of Hermite–Hadamard’s inequality is called trapezoid type inequality. In this paper, we obtain new integral inequalities of trapezoid type for convex functions involving generalized Riemann–Liouville fractional integrals (ψ -Riemann–Liouville fractional integrals). Our obtained inequalities generalize some recent classical integral inequalities and Riemann–Liouville fractional integral inequalities which are established in earlier works.

1. Introduction

A function $g : \mathcal{I} \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is said to be convex on the interval \mathcal{I} , if the inequality

$$g(\eta x + (1 - \eta)y) \leq \eta g(x) + (1 - \eta)g(y) \quad (1)$$

holds for all $x, y \in \mathcal{I}$ and $\eta \in [0, 1]$. We say that g is concave if $-g$ is convex.

For convex functions (1), many equalities and inequalities have been established by many authors; such as Ostrowski type inequality [7], Hardy type inequality [2], Olsen type inequality [8], Gagliardo-Nirenberg type inequality [28], and midpoint and trapezoid type inequalities [21, 22]. But the most important inequality is the Hermite–Hadamard type inequality [3], which is defined by:

$$\begin{aligned} g\left(\frac{u+v}{2}\right) &\leq \frac{1}{v-u} \int_u^v g(x) dx \\ &\leq \frac{g(u) + g(v)}{2}, \end{aligned} \quad (2)$$

where $g : \mathcal{I} \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is assumed to be a convex function on \mathcal{I} with $u, v \in \mathcal{I}$ and $u < v$.

A number of mathematicians in the field of applied and pure mathematics have dedicated their efforts to generalize, refine, counterpart, and extend the Hermite–Hadamard inequality (2) for different classes of convex functions and mappings. For more recent results obtained in view of inequality (2); we refer our readers to [3, 32, 11, 4, 19].

Mathematics subject classification (2020): 26D07, 26D15, 26D10, 26A33.

Keywords and phrases: Riemann–Liouville fractional integral, convex function, Hermite–Hadamard inequality, convex functions, trapezoid formula.

In [13], Sarikaya et al. obtained the following Hermite–Hadamard's inequalities in fractional integral form:

$$\begin{aligned} g\left(\frac{u+v}{2}\right) &\leq \frac{\Gamma(\mu+2)}{2(v-u)^\mu} [I_{u^+}^\mu g(v) + I_{v^-}^\mu g(u)] \\ &\leq \frac{g(u) + g(v)}{2}, \end{aligned} \quad (3)$$

where $g : [u, v] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is assumed to be a positive convex function on $[u, v]$, $g \in L_1[u, v]$ with $u < v$, and $I_{u^+}^\mu$ and $I_{v^-}^\mu$ are the left-sided and right-sided Riemann–Liouville fractional integrals of order $\mu > 0$, these are respectively defined as [10]:

$$\begin{aligned} I_{u^+}^\mu g(x) &= \frac{1}{\Gamma(\mu)} \int_u^x (x-t)^{\mu-1} g(t) dt, \quad x > u, \\ I_{v^-}^\mu g(x) &= \frac{1}{\Gamma(\mu)} \int_x^v (t-x)^{\mu-1} g(t) dt, \quad x < v. \end{aligned} \quad (4)$$

If we take $\mu = 1$ in (3) we obtain (2), so it is clear that inequality (3) is a generalization of Hermite–Hadamard inequality (2).

In view of inequality (3), many inequalities have been established for convex [27, 6], h -convex [13], MT -convex [26, 14], m -convex [31], (s, m) -convex [31], (α, m) -convex [24, 29], F -convex [21], invex and preinvex [9, 32, 12] functions. Furthermore, there are many related inequalities for other kinds of convex functions and fractional operators; so it is important that the readers of this article visit [11, 4, 1, 17, 18, 20, 23].

Let us now state the definition of generalized Riemann–Liouville fractional integrals:

DEFINITION 1. Let $(u, v) \subseteq (-\infty, \infty)$ be a finite or infinite interval of the real-axis \mathbb{R} and $\mu > 0$. Let $\psi(x)$ be an increasing and positive monotone function on the interval (u, v) with a continuous derivative $\psi'(x)$ on the interval (u, v) . Then the left and right-sided ψ -Riemann–Liouville fractional integrals of a function g with respect to another function $\psi(x)$ on $[u, v]$ are defined by [30]:

$$\begin{aligned} I_{u^+}^{\mu:\psi} g(x) &= \frac{1}{\Gamma(\mu)} \int_u^x \psi'(t)(\psi(x) - \psi(t))^{\mu-1} g(t) dt, \\ I_{v^-}^{\mu:\psi} g(x) &= \frac{1}{\Gamma(\mu)} \int_x^v \psi'(t)(\psi(t) - \psi(x))^{\mu-1} g(t) dt. \end{aligned} \quad (5)$$

It is important to be noted that if we set $\psi(x) = x$ in (5), then ψ -Riemann–Liouville fractional integral reduces to Riemann–Liouville fractional integral (4).

Based on this definition, we obtain several inequalities of trapezoid type for convex functions in the present paper.

2. The main results

Our main results depend on the following lemma:

LEMMA 1. Let $g : [u, v] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function and $g'' \in L_1[u, v]$ with $0 \leq u < v$. If $\psi(x)$ is an increasing and positive monotone function on $(u, v]$ and its derivative $\psi'(x)$ is continuous on (u, v) , then for $t_1, t_2, \mu \in (0, 1)$ we have

$$\begin{aligned} \sigma_{\mu, \psi}(g; u, v) &= \frac{1}{2(\mu + 1)(v - u)^\mu} \\ &\times \left(\int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \left[(v - u)^\mu (v - \psi(t_1)) - (v - \psi(t_1))^{1+\mu} \right] \psi'(t_1) (g'' \circ \psi)(t_1) dt_1 \right. \\ &\quad \left. + \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \left[(v - u)^\mu (\psi(t_2) - u) - (\psi(t_2) - u)^{1+\mu} \right] \psi'(t_2) (g'' \circ \psi)(t_2) dt_2 \right), \end{aligned} \quad (6)$$

where

$$\begin{aligned} \sigma_{\mu, \psi}(g; u, v) &= \frac{g(u) + g(v)}{2} \\ &\quad - \frac{\Gamma(\mu + 1)}{2(v - u)^\mu} [I_{u^+}^{\mu: \psi}(g \circ \psi)(\psi^{-1}(v)) + I_{v^-}^{\mu: \psi}(g \circ \psi)(\psi^{-1}(u))]. \end{aligned}$$

Proof. From Definition 1 we have

$$\begin{aligned} \bar{h}_1 &:= \frac{\Gamma(\mu + 1)}{2(v - u)^\mu} I_{u^+}^{\mu: \psi}(g \circ \psi)(\psi^{-1}(v)) \\ &= \frac{\mu}{2(v - u)^\mu} \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \psi'(t_1) (v - \psi(t_1))^{\mu-1} (g \circ \psi)(t_1) dt_1 \\ &= -\frac{1}{2(v - u)^\mu} \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} (g \circ \psi)(t_1) d(v - \psi(t_1))^\mu. \end{aligned}$$

Integrating by parts twice, we have

$$\begin{aligned} \bar{h}_1 &= \frac{1}{2} g(u) + \frac{1}{2(\mu + 1)(v - u)^\mu} \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \psi'(t_1) (v - \psi(t_1))^\mu (g' \circ \psi)(t_1) dt_1 \\ &= \frac{1}{2} g(u) + \frac{v - u}{2(\mu + 1)} g'(u) \\ &\quad + \frac{1}{2(\mu + 1)(v - u)^\mu} \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \psi'(t_1) (v - \psi(t_1))^{1+\mu} (g'' \circ \psi)(t_1) dt_1. \end{aligned} \quad (7)$$

Similarly,

$$\begin{aligned}\bar{h}_2 &:= \frac{\Gamma(\mu+1)}{2(v-u)^\mu} I_{v^-}^{\mu:\psi}(g \circ \psi)(\psi^{-1}(u)) \\ &= \frac{1}{2}g(v) - \frac{v-u}{2(\mu+1)}g'(v) \\ &\quad + \frac{1}{2(\mu+1)(v-u)^\mu} \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \psi'(t_2)(\psi(t_2)-u)^{\mu+1}(g'' \circ \psi)(t_2) dt_2.\end{aligned}\quad (8)$$

Integrating by parts, we have

$$\begin{aligned}\bar{h}_3 &:= \frac{1}{2(\mu+1)} \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \psi'(t_1)(v-\psi(t_1))(g'' \circ \psi)(t_1) dt_1 \\ &= -\frac{v-u}{2(\mu+1)}g'(u) + \frac{g(v)-g(u)}{2(\mu+1)},\end{aligned}\quad (9)$$

and

$$\begin{aligned}\bar{h}_4 &:= \frac{1}{2(\mu+1)} \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \psi'(t_2)(\psi(t_2)-u)(g'' \circ \psi)(t_2) dt_1 \\ &= \frac{v-u}{2(\mu+1)}g'(v) - \frac{g(v)-g(u)}{2(\mu+1)}.\end{aligned}\quad (10)$$

It follows from (7)–(10) that

$$\begin{aligned}&\frac{g(u)+g(v)}{2} - (\bar{h}_1 + \bar{h}_2 + \bar{h}_3 + \bar{h}_4) \\ &= \frac{1}{2(\mu+1)(v-u)^\mu} \left(\int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \left[(v-u)^\mu (v-\psi(t_1)) - (v-\psi(t_1))^{\mu+1} \right] \psi'(t_1)(g'' \circ \psi)(t_1) dt_1 \right. \\ &\quad \left. + \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \left[(v-u)^\mu (\psi(t_2)-u) - (\psi(t_2)-u)^{\mu+1} \right] \psi'(t_2)(g'' \circ \psi)(t_2) dt_2 \right),\end{aligned}$$

and this completes the proof of Lemma 1. \square

REMARK 1. By letting $t_1 = t_2 = t$, then Lemma 1 becomes

$$\begin{aligned}\sigma_{\mu,\psi}(g; u, v) &= \frac{1}{2(\mu+1)(v-u)^\mu} \\ &\times \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \left[(v-u)^{\mu+1} - (v-\psi(t))^{\mu+1} - (\psi(t)-u)^{\mu+1} \right] \psi'(t)(g'' \circ \psi)(t) dt.\end{aligned}\quad (11)$$

COROLLARY 1. With the similar assumptions of Lemma 1 if $\psi(x) = x$, we have

$$\begin{aligned} & \frac{g(u) + g(v)}{2} - \frac{\Gamma(\mu + 1)}{2(v-u)^\mu} [I_{u^+}^\mu g(v) + I_v^\mu g(u)] \\ &= \frac{(v-u)^2}{2(\mu+1)} \int_0^1 t(1-t^\mu) \left[g''(tu + (1-t)v) + g''((1-t)u + tv) \right] dt, \end{aligned}$$

which is obtained by Dragomir et al. [5].

THEOREM 1. Let $g : [u, v] \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function and $g'' \in L_1[u, v]$ with $0 \leq u < v$. Suppose that $|g''|$ is convex on $[u, v]$, $\psi(x)$ is an increasing and positive monotone function on (u, v) and its derivative $\psi'(x)$ is continuous on (u, v) , then for $\mu \in (0, 1)$ we have

$$\begin{aligned} |\sigma_{\mu, \psi}(g; u, v)| &\leq \frac{(v-u)^2}{2(\mu+1)(\mu+2)} \left(\frac{|g''(u)| + |g''(v)|}{2} \right) \\ &\leq \frac{(v-u)^2}{\mu+1} \mathcal{B}(2, \mu+1) \left(\frac{|g''(u)| + |g''(v)|}{2} \right), \end{aligned} \quad (12)$$

where \mathcal{B} denotes the well-known Euler Beta function.

Proof. By using Lemma 1 with $x_1 = \frac{v-\psi(t_1)}{v-u}$ and $x_2 = \frac{\psi(t_2)-u}{v-u}$, we get

$$\begin{aligned} \sigma_{\mu, \psi}(g; u, v) &= \frac{(v-u)^2}{2(\mu+1)} \left(\int_0^1 \left(x_1 - x_1^{\mu+1} \right) g''(x_1 u + (1-x_1)v) dx_1 \right. \\ &\quad \left. + \int_0^1 \left(x_2 - x_2^{\mu+1} \right) g''((1-x_2)u + x_2 v) dx_2 \right). \end{aligned} \quad (13)$$

Let $t = x_1 = x_2$, then (13) becomes

$$\begin{aligned} & \sigma_{\mu, \psi}(g; u, v) \\ &= \frac{(v-u)^2}{2(\mu+1)} \left[\int_0^1 (t - t^{\mu+1}) \left(g''(tu + (1-t)v) + g''((1-t)u + tv) \right) dt \right]. \end{aligned} \quad (14)$$

Using properties of modulus and the convexity of $|g''|$, we get

$$\begin{aligned} |\sigma_{\mu, \psi}(g; u, v)| &\leq \frac{(v-u)^2}{2(\mu+1)} \left[\int_0^1 |t - t^{\mu+1}| \left(|g''(tu + (1-t)v)| + |g''((1-t)u + tv)| \right) dt \right] \\ &\leq \frac{(v-u)^2}{2(\mu+1)} \left(|g''(u)| \int_0^1 t (t - t^{\mu+1}) dt + |g''(v)| \int_0^1 (1-t) (t - t^{\mu+1}) dt \right. \\ &\quad \left. + |g''(v)| \int_0^1 (t - t^{\mu+1}) dt + |g''(u)| \int_0^1 (1-t) (t - t^{\mu+1}) dt \right) \\ &= \frac{(v-u)^2}{2(\mu+1)(\mu+2)} \left(\frac{|g''(u)| + |g''(v)|}{2} \right), \end{aligned}$$

which proves the first inequality of (12).

We know for each $x_1, x_2 \in [0, 1]$ and $\mu \in (0, 1)$ that $|x_1^\mu + x_2^\mu| \leq |x_1 + x_2|^\mu$, so we have

$$\int_0^1 t^2 (1-t^\mu) dt + \int_0^1 t(1-t)(1-t^\mu) dt \leq \int_0^1 t(1-t)^\mu dt = \mathcal{B}(2, \mu+1),$$

which proves the second inequality of (12). Thus the proof of Theorem 1 is completed. \square

COROLLARY 2. *With the similar assumptions of Theorem 1*

1. if $\psi(x) = x$, we have

$$\begin{aligned} & \left| \frac{g(u) + g(v)}{2} - \frac{\Gamma(\mu+2)}{2(v-u)^\mu} [I_{u^+}^\mu g(v) + I_{v^-}^\mu g(u)] \right| \\ & \leq \frac{(v-u)^2}{2(\mu+1)(\mu+2)} \left(\frac{|g''(u)| + |g''(v)|}{2} \right) \\ & \leq \frac{(v-u)^2}{\mu+1} \mathcal{B}(2, \mu+1) \left(\frac{|g''(u)| + |g''(v)|}{2} \right), \end{aligned}$$

this is obtained by Dragomir et al. [5].

2. if $\psi(x) = x$ and $\mu = 1$, we have

$$\left| \frac{g(u) + g(v)}{2} - \frac{1}{v-u} \int_u^v g(x) dx \right| \leq \frac{(v-u)^2}{12} \left(\frac{|g''(u)| + |g''(v)|}{2} \right),$$

which is obtained by Ozdemir et al. [25].

REMARK 2. Let the assumptions of Theorem 1 be hold

1. for $\frac{1}{p} + \frac{1}{q} = 1$, $p > 1$, we have

$$|\sigma_{\mu, \psi}(g; u, v)| \leq \frac{(v-u)^2}{\mu+1} \mathcal{B}^{\frac{1}{p}}(p+1, \mu p+1) \left(\frac{|g''(u)|^q + |g''(v)|^q}{2} \right)^{\frac{1}{q}},$$

such that $|g''|^q$ is convex function on $[u, v]$.

2. for $q > 1$, we have

$$\begin{aligned} |\sigma_{\mu, \psi}(g; u, v)| & \leq \frac{\mu(v-u)^2}{4(\mu+1)(\mu+2)} \left[\left(\frac{2\mu+4}{3\mu+9} |g''(u)|^q + \frac{\mu+5}{3\mu+9} |g''(v)|^q \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\frac{\mu+5}{3\mu+9} |g''(u)|^q + \frac{2\mu+4}{3\mu+9} |g''(v)|^q \right)^{\frac{1}{q}} \right], \end{aligned}$$

such that $|g''|^q$ is a convex function on $[u, v]$.

3. for $\frac{1}{p} + \frac{1}{q} = 1$, $p > 1$

$$|\sigma_{\mu,\psi}(g;u,v)| \leq \frac{(v-u)^2}{\mu+1} \mathcal{B}^{\frac{1}{p}}(p+1, \mu p+1) \left| g''\left(\frac{a+b}{2}\right) \right|,$$

such that $|g''|^q$ is concave function on $[u, v]$.

4. for $\frac{1}{p} + \frac{1}{q} = 1$, $p \geq 1$

$$\begin{aligned} & |\sigma_{\mu,\psi}(g;u,v)| \\ & \leq \frac{\mu(v-u)^2}{4(\mu+1)(\mu+2)} \left[\left| g''\left(\frac{2\mu+4}{3\mu+9}a + \frac{\mu+5}{3\mu+9}b\right) \right| + \left| g''\left(\frac{\mu+5}{3\mu+9}a + \frac{2\mu+4}{3\mu+9}b\right) \right| \right], \end{aligned}$$

such that $|g''|^q$ is concave function on $[u, v]$.

Proof. By making use of the equality (14) and the same method used for Theorem 3–6 of [5], we can easily obtain the items 1–4, respectively. \square

THEOREM 2. *With the assumptions of Theorem 1, we have*

$$\begin{aligned} |\sigma_{\mu,\psi}(g;u,v)| & \leq \frac{(v-u)|g'(v)-g'(u)|}{2(\mu+1)} \\ & + \frac{(v-u)^2}{2(\mu+1)} \left(\frac{\mu+4}{(\mu+2)(\mu+3)} |g''(u)| + \frac{2\mu+5}{(\mu+2)(\mu+3)} |g''(v)| \right). \quad (15) \end{aligned}$$

Proof. Making use of the equality (11), we have

$$\begin{aligned} & |\sigma_{\mu,\psi}(g;u,v)| \leq \frac{1}{2(\mu+1)(v-u)^\mu} \left| \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} (v-u)^{\mu+1} \psi'(t) (g'' \circ \psi)(t) dt \right| \\ & + \frac{1}{2(\mu+1)(v-u)^\mu} \left| \int_{\psi^{-1}(u)}^{\psi^{-1}(v)} \left[(v-\psi(t))^{\mu+1} + (\psi(t)-u)^{\mu+1} \right] \psi'(t) (g'' \circ \psi)(t) dt \right| \\ & := \omega_1 + \omega_2. \quad (16) \end{aligned}$$

By using a simple integrating by parts, we get

$$\omega_1 = \frac{(v-u)|g'(v)-g'(u)|}{2(\mu+1)}. \quad (17)$$

By changing the variable $x = \frac{v-\psi(t)}{v-u}$ and convexity of $|g''|$, we have

$$\begin{aligned} \omega_2 & = \frac{(v-u)^2}{2(\mu+1)} \int_0^1 (x^{\mu+1} + (1-x)^{\mu+1}) |g''(xa + (1-x)b)| dx \\ & = \frac{(v-u)^2}{2(\mu+1)} \left(\frac{\mu+4}{(\mu+2)(\mu+3)} |g''(u)| + \frac{2\mu+5}{(\mu+2)(\mu+3)} |g''(v)| \right). \quad (18) \end{aligned}$$

Hence, the inequalities (16)–(18) rearrange proof of Theorem 2. \square

3. Conclusion

In this paper, we established some new inequalities of trapezoid type for convex functions involving ψ -Riemann–Liouville fractional integrals. Corollary 1–2 and Remark 2 show that our results generalize and extend the obtained results in [5, 25].

REFERENCES

- [1] D. BALEANU, P. O. MOHAMMED, S. ZENG, *Inequalities of trapezoidal type involving generalized fractional integrals*, Accepted.
- [2] P. CIATTI, M. G. COWLING, F. RICCI, *Hardy and uncertainty inequalities on stratified Lie groups*, Adv. Math. 277 (2015), pp. 365–387.
- [3] S. S. DRAGOMIR, C. E. M. PEARCE, *Selected topics on Hermite–Hadamard inequalities and applications*, RGMIA Monographs, Victoria University, 2000.
- [4] S. S. DRAGOMIR, *Hermite–Hadamard inequality for fractional integrals via η -convex functions*, Acta Math. Univ. Comenianae LXXXVI(1) (2017), pp. 153–164.
- [5] S. S. DRAGOMIR, M. I. BHATTI, M. IQBAL, M. MUDDASSAR, *Some new Hermite–Hadamard’s type fractional integral inequalities*, J. Comput. Anal. Appl. 18 (4) (2015), pp. 655–661.
- [6] A. FERNANDEZ, P. O. MOHAMMED, *Hermite–Hadamard inequalities in fractional calculus defined using Mittag–Leffler kernels*, Math. Meth. Appl. Sci. 2020, 1–18, <https://doi.org/10.1002/mma.6188>.
- [7] B. GAVREA, I. GAVREA, *On some Ostrowski type inequalities*, Gen. Math. 18 (1) (2010), pp. 33–44.
- [8] H. GUNAWAN, ERIDANI, *Fractional integrals and generalized Olsen inequalities*, Kyungpook Math. J. 49 (2009), pp. 31–39.
- [9] M. A. HANSON, *On sufficiency of the Kuhn–Tucker conditions*, J. Math. Anal. Appl. 80 (1981), pp. 545–550.
- [10] A. A. KILBAS, H. M. SRIVASTAVA, J. J. TRUJILLO, *Theory and Applications of Fractional Differential Equations*, Elsevier B. V., Amsterdam, Netherlands, 2006.
- [11] T. LIAN, W. TANG, R. ZHOU, *Fractional Hermite–Hadamard inequalities for (s,m) -convex or s -concave functions*, J. Inequal. Appl. 2018 (2018), 240.
- [12] P. O. MOHAMMED, *New integral inequalities for preinvex functions via generalized beta function*, J. Interdiscip. Math. 22 (4) (2019), pp. 539–549.
- [13] P. O. MOHAMMED, *On New Trapezoid Type Inequalities for h -convex Functions via Generalized Fractional Integral*, Turkish J. Anal. Number Theory 6 (4) (2018), pp. 125–128.
- [14] P. O. MOHAMMED, *Some new Hermite–Hadamard type inequalities for MT -convex functions on differentiable coordinates*, J. King Saud Univ. Sci. 30 (2018), pp. 258–262.
- [15] P. O. MOHAMMED, *Inequalities of Type Hermite–Hadamard for Fractional Integrals via Differentiable Convex Functions*, Turkish J. Anal. Number Theory 4 (5) (2016), pp. 135–139.
- [16] P. O. MOHAMMED, *Inequalities of (k,s) , (k,h) -Type For Riemann–Liouville Fractional Integrals*, Appl. Math. E-Notes 17 (2017), pp. 199–206.
- [17] P. O. MOHAMMED, *Hermite–Hadamard inequalities for Riemann–Liouville fractional integrals of a convex function with respect to a monotone function*, Math. Meth. Appl. Sci. (2019), pp. 1–11, <https://doi.org/10.1002/mma.5784>.
- [18] P. O. MOHAMMED, T. ABDELJAWAD, *Modification of certain fractional integral inequalities for convex functions*, Adv. Differ. Equ. 2020 (2020), 69.
- [19] P. O. MOHAMMED, I. BREVIK, *A New Version of the Hermite–Hadamard Inequality for Riemann–Liouville Fractional Integrals*, accepted.
- [20] P. O. MOHAMMED, F. K. HAMASALH, *New Conformable Fractional Integral Inequalities of Hermite–Hadamard Type for Convex Functions*, Symmetry 11 (2) (2019), 263, <https://doi.org/10.3390/sym11020263>.
- [21] P. O. MOHAMMED, M. Z. SARIKAYA, *Hermite–Hadamard type inequalities for F -convex function involving fractional integrals*, J. Inequal. Appl. 2018 (2018), 359.
- [22] P. O. MOHAMMED, M. Z. SARIKAYA, *On generalized fractional integral inequalities for twice differentiable convex functions*, J. Comput. Appl. Math. 372 (2020), 112740.

- [23] P. O. MOHAMMED, M. Z. SARIKAYA, D. BALEANU, *On the Generalized Hermite-Hadamard Inequalities via the Tempered Fractional Integrals*, Symmetry 12 (4) (2020), 595, <https://doi:10.3390/sym12040595>.
- [24] F. QI, P. O. MOHAMMED, J. C. YAO, Y. H. YAO, *Generalized fractional integral inequalities of Hermite–Hadamard type for (α, m) -convex functions*, J. Inequal. Appl. 2019 (2019), 135.
- [25] M. E. OZDEMIR, M. AVCI, H. KAVURMACI, *Hermite–Hadamard-type inequalities via (α, m) -convexity*, Comput. Math. Appl. 61 (2011), pp. 2614–2620.
- [26] J. PARK, *Some Hermite–Hadamard type inequalities for MT-convex functions via classical and Riemann–Liouville fractional integrals*, Appl. Math. Sci. 9 (101) (2015), pp. 5011–5026.
- [27] M. Z. SARIKAYA, E. SET, H. YALDIZ, N. BAŞAK, *Hermite–Hadamard’s inequalities for fractional integrals and related fractional inequalities*, Math. Comput. Model. 57 (2013), pp. 2403–2407.
- [28] Y. SAWANO, H. WADADE, *On the Gagliardo–Nirenberg type inequality in the critical Sobolev–Morrey space*, J. Fourier Anal. Appl. 19 (1) (2013), pp. 20–47.
- [29] E. SET, M. E. OZDEMIR, M. Z. SARIKAYA, F. KARAKOC, *Hermite–Hadamard type inequalities for (α, m) -convex functions via fractional integrals*, Moroccan J. Pure and Appl. Anal. 3 (1) (2017), pp. 15–21.
- [30] J. V. C. SOUSA, E. C. OLIVEIRA, *On the Ψ -Hilfer fractional derivative*, Commun. Nonlinear Sci. Numer. Simul. 60, pp. 72–91.
- [31] J. WANGA, X. LIA, M. FECKANB, Y. ZHOU, *Hermite–Hadamard-type inequalities for Riemann–Liouville fractional integrals via two kinds of convexity*, Applicable Analysis 92(11) (2013), pp. 2241–2253.
- [32] A. WEIR, B. MOND, *Preinvex functions in multiple objective optimization*, J. Math. Anal. Appl. 136 (1988), pp. 29–38.

(Received May 1, 2019)

*Pshtiwan Othman Mohammed
Department of Mathematics
College of Education, University of Sulaimani
Sulaimani, Kurdistan Region, Iraq
e-mail: pshtiwansangawi@gmail.com*