

A GENERAL NONLINEAR VERSION OF ROTH'S THEOREM ON THE REAL LINE

XIANG LI, DUNYAN YAN, HAIXIA YU AND XINGSONG ZHANG*

Abstract. Let $N > 1$ be a real number and $\varepsilon > 0$ be given. In this paper, we will prove that, for a measurable subset S of $[0, N]$ with positive density ε , there must be patterns of the form $(x, x+t, x+\gamma(t))$ such that

$$x, x+t, x+\gamma(t) \in S,$$

where γ is convex and has some curvature constraints, $t > \delta(\varepsilon, \gamma)\gamma^{-1}(N)$ and $\delta(\varepsilon, \gamma)$ is a positive constant depending only on ε and γ , γ^{-1} is the inverse function of γ . Our result extends Bourgain's result [2] to the general curve γ . We use Bourgain's energy pigeonholing argument and Li's σ -uniformity argument.

Mathematics subject classification (2020): Primary 42B20; Secondary 05D10.

Keywords and phrases: Roth's theorem, convex curve, stationary phase method, σ -uniformity.

REFERENCES

- [1] V. BERGELSON AND A. LEIBMAN, *Polynomial extensions of van der Waerden's and Szemerédi's theorems*, J. Amer. Math. Soc. 9 (1996), no. 3, 725–753.
- [2] J. BOURGAIN, *A nonlinear version of Roth's theorem for sets of positive density in the real line*, J. Analyse Math. 50 (1988), 169–181.
- [3] J. BOURGAIN, *A Szemerédi type theorem for sets of positive density in R^k* , Israel J. Math. 54 (1986), no. 3, 307–316.
- [4] J. BOURGAIN, M.-C. CHANG, *Nonlinear Roth type theorems in finite fields*, Israel J. Math. 221 (2017), no. 2, 853–867.
- [5] A.-P. CALDERÓN, *Cauchy integrals on Lipschitz curves and related operators*, Proc. Nat. Acad. Sci. U.S.A. 74 (1977), no. 4, 1324–1327.
- [6] A. CARBERY, M. CHRIST, J. VANCE, S. WAINGER AND D. WATSON, *Operators associated to flat plane curves: L^p estimates via dilation methods*, Duke Math. J. 59 (1989), no. 3, 675–700.
- [7] A. CARBERY, J. VANCE, S. WAINGER AND D. WATSON, *The Hilbert transform and maximal functions along flat curves, dilations, and differential equations*, Amer. J. Math. 116 (1994), no. 5, 1203–1239.
- [8] H. CARLSSON, M. CHRIST, A. CORDOBA, J. DUANDIKOETXEA, J. L. RUBIO DE FRANCIA, J. VANCE, S. WAINGER AND D. WEINBERG, *L^p estimates for maximal functions and Hilbert transforms along flat convex curves in \mathbb{R}^2* , Bull. Amer. Math. Soc. (N.S.) 14 (1986), no. 2, 263–267.
- [9] M. CHRIST, X. LI, T. TAO AND C. THIELE, *On multilinear oscillatory integrals, nonsingular and singular*, Duke Math. J. 130 (2005), no. 2, 321–351.
- [10] M. CHRIST, A. NAGEL, E. M. STEIN AND S. WAINGER, *Singular and maximal Radon transforms: analysis and geometry*, Ann. of Math. (2) 150 (1999), no. 2, 489–577.
- [11] B. COOK, Á. MAGYAR AND M. PRAMANIK, *A Roth-type theorem for dense subsets of \mathbb{R}^d* , Bull. Lond. Math. Soc. 49 (2017), no. 4, 676–689.
- [12] D. DONG, X. LI AND W. SAWIN, *Improved estimates for polynomial Roth type theorems in finite fields*, arXiv:1709.00080, 2017.
- [13] P. DURCIK, S. GUO AND J. ROOS, *A polynomial Roth theorem on the real line*, Trans. Amer. Math. Soc. 371 (2019), no. 10, 6973–6993.

- [14] P. ERDŐS AND P. TURAŃ, *On Some Sequences of Integers*, J. London Math. Soc. 11 (1936), no. 4, 261–264.
- [15] E. B. FABES AND N. M. RIVIÈRE, *Singular integrals with mixed homogeneity*, Studia Math. 27 (1966), 19–38.
- [16] R. FRASER, S. GUO AND M. PRAMANIK, *Polynomial Roth theorems on sets of fractional dimensions*, arXiv:1904.11123, 2019.
- [17] H. FURSTENBERG, *Ergodic behavior of diagonal measures and a theorem of Szemerédi on arithmetic progressions*, J. Analyse Math. 31 (1977), 204–256.
- [18] H. FURSTENBERG AND Y. KATZNELSON, *An ergodic Szemerédi theorem for commuting transformations*, J. Analyse Math. 34 (1978), 275–291 (1979).
- [19] W. T. GOWERS, *A new proof of Szemerédi theorem*, Geom. Funct. Anal. 11 (2001), no. 3, 465–588.
- [20] W. T. GOWERS, *A new proof of Szemerédi theorem for arithmetic progressions of length four*, Geom. Funct. Anal. 8 (1998), no. 3, 529–551.
- [21] B. GREEN, *On arithmetic structures in dense sets of integers*, Duke Math. J. 114 (2002), no. 2, 215–238.
- [22] B. GREEN AND T. TAO, *The primes contain arbitrarily long arithmetic progressions*, Ann. of Math. (2) 167 (2008), no. 2, 481–547.
- [23] J. GUO AND L. XIAO, *Bilinear Hilbert transforms associated with plane curves*, J. Geom. Anal. 26 (2016), no. 2, 967–995.
- [24] L. HÖRMANDER, *Oscillatory integrals and multipliers on L^p* , Ark. Mat. 11 (1973), 1–11.
- [25] B. F. JONES, *A class of singular integrals*, Amer. J. Math. 86 (1964), 441–462.
- [26] M. T. LACEY AND C. THIELE, *L^p estimates on the bilinear Hilbert transform for $2 < p < \infty$* , Ann. of Math. (2) 146 (1997), no. 3, 693–724.
- [27] M. T. LACEY AND C. THIELE, *On Calderón’s conjecture*, Ann. of Math. (2) 149 (1999), no. 2, 475–496.
- [28] J. LI AND H. YU, *L^p boundedness of Hilbert transforms associated with variable plane curves*, arXiv: 1806.08589, 2018.
- [29] X. LI, *Bilinear Hilbert transforms along curves I: The monomial case*, Anal. PDE 6 (2013), no. 1, 197–220.
- [30] X. LI AND L. XIAO, *Uniform estimates for bilinear Hilbert transforms and bilinear maximal functions associated to polynomials*, Amer. J. Math. 138 (2016), no. 4, 907–962.
- [31] V. LIE, *On the boundedness of the bilinear Hilbert transform along “non-flat” smooth curves*, Amer. J. Math. 137 (2015), no. 2, 313–363.
- [32] V. LIE, *On the boundedness of the bilinear Hilbert transform along “non-flat” smooth curves. The Banach triangle case (L^r , $1 \leq r < \infty$)*, Rev. Mat. Iberoam. 34 (2018), no. 1, 331–353.
- [33] A. NAGEL, J. VANCE, S. WAINGER AND D. WEINBERG, *Hilbert transforms for convex curves*, Duke Math. J. 50 (1983), no. 3, 735–744.
- [34] K. F. ROTH, *On certain sets of integers*, J. London Math. Soc. 28 (1953), 104–109.
- [35] E. M. STEIN, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Mathematical Series 43, Monographs in Harmonic Analysis III, Princeton University Press, Princeton, NJ, 1993.
- [36] E. SZEMERÉDI, *On sets of integers containing no four elements in arithmetic progression*, Acta Math. Acad. Sci. Hungar. 20 (1969), 89–104.
- [37] E. SZEMERÉDI, *On sets of integers containing no k elements in arithmetic progression*, Acta Arith. 27 (1975), 199–245.
- [38] T. TAO AND T. ZIEGLER, *A multi-dimensional Szemerédi theorem for the primes via a correspondence principle*, Israel J. Math. 207 (2015), no. 1, 203–228.
- [39] T. TAO AND T. ZIEGLER, *The primes contain arbitrarily long polynomial progressions*, Acta Math. 201 (2008), no. 2, 213–305.