

TWO NEW LOWER BOUNDS FOR THE SMALLEST SINGULAR VALUE

XU SHUN

(Communicated by M. Krnić)

Abstract. In this paper, we obtain two new lower bounds for the smallest singular value of nonsingular matrices which is better than the bound presented by Zou [1], Lin and Xie [2] under certain circumstances.

1. Introduction

Let M_n ($n \geq 2$) be the space of $n \times n$ complex matrices. Let σ_i ($i = 1, \dots, n$) be the singular values of $A \in M_n$ which is nonsingular and suppose that $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_{n-1} \geq \sigma_n > 0$. For $A = [a_{ij}] \in M_n$, the Frobenius norm of A is defined by

$$\|A\|_F = \left(\sum_{i,j=1}^n |a_{ij}|^2 \right)^{1/2} = \text{tr}(A^H A)^{\frac{1}{2}}$$

where A^H is the conjugate transpose of A . The relationship between the Frobenius norm and singular values is

$$\|A\|_F^2 = \sigma_1^2 + \sigma_2^2 + \dots + \sigma_n^2.$$

It is well known that lower bounds for the smallest singular value σ_n of a nonsingular matrix $A \in M_n$ have many potential theoretical and practical applications [3, 4]. Yu and Gu [5] obtained a lower bound for σ_n as follows:

$$\sigma_n \geq |\det A| \cdot \left(\frac{n-1}{\|A\|_F^2} \right)^{(n-1)/2} = l > 0.$$

The above inequality is also shown in [6]. In [1], Zou improved the above inequality by showing that

$$\sigma_n \geq |\det A| \left(\frac{n-1}{\|A\|_F^2 - l^2} \right)^{(n-1)/2} = l_0.$$

In [2], Lin, Minghua and Xie, Mengyan improve a lower bound for smallest singular value of matrices by showing that a is the smallest positive solution to the equation

$$x^2 (\|A\|_F^2 - x^2)^{n-1} = |\det A|^2 (n-1)^{n-1}$$

Mathematics subject classification (2020): 15A18, 15A42, 15A60.

Keywords and phrases: Singular values, Frobenius norm, determinant.

and $\sigma \geq a > l_0$.

In this paper, we obtain two new lower bounds for the smallest singular value of nonsingular matrices. We give some numerical examples which will show that our result is better than l_0 and a under certain circumstances.

2. Main results

LEMMA 1. *Let*

$$l_0 = |\det A| \left(\frac{n-1}{\|A\|_F^2 - l^2} \right)^{(n-1)/2}$$

then $\sigma_n > l_0$.

Proof. In [1], we have

$$\sigma_n \geq |\det A| \left(\frac{n-1}{\|A\|_F^2 - \sigma_n^2} \right)^{(n-1)/2}$$

since $\sigma_n \geq l_0 > l$, thus

$$\begin{aligned} \sigma &\geq |\det A| \left(\frac{n-1}{\|A\|_F^2 - \sigma_n^2} \right)^{(n-1)/2} \\ &\geq |\det A| \left(\frac{n-1}{\|A\|_F^2 - l_0^2} \right)^{(n-1)/2} \\ &> |\det A| \left(\frac{n-1}{\|A\|_F^2 - l^2} \right)^{(n-1)/2} = l_0 \end{aligned}$$

so $\sigma_n > l_0$. \square

THEOREM 1. *Let $A \in M_n$ be nonsingular. Then*

$$\left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - n l_0^2} \right)^{n-1} \right)^{1/2} = l_1$$

then $\sigma_n \geq l_1$, where

$$l = |\det A| \left(\frac{n-1}{\|A\|_F^2} \right)^{\frac{n-1}{2}}, \quad l_0 = |\det A| \left(\frac{n-1}{\|A\|_F^2 - l^2} \right)^{\frac{n-1}{2}}.$$

Proof. Let $0 < \lambda < \sigma_n^2$, then

$$|(\lambda - \sigma_1^2)(\lambda - \sigma_2^2) \cdots (\lambda - \sigma_{n-1}^2)| \leq \left(\frac{\sigma_1^2 + \cdots + \sigma_{n-1}^2 - (n-1)\lambda}{n-1} \right)^{n-1}.$$

Since

$$\begin{aligned} |(\lambda - \sigma_1^2)(\lambda - \sigma_2^2) \cdots (\lambda - \sigma_{n-1}^2)| &= \frac{|(\lambda - \sigma_1^2)(\lambda - \sigma_2^2) \cdots (\lambda - \sigma_n^2)|}{\sigma_n^2 - \lambda} \\ &= \frac{|\det(\lambda I_n - A^H A)|}{\sigma_n^2 - \lambda} \end{aligned}$$

then

$$\begin{aligned} \frac{|\det(\lambda I_n - A^H A)|}{\sigma_n^2 - \lambda} &\leq \left(\frac{\sigma_1^2 + \cdots + \sigma_{n-1}^2 - (n-1)\lambda}{n-1} \right)^{n-1} \\ \sigma_n^2 &\geq \lambda + |\det(\lambda I_n - A^H A)| \left(\frac{n-1}{\sigma_1^2 + \cdots + \sigma_{n-1}^2 - (n-1)\lambda} \right)^{n-1} \\ \sigma_n &\geq \left(\lambda + |\det(\lambda I_n - A^H A)| \left(\frac{n-1}{\sigma_1^2 + \cdots + \sigma_{n-1}^2 - (n-1)\lambda} \right)^{n-1} \right)^{1/2}. \end{aligned}$$

By Lemma 1, $l_0 < \sigma_n$, $l_0^2 < \sigma_n^2$, let $\lambda = l_0^2$, then

$$\sigma_n \geq \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - \sigma_n^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2}. \quad (1)$$

Therefore

$$\sigma_n \geq \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - nl_0^2} \right)^{n-1} \right)^{1/2}. \quad \square$$

THEOREM 2. Let $A \in M_n$ be nonsingular. Let

$$b_{k+1} = \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)l_0^2 - b_k^2} \right)^{n-1} \right)^{1/2}, \quad k = 1, 2, \dots$$

with $l = |\det A| \left(\frac{n-1}{\|A\|_F^2} \right)^{\frac{n-1}{2}}$, $l_0 = |\det A| \left(\frac{n-1}{\|A\|_F^2 - l^2} \right)^{\frac{n-1}{2}}$

$$b_1 = \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2}$$

then $0 < b_k < b_{k+1} \leq \sigma_n$, $k = 1, 2, \dots$, $\lim_{k \rightarrow \infty} b_k$ exists.

Proof. We show by induction on k that

$$\sigma_n \geq b_{k+1} > b_k > 0.$$

By (1), we have

$$\begin{aligned}\sigma_n &\geq \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - \sigma_n^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} \\ &\geq \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} = b_1\end{aligned}$$

so $\sigma_n \geq b_1$, then

$$\begin{aligned}\sigma_n &\geq \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - \sigma_n^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} \\ &\geq \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)l_0^2 - b_1^2} \right)^{n-1} \right)^{1/2} = b_2 \\ &> \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} = b_1 > 0.\end{aligned}$$

When $k = 1$, we have

$$\sigma_n \geq b_2 > b_1 > 0.$$

Assume that our claim is true for $k = m$, that is $\sigma_n \geq b_{m+1} > b_m > 0$. Now we consider the case when $k = m + 1$. By (1), we have

$$\begin{aligned}\sigma_n &\geq \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - \sigma_n^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} \\ &\geq \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - b_{m+1}^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} = b_{m+2} \\ &> \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - b_m^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} = b_{m+1} > 0.\end{aligned}$$

Hence $\sigma_n \geq b_{m+2} > b_{m+1} > 0$. This proves $\sigma_n \geq b_{k+1} > b_k > 0$, $k = 1, 2, \dots$. By the well known monotone convergence theorem, $\lim_{k \rightarrow \infty} b_k$ exists. \square

THEOREM 3. Let $b = \lim_{k \rightarrow \infty} b_k$,

$$f(x) = \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - x^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2}$$

then b is the smallest positive solution to the equation $x = f(x)$, and $\sigma_n \geq b$.

Proof. Let x_0 is the smallest positive solution to the equation $x = f(x)$, we show by induction on k that $x_0 > b_k$, $k = 1, 2, \dots$. When $k = 1$

$$\begin{aligned} x_0 &= \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - x_0^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} \\ &> \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} = b_1. \end{aligned}$$

Assume that our claim is true for $k = m$, that is $\sigma_n > b_m$. Now we consider the case when $k = m + 1$.

$$\begin{aligned} x_0 &= \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - x_0^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} \\ &> \left(l_0^2 + |\det(l_0^2 I_n - A^H A)| \left(\frac{n-1}{\|A\|_F^2 - b_m^2 - (n-1)l_0^2} \right)^{n-1} \right)^{1/2} = b_{m+1}. \end{aligned}$$

Hence $x_0 > b_{m+1}$. This proves $x_0 > b_k$, $k = 1, 2, \dots$. Since b is a positive solution to the equation $x = f(x)$ and $x_0 > b_k$, $k = 1, 2, \dots$, then $b = x_0$. Therefore b is the smallest positive solution to the equation $x = f(x)$ and $\sigma_n \geq b$. \square

Therefore we obtain two new lower bounds l_1 and b for the smallest singular value of nonsingular matrices.

3. Numerical examples

We use Examples 1 and Example 2 to compare the values of l, l_0, l_1 .

EXAMPLE 1. Let

$$A = \begin{bmatrix} 4 & -4 & -3 \\ 3 & 4 & 2 \\ 4 & 1 & 0 \end{bmatrix}.$$

Then $\sigma_{\min} = 0.0231$, and

$$l = 0.0229885$$

$$l_0 = 0.0229886.$$

Our result:

$$l_1 = 0.0230691.$$

EXAMPLE 2. Let

$$A = \begin{bmatrix} 4 & 0 & 0 \\ -1 & 5 & 0 \\ 0 & 5 & 4 \end{bmatrix}.$$

Then

$$l = 1.92771$$

$$l_0 = 2.01806.$$

Our result:

$$l_1 = 2.31515.$$

Next we use the following example to compare the values of a, b, l_1 .

EXAMPLE 3. Let

$$A = \begin{bmatrix} 3 & 2 & 0 \\ 1 & 9 & 5 \\ 0 & 5 & 7 \end{bmatrix}.$$

Then

$$a = 1.0367.$$

Our result:

$$l_1 = 1.3434$$

$$b = 1.3455.$$

Acknowledgements. I would like to express my sincere gratitude to professor Xiao-Wu Chen from University of Science and Technology of China.

REFERENCES

- [1] L. ZOU, *A lower bound for the smallest singular value*, J. Math. Inequal 6 (4) (2012) 625–629.
- [2] M. LIN, M. XIE, *On some lower bounds for smallest singular value of matrices*, Applied Mathematics Letters 121 (2021) 107411.
- [3] R. A. HORN, C. R. JOHNSON, *Matrix analysis*, New York.
- [4] R. A. HORN, R. A. HORN, C. R. JOHNSON, *Topics in matrix analysis*, Cambridge University Press, 1994.
- [5] Y. YISHENG, G. DUNHE, *A note on a lower bound for the smallest singular value*, Linear algebra and its Applications 253 (1–3) (1997) 25–38.
- [6] G. PIAZZA, T. POLITI, *An upper bound for the condition number of a matrix in spectral norm*, Journal of Computational and Applied Mathematics 143 (1) (2002) 141–144.

(Received August 4, 2021)

Xu Shun
 School of Mathematical Sciences
 University of Science and Technology of China
 Hefei, Anhui, 230026, P. R. China
 e-mail: xushun@mail.ustc.edu.cn