

EQUIVALENT CONDITIONS OF OPTIMAL HALF DISCRETE HILBERT TYPE MULTIPLE INTEGRAL INEQUALITIES WITH QUASI HOMOGENEOUS KERNEL AND APPLICATIONS

YONG HONG, BING HE AND MINGJUN FENG*

Abstract. Let $G(u, v)$ be a λ -order homogeneous function. In this paper, by discussing optimal matching parameters of the half discrete Hilbert type multiple integral inequality with quasi-homogeneous kernel $K(n, \|x\|_{\rho, m}) = G(n^{\lambda_1}, \|x\|_{\rho, m}^{\lambda_2})$, several equivalent conditions of the optimal matching parameters are obtained, and a basic theoretical problem of the half discrete Hilbert type inequality is solved. Finally, their applications to operator boundedness and operator norm are discussed.

Mathematics subject classification (2020): 26D15, 47A07.

Keywords and phrases: Quasi-homogeneous kernel, half discrete Hilbert type multiple integral inequality, the best constant factor, optimal matching parameter, equivalent condition, operator norm.

REFERENCES

- [1] L. Z. XU, Y. K. GUO, *Note on Hardy-Riesz's extension of Hilbert's inequality*, Chin. Quart. J. Math., 1991, 6 (1): 75–77.
- [2] O. F. BREVIG, *Sharp norm estimates for composition operators and Hilbert-type inequalities*, B. Lond. Math. Soc., 2017, 49 (6), 965–978.
- [3] T. BATBOLD, M. KRNIĆ AND P. VUKOVIĆ, *A unified approach to fractal Hilbert-type inequalities*, J. Inequal. Appl., 2019, doi:10.1186/s13660-019-2076-9.
- [4] B. YANG, M. HUANG AND Y. ZHONG, *On a parametric Mulholland-type inequality and applications*, Abstr. Appl. Anal., 2019, doi:10.1155/2019/8317029.
- [5] L. HE, H. LIU AND B. YANG, *On a more accurate reverse Mulholland-type inequality with parameters*, J. Inequal. Appl., 2019, doi:10.1186/s13660-019-2139-y.
- [6] Y. HONG, *On the norm of a singular multiple integral operator with homogeneous kernel and its application*, Chinese Annals of Mathematics, 2011, 32 (5), 599–606.
- [7] Y. HONG, *A Hilbert-Type Integral Inequality with Quasi-homoeneous Kernel and Several Functions*, Acta Mathematics Sinica, Chinese Series, 2014, 57 (5), 833–840.
- [8] Y. HONG, B. HE AND B. YANG, *Necessary and sufficient conditions for the validity of Hilbert type integral inequalities with a class of quasi-homogeneous kernels and its application in operator theory*, J. Math. Inequal., 2018, 12 (3), 777–788.
- [9] Y. HONG, Q. HUANG AND B. YANG, et al., *The necessary and sufficient conditions for the existence of a kind of Hilbert-type multiple integral inequality with the non-homogeneous kernel and its applications*, J. Inequal. Appl., 2017, doi:10.1186/s13660-017-1592-8.
- [10] M. KRNIĆ AND P. VUKOVIĆ, *A class of Hilbert-type inequalities obtained via the improved Young inequality*, Results math., 2017, 71, 185–196.
- [11] R. LUO AND B. YANG, *Parameterized discrete Hilbert-type inequalities with intermediate variables*, J. Inequal. Appl., 2019, doi:10.1186/s13660-019-2095-6.
- [12] A. OSEKOWSKI, *Inequalities for Hilbert operator and its extensions: The probabilistic approach*, Ann. Probab., 2017, 45 (1), 535–563.

- [13] M. TH. RASSIAS, B. YANG AND A. RAIGORODSKII, *Two Kinds of the Reverse Hardy-Type Integral Inequalities with the Equivalent Forms Related to the Extended Riemann Zeta Function*, Appl. Anal. Discrete Math., 2018, 12, 273–296.
- [14] S. H. SAKER, A. A. EL-DEEB AND H. M. REZK, et al., *On Hilbert's inequality on time scales*, Appl. Anal. Discr. Math., 2017, 11 (2), 399–423.
- [15] B. TSERENDORJ, A. VANDANJAV AND L. E. AZAR, *A new discrete Hilbert-type inequality involving partial sums*, J. Inequal. Appl., 2019, doi:10.1186/s13660-019-2087-6.
- [16] G. M. FICHTINGOLOZ, *A course in differential and integral calculus*, People's Education Press, Beijing, China, 1957, 404–423.