

INCREASING PROPERTY AND LOGARITHMIC CONVEXITY OF TWO FUNCTIONS INVOLVING DIRICHLET ETA FUNCTION

DONGKYU LIM* AND FENG QI

Dedicated to retired Professor Shi-Ying Yuan, former President of Henan Polytechnic University, China

(Communicated by L. Mihoković)

Abstract. In the paper, with the help of an integral representation of the Dirichlet eta function and by means of a monotonicity rule for a ratio of two integrals with a parameter, the authors find increasing property and logarithmic convexity of two functions involving the gamma function, the extended binomial coefficient, and the Dirichlet eta function.

1. Introduction

In this paper, we use the notation

$$\mathbb{N} = \{1, 2, \dots\}, \quad \mathbb{N}_0 = \{0, 1, 2, \dots\}, \quad \mathbb{N}_- = \{-1, -2, \dots\}, \quad \mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}.$$

It is well known (see [1, Chapter 6], [21, Chapter 3]) that the classical Euler's gamma function $\Gamma(z)$ can be defined by

$$\Gamma(z) = \lim_{n \rightarrow \infty} \frac{n! n^z}{\prod_{k=0}^n (z+k)}, \quad z \in \mathbb{C} \setminus \{0, -1, -2, \dots\}.$$

According to [6, Fact 13.3], for $z \in \mathbb{C}$ such that $\Re(z) > 1$, the Riemann zeta function $\zeta(z)$ can be defined by

$$\zeta(z) = \sum_{k=1}^{\infty} \frac{1}{k^z} = \frac{1}{1-2^{-z}} \sum_{k=1}^{\infty} \frac{1}{(2k-1)^z} = \frac{1}{1-2^{1-z}} \sum_{k=1}^{\infty} (-1)^{k-1} \frac{1}{k^z}. \quad (1.1)$$

In [21, Section 3.5, pp. 57–58], the analytic continuation of the Riemann zeta function $\zeta(z)$ into the punctured complex plane $\mathbb{C} \setminus \{1\}$ is discussed: the only singularity $z = 1$ is a simple pole with residue 1.

Basing on the last equation in (1.1), ones consider the Dirichlet eta function

$$\eta(z) = \left(1 - \frac{1}{2^{z-1}}\right) \zeta(z), \quad \Re(z) > 0.$$

Mathematics subject classification (2020): Primary 11M06; Secondary 11B73, 11M41, 26A48, 26A51, 33B15.

Keywords and phrases: Dirichlet eta function, integral representation, increasing property, logarithmic convexity, monotonicity rule, extended binomial coefficient, gamma function.

* Corresponding author.

It is also known as alternating zeta function. The Dirichlet eta function $\eta(z)$ has an integral representation

$$\eta(x) = \frac{1}{\Gamma(x)} \int_0^\infty \frac{t^{x-1}}{e^t + 1} dt, \quad x > 0 \tag{1.2}$$

in [8, p. 1046, 9.513.1] and [9, p. 604, 25.5.3]. In 1998, Wang [22] obtained that the Dirichlet eta function $\eta(x)$ is logarithmically concave on $(0, \infty)$. In 2009, Cerone and Dragomir [7] established many inequalities and properties for the Riemann eta function $\zeta(x)$ and the Dirichlet eta function $\eta(x)$. In 2015, Adell and Lekuona [2] and Alzer and Kwong [3] strengthened the logarithmic concavity in [22] to a concavity of the Dirichlet eta function $\eta(x)$ on $(0, \infty)$. In 2018, Qi [16, 18] used the logarithmic concavity of the Dirichlet eta function $\eta(x)$ in [22] to establish a double inequality of the ratio $\frac{|B_{2(n+1)}|}{|B_{2n}|}$ for $n \in \mathbb{N}$, where the Bernoulli numbers B_{2n} for $n \geq 0$ are generated by

$$\frac{z}{e^z - 1} = \sum_{n=0}^\infty B_n \frac{z^n}{n!} = 1 - \frac{z}{2} + \sum_{n=1}^\infty B_{2n} \frac{z^{2n}}{(2n)!}, \quad |z| < 2\pi.$$

Very recently, the double inequality established in [16, 18] was extended, sharpened, discussed in the papers [20, 25, 26].

In this paper, we consider

1. the functions

$$x \mapsto \binom{x + \alpha + \ell}{\alpha} \frac{\eta(x + \alpha)}{\eta(x)}, \quad \ell = 0, 1 \tag{1.3}$$

and their monotonicity on $(0, \infty)$, where $\alpha > 0$ is a constant,

$$\binom{z}{w} = \begin{cases} \frac{\Gamma(z+1)}{\Gamma(w+1)\Gamma(z-w+1)}, & z \notin \mathbb{N}_-, \quad w, z-w \notin \mathbb{N}_- \\ 0, & z \notin \mathbb{N}_-, \quad w \in \mathbb{N}_- \text{ or } z-w \in \mathbb{N}_- \\ \frac{\langle z \rangle_w}{w!}, & z \in \mathbb{N}_-, \quad w \in \mathbb{N}_0 \\ \frac{\langle z \rangle_{z-w}}{(z-w)!}, & z, w \in \mathbb{N}_-, \quad z-w \in \mathbb{N}_0 \\ 0, & z, w \in \mathbb{N}_-, \quad z-w \in \mathbb{N}_- \\ \infty, & z \in \mathbb{N}_-, \quad w \notin \mathbb{Z} \end{cases} \tag{1.4}$$

denotes the extended binomial coefficient, and

$$\langle \beta \rangle_n = \prod_{k=0}^{n-1} (\beta - k) = \begin{cases} \beta(\beta - 1) \cdots (\beta - n + 1), & n \in \mathbb{N} \\ 1, & n = 0 \end{cases}$$

for $\beta \in \mathbb{C}$ is called the falling factorial;

2. the functions $\Gamma(x + \ell)\eta(x)$ on $(0, \infty)$ for $\ell = 1, 2$ and their logarithmic convexity.

2. A monotonicity rule

For proving our main results in this paper, we need the following lemma, a monotonicity rule for the ratio of two integrals with a parameter, which can be found in [17, Lemma 2.7 and Remark 6.3] and [19, Remark 7.2].

LEMMA 2.1. *Let $U(t), V(t) > 0$, and $W(t, x) > 0$ be integrable in $t \in (a, b)$,*

1. *if the ratios $\frac{t \partial W(t, x) / \partial x}{W(t, x)}$ and $\frac{U(t)}{V(t)}$ are both increasing or both decreasing in $t \in (a, b)$, then the ratio*

$$R(x) = \frac{\int_a^b W(t, x) U(t) dt}{\int_a^b W(t, x) V(t) dt}$$

is increasing in x ;

2. *if one of the ratios $\frac{\partial W(t, x) / \partial x}{W(t, x)}$ and $\frac{U(t)}{V(t)}$ is increasing and another is decreasing in $t \in (a, b)$, then the ratio $R(x)$ is decreasing in x .*

REMARK 2.1. Lemma 4 in [24] states that, if the ratio $\frac{A(x)}{B(x)}$ is increasing on $(0, \infty)$, then the ratio $\frac{\int_0^\infty A(x)e^{-xt} dx}{\int_0^\infty B(x)e^{-xt} dx}$ is decreasing in $t \in (0, \infty)$. This monotonicity rule for the ratio of two Laplace transforms can be deduced from Lemma 2.1 by setting $(a, b) = (0, \infty)$, $U(x) = A(x)$, $V(x) = B(x)$, and $W(x, t) = e^{-xt}$. This means that Lemma 2.1 is a generalization of [24, Lemma 4].

There have been a number of literature, such as [4, pp. 10–11, Theorem 1.25] and [5, 10, 11, 12, 13, 14, 15, 23], dedicated to investigation and application of various monotonicity rules for ratios of two functions, of two integrals, of two Laplace transforms, and the like.

3. Increasing property and logarithmic convexity

We are now in a position to state and prove our main results in this paper.

THEOREM 3.1. *Let $\alpha > 0$ be a constant and let $\ell \geq 0$ be an integer. Then the two functions defined in (1.3) are increasing from $(0, \infty)$ onto $(0, \infty)$. Consequently, for fixed $\ell = 1, 2$, the functions $\Gamma(x + \ell)\eta(x)$ are logarithmically convex in $x \in (0, \infty)$.*

Proof. With the aid of the recurrence relation $\Gamma(z + 1) = z\Gamma(z)$ and the integral representation (1.2), integrating by parts yields

$$\begin{aligned} \frac{\Gamma(x + \alpha + 1)}{\Gamma(x + 1)} \frac{\eta(x + \alpha)}{\eta(x)} &= \frac{\Gamma(x + \alpha + 1)}{\Gamma(x + 1)} \frac{1}{\Gamma(x + \alpha)} \frac{\int_0^\infty \frac{t^{x + \alpha - 1}}{e^t + 1} dt}{\frac{1}{\Gamma(x)} \int_0^\infty \frac{t^{x - 1}}{e^t + 1} dt} \\ &= \frac{(x + \alpha) \int_0^\infty \frac{t^{x + \alpha - 1}}{e^t + 1} dt}{x \int_0^\infty \frac{t^{x - 1}}{e^t + 1} dt} \end{aligned}$$

$$\begin{aligned}
 &= \frac{\int_0^\infty \frac{1}{e^t+1} \frac{dt^{x+\alpha}}{dt} dt}{\int_0^\infty \frac{1}{e^t+1} \frac{dt^x}{dt} dt} \\
 &= \frac{\int_0^\infty \left[\frac{e^t}{(e^t+1)^2} t^\alpha \right] t^x dt}{\int_0^\infty \frac{e^t}{(e^t+1)^2} t^x dt}.
 \end{aligned}$$

Applying Lemma 2.1 to

$$U(t) = \frac{e^t}{(e^t+1)^2} t^\alpha, \quad V(t) = \frac{e^t}{(e^t+1)^2} > 0, \quad W(t,x) = t^x > 0,$$

and $(a,b) = (0, \infty)$, since $\frac{U(t)}{V(t)} = t^\alpha$ and $\frac{\partial W(t,x)/\partial x}{W(t,x)} = \ln t$ are both increasing on $(0, \infty)$, we conclude by (1.4) that the function

$$\begin{aligned}
 \frac{\int_0^\infty \left[\frac{e^t}{(e^t+1)^2} t^\alpha \right] t^x dt}{\int_0^\infty \frac{e^t}{(e^t+1)^2} t^x dt} &= \frac{\Gamma(x+\alpha+1)}{\Gamma(x+1)} \frac{\eta(x+\alpha)}{\eta(x)} \\
 &= \Gamma(\alpha+1) \binom{x+\alpha}{\alpha} \frac{\eta(x+\alpha)}{\eta(x)}
 \end{aligned}$$

is increasing in $x \in (0, \infty)$.

Similarly, integrating by parts, we obtain

$$\begin{aligned}
 \frac{\Gamma(x+\alpha+2)}{\Gamma(x+2)} \frac{\eta(x+\alpha)}{\eta(x)} &= \frac{\Gamma(x+\alpha+2)}{\Gamma(x+\alpha+1)} \frac{\Gamma(x+\alpha+1)}{\Gamma(x+1)} \frac{\eta(x+\alpha)}{\eta(x)} \\
 &= \frac{\Gamma(x+\alpha+2)}{\Gamma(x+1)} \frac{\int_0^\infty \frac{e^t}{(e^t+1)^2} t^{x+\alpha} dt}{\int_0^\infty \frac{e^t}{(e^t+1)^2} t^x dt} \\
 &= \frac{\Gamma(x+\alpha+2)}{\Gamma(x+1)} \frac{\int_0^\infty \left(\frac{1}{e^t+1}\right)' t^{x+\alpha} dt}{\int_0^\infty \left(\frac{1}{e^t+1}\right)' t^x dt} \\
 &= \frac{\Gamma(x+\alpha+2)}{\Gamma(x+2)} (x+\alpha+1) \frac{\int_0^\infty \left(\frac{1}{e^t+1}\right)' t^{x+\alpha} dt}{(x+1) \int_0^\infty \left(\frac{1}{e^t+1}\right)' t^x dt} \\
 &= \frac{\int_0^\infty \left(\frac{1}{e^t+1}\right)' \frac{dt^{x+\alpha+1}}{dt} dt}{\int_0^\infty \left(\frac{1}{e^t+1}\right)' \frac{dt^{x+1}}{dt} dt} \\
 &= \frac{\int_0^\infty \left(\frac{1}{e^t+1}\right)'' t^{x+\alpha+1} dt}{\int_0^\infty \left(\frac{1}{e^t+1}\right)'' t^{x+1} dt} \\
 &= \frac{\int_0^\infty \left[\frac{e^t(e^t-1)}{(e^t+1)^3} t^{\alpha+1} \right] t^x dt}{\int_0^\infty \left[\frac{e^t(e^t-1)}{(e^t+1)^3} t \right] t^x dt}.
 \end{aligned}$$

Applying Lemma 2.1 to

$$U(t) = \frac{e^t(e^t-1)}{(e^t+1)^3} t^{\alpha+1}, \quad V(t) = \frac{e^t(e^t-1)}{(e^t+1)^3} t > 0, \quad W(t,x) = t^x > 0,$$

and $(a, b) = (0, \infty)$, since $\frac{U(t)}{V(t)} = t^\alpha$ and $\frac{\partial W(t,x)/\partial x}{W(t,x)} = \ln t$ are both increasing on $(0, \infty)$, we conclude by (1.4) that the function

$$\begin{aligned} \frac{\int_0^\infty \left[\frac{e^t(e^t-1)}{(e^t+1)^3} t^{\alpha+1} \right] t^x dt}{\int_0^\infty \left[\frac{e^t(e^t-1)}{(e^t+1)^3} t \right] t^x dt} &= \frac{\Gamma(x+\alpha+2)}{\Gamma(x+2)} \frac{\eta(x+\alpha)}{\eta(x)} \\ &= \Gamma(\alpha+1) \binom{x+\alpha+1}{\alpha} \frac{\eta(x+\alpha)}{\eta(x)} \end{aligned}$$

is increasing in $x \in (0, \infty)$.

Because the function $\frac{\Gamma(x+\alpha+\ell)}{\Gamma(x+\ell)} \frac{\eta(x+\alpha)}{\eta(x)}$ for $\ell = 1, 2$ is increasing in $x \in (0, \infty)$, its first derivative

$$\begin{aligned} &\left[\frac{\Gamma(x+\alpha+\ell)}{\Gamma(x+\ell)} \frac{\eta(x+\alpha)}{\eta(x)} \right]' \\ &= \frac{[\Gamma(x+\alpha+\ell)\eta(x+\alpha)]'[\Gamma(x+\ell)\eta(x)] - [\Gamma(x+\alpha+\ell)\eta(x+\alpha)][\Gamma(x+\ell)\eta(x)]'}{[\Gamma(x+\ell)\eta(x)]^2} \end{aligned}$$

is positive for $x \in (0, \infty)$. Hence, we have

$$\frac{[\Gamma(x+\alpha+\ell)\eta(x+\alpha)]'}{\Gamma(x+\alpha+\ell)\eta(x+\alpha)} > \frac{[\Gamma(x+\ell)\eta(x)]'}{[\Gamma(x+\ell)\eta(x)]},$$

that is, the logarithmic derivative

$$(\ln[\Gamma(x+\ell)\eta(x)])' = \frac{[\Gamma(x+\ell)\eta(x)]'}{[\Gamma(x+\ell)\eta(x)]}$$

is increasing in $x \in (0, \infty)$. Consequently, for $\ell = 1, 2$, the function $\Gamma(x+\ell)\eta(x)$ is logarithmically convex in $(1, \infty)$. The proof of Theorem 3.1 is complete. \square

REMARK 3.1. Because the third derivative

$$\left(\frac{1}{e^t+1} \right)''' = -\frac{e^t[(e^t-1)^2-2e^t]}{(e^t+1)^4}$$

is not monotonic and does not keep the same sign on $(0, \infty)$, we can not find any more result by the method used in this paper.

Acknowledgements. The authors appreciate anonymous referees for their careful corrections to and valuable comments on the original version of this paper.

Funding. The first and corresponding author, Dr. Dongkyu Lim, was partially supported by the National Research Foundation of Korea under Grant NRF-2021R1C1C1 010902, Republic of Korea.

REFERENCES

- [1] M. ABRAMOWITZ AND I. A. STEGUN (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series **55**, 10th printing, Washington, 1972.
- [2] J. A. ADELL AND A. LEKUONA, *Dirichlet's eta and beta functions: concavity and fast computation of their derivatives*, *J. Number Theory* **157** (2015), 215–222; available online at <https://doi.org/10.1016/j.jnt.2015.05.006>.
- [3] H. ALZER AND M. K. KWONG, *On the concavity of Dirichlet's eta function and related functional inequalities*, *J. Number Theory* **151** (2015), 172–196; available online at <https://doi.org/10.1016/j.jnt.2014.12.009>.
- [4] G. D. ANDERSON, M. K. VAMANAMURTHY, AND M. VUORINEN, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.
- [5] G. ANDERSON, M. VAMANAMURTHY, AND M. VUORINEN, *Monotonicity rules in calculus*, *Amer. Math. Monthly* **113** (2006), no. 9, 805–816; available online at <https://doi.org/10.2307/27642062>.
- [6] D. S. BERNSTEIN, *Scalar, Vector, and Matrix Mathematics: Theory, Facts, and Formulas*, Revised and expanded edition, Princeton University Press, Princeton, NJ, 2018.
- [7] P. CERONE AND S. S. DRAGOMIR, *Some convexity properties of Dirichlet series with positive terms*, *Math. Nachr.* **282** (2009), no. 7, 964–975; available online at <https://doi.org/10.1002/mana.200610783>.
- [8] I. S. GRADSHTEYN AND I. M. RYZHIK, *Table of Integrals, Series, and Products*, Translated from the Russian, Translation edited and with a preface by Daniel Zwillinger and Victor Moll, Eighth edition, Revised from the seventh edition, Elsevier/Academic Press, Amsterdam, 2015; available online at <https://doi.org/10.1016/B978-0-12-384933-5.00013-8>.
- [9] F. W. J. OLVER, D. W. LOZIER, R. F. BOISVERT, AND C. W. CLARK (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010; available online at <http://dlmf.nist.gov/>.
- [10] I. PINELIS, *L'Hospital rules for monotonicity and the Wilker-Anglesio inequality*, *Amer. Math. Monthly* **111** (2004), no. 10, 905–909; available online at <http://dx.doi.org/10.2307/4145099>.
- [11] I. PINELIS, *L'Hospital-type rules for monotonicity, and the Lambert and Saccheri quadrilaterals in hyperbolic geometry*, *J. Inequal. Pure Appl. Math.* **6** (2005), no. 4, Article 99, 12 pages; available online at <https://emis.de/journals/JIPAM/article573.html>.
- [12] I. PINELIS, *L'Hospital type rules for monotonicity: applications to probability inequalities for sums of bounded random variables*, *J. Inequal. Pure Appl. Math.* **3** (2002), no. 1, Article 7, 9 pages; available online at <https://emis.de/journals/JIPAM/article158.html>.
- [13] I. PINELIS, *L'Hospital-type rules for monotonicity: discrete case*, *Math. Inequal. Appl.* **11** (2008), no. 4, 647–653; available online at <https://doi.org/10.7153/mia-11-54>.
- [14] I. PINELIS, “Non-strict” *L'Hospital-type rules for monotonicity: intervals of constancy*, *J. Inequal. Pure Appl. Math.* **8** (2007), no. 1, Article 14, 8 pages; available online at <https://emis.de/journals/JIPAM/article827.html>.
- [15] I. PINELIS, *On L'Hospital-type rules for monotonicity*, *J. Inequal. Pure Appl. Math.* **7** (2006), no. 2, Article 40, 19 pages; available online at <https://emis.de/journals/JIPAM/article657.html>.
- [16] F. QI, *A double inequality for the ratio of two non-zero neighbouring Bernoulli numbers*, *J. Comput. Appl. Math.* **351** (2019), 1–5; available online at <https://doi.org/10.1016/j.cam.2018.10.049>.
- [17] FENG QI, *Decreasing properties of two ratios defined by three and four polygamma functions*, *Comptes Rendus Mathématique Académie des Sciences Paris* **360** (2022), 89–101; available online at <https://doi.org/10.5802/crmath.296>.
- [18] F. QI, *Notes on a double inequality for ratios of any two neighbouring non-zero Bernoulli numbers*, *Turkish J. Anal. Number Theory* **6** (2018), no. 5, 129–131; available online at <https://doi.org/10.12691/tjant-6-5-1>.

- [19] F. QI, W.-H. LI, S.-B. YU, X.-Y. DU, AND B.-N. GUO, *A ratio of finitely many gamma functions and its properties with applications*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM. **115** (2021), no. 2, Paper No. 39, 14 pages; available online at <https://doi.org/10.1007/s13398-020-00988-z>.
- [20] Y. SHUANG, B.-N. GUO, AND F. QI, *Logarithmic convexity and increasing property of the Bernoulli numbers and their ratios*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **115** (2021), no. 3, Paper No. 135, 12 pages; available online at <https://doi.org/10.1007/s13398-021-01071-x>.
- [21] N. M. TEMME, *Special Functions: An Introduction to Classical Functions of Mathematical Physics*, A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1996; available online at <https://doi.org/10.1002/9781118032572>.
- [22] K. C. WANG, *The logarithmic concavity of $(1 - 2^{1-r})\zeta(r)$* , J. Changsha Comm. Univ. **14** (1998), no. 2, 1–5, (Chinese).
- [23] S. WU AND L. DEBNATH, *A generalization of L'Hôpital-type rules for monotonicity and its application*, Appl. Math. Lett. **22** (2009), no. 2, 284–290; available online at <https://doi.org/10.1016/j.aml.2008.06.001>.
- [24] Z.-H. YANG AND J.-F. TIAN, *Monotonicity and inequalities for the gamma function*, J. Inequal. Appl. **2017**, paper no. 317, 15 pages; available online at <https://doi.org/10.1186/s13660-017-1591-9>.
- [25] Z.-H. YANG AND J.-F. TIAN, *Sharp bounds for the ratio of two zeta functions*, J. Comput. Appl. Math. **364** (2020), 112359, 14 pages; available online at <https://doi.org/10.1016/j.cam.2019.112359>.
- [26] L. ZHU, *New bounds for the ratio of two adjacent even-indexed Bernoulli numbers*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **114** (2020), no. 2, Paper No. 83, 13 pages; available online at <https://doi.org/10.1007/s13398-020-00814-6>.

(Received June 18, 2021)

Dongkyu Lim
 Department of Mathematics Education
 Andong National University
 Andong 36729, Republic of Korea
 e-mail: dgrim84@gmail.com, dklim@andong.ac.kr
<https://orcid.org/0000-0002-0928-8480>

Feng Qi
 Institute of Mathematics
 Henan Polytechnic University
 Jiaozuo 454010, Henan, China
 and
 School of Mathematical Sciences
 Tiangong University
 Tianjin 300387, China
 e-mail: qifeng618@gmail.com,
 qifeng618@hotmail.com, qifeng618@qq.com
<https://qifeng618.wordpress.com>
<https://orcid.org/0000-0001-6239-2968>