

# A MATRIX INEQUALITY FOR UNITARILY INVARIANT NORMS

XIN JIN, FENG ZHANG AND JINLI XU\*

(Communicated by M. Krnić)

*Abstract.* In this paper, we present an inequality of matrix norms, which is a generalization of the inequality shown by Zou [Linear Algebra Appl. 562, 154–162].

## 1. Introduction

As usual, the set of  $n \times n$  complex matrices is denoted by  $M_n$ . The identity matrix of  $M_n$  is denoted by  $I$ . For  $A \in M_n$ ,  $s_i(A)$  is the  $i$ -th largest singular value of  $A$  and  $s(A) = (s_1(A), \dots, s_n(A))$ .  $\lambda_i(A)$  is the  $i$ -th largest eigenvalue of  $A$  and  $\lambda(A) = (\lambda_1(A), \dots, \lambda_n(A))$ .  $s_i(A) = \lambda_i(|A|)$  for  $|A| = (A^*A)^{\frac{1}{2}}$ , where  $A^*$  is the conjugate transpose of  $A$ .  $A \geq B$  means that  $A - B$  is positive semidefinite. The direct sum of  $A$  and  $B$  is denoted by  $A \oplus B$ . The block matrix is presented by  $[X_{ij}]$  which  $X_{ij}$  is the  $i, j$ -th block.

Now we introduce the definition of majorization. Given a real vector  $x = (x_1, x_2, \dots, x_n) \in R^n$ , we rearrange its components as  $x_{[1]} \geq x_{[2]} \geq \dots \geq x_{[n]}$ .

For  $x = (x_1, x_2, \dots, x_n)$ ,  $y = (y_1, y_2, \dots, y_n) \in R^n$ , if

$$\sum_{i=1}^k x_{[i]} \leq \sum_{i=1}^k y_{[i]}, k = 1, 2, \dots, n,$$

then we say that  $x$  is weakly majorized by  $y$  and denote  $x \prec_w y$ . If  $x \prec_w y$  and  $\sum_{i=1}^n x_i = \sum_{i=1}^n y_i$  hold, then we say that  $x$  is majorized by  $y$  and denote  $x \prec y$ .

Let  $\|\cdot\|$  be any unitary invariant norm. Due to Ky fan's result (see [2]), it's known that  $\|X\| \leq \|Y\|$  if and only if  $s(X) \prec_w s(Y)$  for  $X, Y \in M_n$ . Let  $f$  be convex and increasing function on  $[0, +\infty)$ . If  $x \prec_w y$ , then

$$f(x) \prec_w f(y). \quad (1)$$

For more detail (see [2]).

For  $a \in [0, 1]$ , in [1], Zou proved

$$\left\| |AXB^*|^{2r} \right\| \leq \left\| |aA^*AX + (1-a)XB^*B|^{rp} \right\|^{\frac{1}{p}} \times \left\| |(1-a)A^*AX + aXB^*B|^{rq} \right\|^{\frac{1}{q}} \quad (2)$$

*Mathematics subject classification (2020):* 15A60, 47A63.

*Keywords and phrases:* Arithmetic-geometric mean inequality, unitarily invariant norms, majorization.

\* Corresponding author.

for  $r \geq \max\left\{\frac{1}{p}, \frac{1}{q}\right\}$ . Zou explained that the insertion of  $X$  is no idle generalization, a judicious choice can lead to powerful perturbation theorems. Our aim here is to obtain a stronger version of inequality (2) in the same spirit.

In this paper, we present a generalization of inequality (2).

## 2. Main result

In this section, we first show some Lemmas used in our proof.

LEMMA 1. [3] Let  $A, B \in M_n$  be Hermitian matrices. Then

$$\lambda(A+B) \prec \lambda(A) + \lambda(B).$$

LEMMA 2. [4] Let  $H = [A_{ij}] \in M_{sn}$  ( $s \geq 2$ ) be positive semidefinite matrix with  $A_{ij} = -A_{ij}^*$  ( $i \neq j, i, j = 1, 2, \dots, s$ ). Then

$$\lambda(H) \prec \lambda\left(\sum_{i=1}^s A_{ii} \oplus 0\right). \quad (3)$$

THEOREM 1. Let  $A_i, B_i, X \in M_n$ ,  $p > 1$ ,  $q > 1$  and  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $r \geq \max\left\{\frac{1}{p}, \frac{1}{q}\right\}$ ,  $a \in [0, 1]$ . Then

$$\begin{aligned} \left\| \left| \sum_{i=1}^n A_i X B_i^* \right|^{2r} \right\| &\leq \left\| \left| \sum_{i=1}^n a A_i^* A_i X + \sum_{i=1}^n (1-a) X B_i^* B_i \right|^{rp} \right\|^{\frac{1}{p}} \\ &\times \left\| \left| \sum_{i=1}^n (1-a) A_i^* A_i X + \sum_{i=1}^n a X B_i^* B_i \right|^{rq} \right\|^{\frac{1}{q}} \end{aligned}$$

holds under the conditions that  $A_i^* A_j = -A_j^* A_i$ ,  $B_i^* B_j = -B_j^* B_i$  ( $i \neq j$ ) is skew-Hermitian.

*Proof.* We first discuss the case when  $X$  is positive semidefinite. By inequality (2), we obtain

$$\begin{aligned} &\left\| \left| \sum_{i=1}^n A_i X B_i^* \oplus 0 \right|^{2r} \right\| \\ &= \left\| \left[ \begin{array}{cccc} A_1 X^{\frac{1}{2}} & A_2 X^{\frac{1}{2}} & \cdots & A_n X^{\frac{1}{2}} \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{array} \right] \left[ \begin{array}{cccc} B_1 X^{\frac{1}{2}} & B_2 X^{\frac{1}{2}} & \cdots & B_n X^{\frac{1}{2}} \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{array} \right]^* \right|^{2r} \right\| \\ &\leq \left\| \left( a \left[ X^{\frac{1}{2}} A_i^* A_j X^{\frac{1}{2}} \right] + (1-a) \left[ X^{\frac{1}{2}} B_i^* B_j X^{\frac{1}{2}} \right] \right)^{rp} \right\|^{\frac{1}{p}} \end{aligned}$$

$$\begin{aligned}
& \times \left\| \left( (1-a) \left[ X^{\frac{1}{2}} A_i^* A_j X^{\frac{1}{2}} \right] + a \left[ X^{\frac{1}{2}} B_i^* B_j X^{\frac{1}{2}} \right] \right)^{rq} \right\|^{\frac{1}{q}} \\
& \leqslant \left\| \left( a \sum_{i=1}^n X^{\frac{1}{2}} A_i^* A_i X^{\frac{1}{2}} + (1-a) \sum_{i=1}^n X^{\frac{1}{2}} B_i^* B_i X^{\frac{1}{2}} \right)^{rp} \oplus 0 \right\|^{\frac{1}{p}} \\
& \quad \times \left\| \left( (1-a) \sum_{i=1}^n X^{\frac{1}{2}} A_i^* A_i X^{\frac{1}{2}} + a \sum_{i=1}^n X^{\frac{1}{2}} B_i^* B_i X^{\frac{1}{2}} \right)^{rq} \oplus 0 \right\|^{\frac{1}{q}}
\end{aligned}$$

Hence,

$$\begin{aligned}
\left\| \left| \sum_{i=1}^n A_i X B_i^* \right|^{2r} \right\| & \leqslant \left\| \left( \sum_{i=1}^n a X^{\frac{1}{2}} A_i^* A_i X^{\frac{1}{2}} + \sum_{i=1}^n (1-a) X^{\frac{1}{2}} B_i^* B_i X^{\frac{1}{2}} \right)^{rp} \right\|^{\frac{1}{p}} \\
& \quad \times \left\| \left( \sum_{i=1}^n (1-a) X^{\frac{1}{2}} A_i^* A_i X^{\frac{1}{2}} + \sum_{i=1}^n a X^{\frac{1}{2}} B_i^* B_i X^{\frac{1}{2}} \right)^{rq} \right\|^{\frac{1}{q}}.
\end{aligned}$$

By Proposition 9.1.2 in [2] if a product  $AB$  is Hermitian, then  $\|AB\| \leqslant \|\operatorname{Re} BA\|$ . Using this we obtain

$$\begin{aligned}
& \sum_{j=1}^k s_j \left( \sum_{i=1}^n a X^{\frac{1}{2}} A_i^* A_i X^{\frac{1}{2}} + \sum_{i=1}^n (1-a) X^{\frac{1}{2}} B_i^* B_i X^{\frac{1}{2}} \right) \\
& \leqslant \sum_{j=1}^k s_j \left( \operatorname{Re} \sum_{i=1}^n a A_i^* A_i X + \sum_{i=1}^n (1-a) B_i^* B_i X \right) \\
& \leqslant \sum_{j=1}^k s_j \left( \left| \sum_{i=1}^n a A_i^* A_i X + \sum_{i=1}^n (1-a) X B_i^* B_i \right| \right)
\end{aligned}$$

Since  $r \geqslant \frac{1}{p}$ , by inequality (1), we obtain

$$\left\| \left( \sum_{i=1}^n a X^{\frac{1}{2}} A_i^* A_i X^{\frac{1}{2}} + \sum_{i=1}^n (1-a) X^{\frac{1}{2}} B_i^* B_i X^{\frac{1}{2}} \right)^{rp} \right\| \leqslant \left\| \left| \sum_{i=1}^n a A_i^* A_i X + \sum_{i=1}^n (1-a) X B_i^* B_i \right|^{rp} \right\|.$$

Hence,

$$\begin{aligned}
\left\| \left| \sum_{i=1}^n A_i X B_i^* \right|^{2r} \right\| & \leqslant \left\| \left| \sum_{i=1}^n a A_i^* A_i X + \sum_{i=1}^n (1-a) X B_i^* B_i \right|^{rp} \right\|^{\frac{1}{p}} \\
& \quad \times \left\| \left| \sum_{i=1}^n (1-a) A_i^* A_i X + \sum_{i=1}^n a X B_i^* B_i \right|^{rq} \right\|^{\frac{1}{q}}.
\end{aligned}$$

Next we consider the case when  $X$  is any matrix. By the singular value decomposition we know there exist unitary matrices  $U, V$  such that  $X = UDV^*$ , then

$$\begin{aligned}
& \left\| \left| \sum_{i=1}^n A_i X B_i^* \right|^{2r} \right\| \\
&= \left\| \left| \sum_{i=1}^n A_i U D (B_i V)^* \right|^{2r} \right\| \\
&\leq \left\| \left| \sum_{i=1}^n a U^* A_i^* A_i U D + \sum_{i=1}^n (1-a) D V^* B_i^* B_i V \right|^{rp} \right\|^{\frac{1}{p}} \\
&\quad \times \left\| \left| \sum_{i=1}^n (1-a) U^* A_i^* A_i U D + \sum_{i=1}^n a D V^* B_i^* B_i V \right|^{rq} \right\|^{\frac{1}{q}} \\
&= \left\| \left| \sum_{i=1}^n a A_i^* A_i U D V^* + \sum_{i=1}^n (1-a) U D V^* B_i^* B_i \right|^{rp} \right\|^{\frac{1}{p}} \\
&\quad \times \left\| \left| \sum_{i=1}^n (1-a) A_i^* A_i U D V^* + \sum_{i=1}^n a U D V^* B_i^* B_i \right|^{rq} \right\|^{\frac{1}{q}} \\
&\leq \left\| \left| \sum_{i=1}^n a A_i^* A_i X + \sum_{i=1}^n (1-a) X B_i^* B_i \right|^{rp} \right\|^{\frac{1}{p}} \times \left\| \left| \sum_{i=1}^n (1-a) A_i^* A_i X + \sum_{i=1}^n a X B_i^* B_i \right|^{rq} \right\|^{\frac{1}{q}}. \quad \square
\end{aligned}$$

REMARK 1. Let  $n = 2$ ,  $A_1 = B_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ ,  $A_2 = B_2 = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ . Then

$$\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = A_1^* A_2 \neq A_2^* A_1 = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}, B_1^* B_2 \neq B_2^* B_1$$

and

$$A_1 A_1^* + A_2 A_2^* = \begin{bmatrix} 3 & 0 \\ 0 & 0 \end{bmatrix}, A_1^* A_1 + A_2^* A_2 = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}.$$

A trivial verification shows that  $\lambda_1(A_1 A_1^* + A_2 A_2^*) > \lambda_1(A_1^* A_1 + A_2^* A_2)$ .

REMARK 2. For  $n = 2$ , let  $A_1 = B_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ ,  $A_2 = B_2 = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ ,  $X_1 = I$ ,  $X_2 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ . Then

$$\lambda_1(A_1 X_1 A_1^* + A_2 X_2 A_2^*) > \lambda_1\left(A_1^* A_1 + \frac{A_2^* A_2 X_2 + X_2 A_2^* A_2}{2}\right). \quad (4)$$

Inequality (4) implies that

$$\begin{aligned} \left\| \left| \sum_{i=1}^n A_i X_i B_i^* \right|^{2r} \right\| &\leq \left\| \left| \sum_{i=1}^n a A_i^* A_i X_i + \sum_{i=1}^n (1-a) X_i B_i^* B_i \right|^{rp} \right\|^{\frac{1}{p}} \\ &\times \left\| \left| \sum_{i=1}^n (1-a) A_i^* A_i X_i + \sum_{i=1}^n a X_i B_i^* B_i \right|^{rq} \right\|^{\frac{1}{q}} \end{aligned}$$

isn't always true without  $X_i = X_j$  ( $i \neq j$ ).

**COROLLARY 1.** Let  $A, B, X \in M_n$ ,  $p > 1$ ,  $q > 1$  and  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $r \geq \max \left\{ \frac{1}{p}, \frac{1}{q} \right\}$ ,  $a \in [0, 1]$ . Then

$$\left\| |AXB^*|^{2r} \right\| \leq \left\| |aA^*AX + (1-a)XB^*B|^{rp} \right\|^{\frac{1}{p}} \times \left\| |(1-a)A^*AX + aXB^*B|^{rq} \right\|^{\frac{1}{q}}.$$

*Acknowledgement.* The author would like to thank the referees for their valuable comments which have improve this paper significantly. Jinli Xu is supported by the Fundamental Research Funds for the Central Universities (No. 2572019BC07), the National Natural Science Foundation of China (No. 11701075).

## REFERENCES

- [1] L. ZOU, *Unification of the arithmetic-geometric mean and Hölder inequalities for unitarily invariant norms*, Linear Algebra Appl. **562**, 1 (2019), 154–162.
- [2] R. BHATIA, *Matrix Analysis*, Springer-Verlag, New York, 1997.
- [3] K. FAN, *On a theorem of Weyl concerning eigenvalues of linear transformations I*, Proc. Natl. Acad. Sci. USA. **35**, 11 (1949), 652–655.
- [4] F. ZHANG, AND J. L. XU, *An eigenvalue inequality for positive semidefinite  $k \times k$  block matrices*, J. Math. Ineq. **14**, 4 (2020), 1383–1388.

(Received June 21, 2021)

*Xin Jin*

*Department of Mathematics  
Northeast Forestry University  
Harbin 150040, P. R. China  
e-mail: 2321355910@qq.com*

*Feng Zhang*  
*Department of Higher Mathematics  
Jiangxi College of Science  
Xinyu 338000, P. R. China  
e-mail: fzhang1024@163.com*

*Jinli Xu*  
*Department of Mathematics  
Northeast Forestry University  
Harbin 150040, P. R. China  
e-mail: jcclixv@qq.com*