

APPROXIMATION PROPERTIES OF THE RIEMANN-LIOUVILLE FRACTIONAL INTEGRAL TYPE SZÁSZ-MIRAKYAN-KANTOROVICH OPERATORS

NAZIM MAHHMUDOV AND MUSTAFA KARA

(Communicated by I. Raşa)

Abstract. In the present paper, we introduce the Riemann-Liouville fractional integral type Szász-Mirakyan-Kantorovich operators. We investigate the order of convergence by using Lipschitz-type maximal functions, second order modulus of smoothness and Peetre's K-functional. Weighted approximation properties of these operators in terms of modulus of continuity have been discussed. Then, Vorononskaja-type theorem are obtained. Moreover, bivariate the Riemann-Liouville fractional integral type Szász-Mirakyan-Kantorovich operators are constructed. The last section is devoted to graphical representation and numerical results for these operators.

1. Introduction

Approximation theory plays an important role in mathematical analysis problems and in many fields of science. In particular, linear positive operators have an important role in approximation theory. One of the best known of these operators is the Szász–Mirakjan operator. The classical Szász–Mirakjan operator (see [8] and [9]) is defined for bounded functions $\varphi(\tau)$ on $[0, \infty)$ by the formula

$$S_n(\varphi; \tau) = \sum_{k=0}^n s_{n,k}(\tau) \varphi\left(\frac{k}{n}\right)$$

where $n \in \mathbb{N}$, $\tau \in [0, \infty)$ and $s_{n,k}(\tau) = e^{-n\tau} \frac{(n\tau)^k}{k!}$. In 1954, Butzer [10] generalized into integral modification of the Szász–Mirakjan operators as follows:

$$K_n(\varphi; \tau) = n \sum_{k=0}^{\infty} s_{n,k}(\tau) \int_{\frac{k}{n}}^{\frac{k+1}{n}} \varphi(t) dt. \quad (1)$$

The above operators can be rewritten as follows:

$$K_n(\varphi; \tau) = \sum_{k=0}^{\infty} s_{n,k}(\tau) \int_0^1 \varphi\left(\frac{k+t}{n}\right) dt. \quad (2)$$

Mathematics subject classification (2020): 41A25, 41A36, 47A58.

Keywords and phrases: Szász–Mirakyan–Kantorovich operators, Riemann–Liouville integral.

In the literature, there are a lot of studies that involve Szász operators, Szász-Kantorovich operators and their generalizations. For instance, see [1]–[7] and [20]–[31].

Recently, Fractional calculus and its applications have been paid more and more attention. Fractional calculus deals with the study of fractional degree derivative and integral operators on complex or real fields and their applications. There are several known forms of fractional integrals. The most well-known is:

$$(I_{a^+}^\alpha \varphi)(\tau) = \frac{1}{\Gamma(\alpha)} \int_a^\tau \frac{\varphi(t)}{(\tau-t)^{1-\alpha}} dt \quad (\tau > a; \operatorname{Re}(\alpha) > 0)$$

$$\text{where } \Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt.$$

The aim of the present paper is to construct the Riemann-Liouville fractional integral type Szász-Mirakyan-Kantorovich operators and discuss their approximation properties.

The paper is organized as follows. In Section 2, we introduce the Riemann-Liouville fractional integral type Szász-Mirakyan-Kantorovich operators and give some moments and central moments. In section 3, Korovkin-type approximation theorem is given, and the rate of convergence of these types of operators is obtained for Lipschitz-type maximal functions, second order modulus of smoothness and Peetre's K-functional. In Section 4, weighted approximation properties of the Riemann-Liouville fractional integral type Szász-Mirakyan-Kantorovich operators in terms of modulus of continuity has been discussed. In Section 5, we provide the quantitative Vorononskaja-type asymptotic formula. In section 6, we introduce bivariate the Riemann-Liouville fractional integral type Szász-Mirakyan-Kantorovich operators and their approximation behaviors. Finally, some graphs and error estimation tables are given for the convergence of Szász-Mirakyan-Kantorovich operators of the Riemann-Liouville fractional integral type.

2. The Riemann-Liouville fractional integral type Szász-Mirakyan-Kantorovich operators

Now, for the Szász-Mirakjan operators, the following Lemma is known (see [11] and [12]).

LEMMA 1. ([11] and [12]) *Let $e_i(\tau) = \tau^i$, $i = 1, 2, 3, 4$. Then, for each $\tau \geq 0$, we have*

$$\begin{aligned} S_n(1; \tau) &= 1 \\ S_n(t; \tau) &= \tau \\ S_n(t^2; \tau) &= \tau^2 + \frac{\tau}{n} \\ S_n(t^3; \tau) &= \tau^3 + \frac{3\tau^2}{n} + \frac{\tau}{n^2} \end{aligned}$$

$$S_n(t^4; \tau) = \tau^4 + \frac{6\tau^3}{n} + \frac{7\tau^2}{n^2} + \frac{\tau}{n^3}$$

DEFINITION 2. Let $\alpha > 0$ and $n \in \mathbb{N}$. For $\varphi \in C[0, \infty)$, Riemann-Liouville integral type Szász-Mirakyan-Kantorovich operators can be defined by

$$K_n^{(\alpha)}(\varphi; \tau) = \sum_{k=0}^{\infty} \alpha s_{n,k}(\tau) \int_0^1 \frac{\varphi\left(\frac{k+t}{n}\right)}{(1-t)^{1-\alpha}} dt. \quad (3)$$

where $s_{n,k}(\tau) = e^{-n\tau} \frac{(n\tau)^k}{k!}$.

If $\alpha = 1$, then the operator reduces to classical Szász-Mirakyan-Kantorovich operators.

The moments of the $K_n^{(\alpha)}$ operators plays significant role in our main results. We derive the following formula to obtain them.

LEMMA 3. Let $\alpha > 0, m \in \mathbb{N} \cup \{0\}$ and $n \in \mathbb{N}$. Then for $\tau \geq 0$, we have

$$K_n^{(\alpha)}(t^m; \tau) = \sum_{j=0}^m \binom{m}{j} \frac{\alpha n^j B(m-j+1, \alpha)}{n^m} S_n(t^j; \tau) \quad (4)$$

where

$$S_n(\varphi; \tau) = \sum_{k=0}^n s_{n,k}(\tau) \varphi\left(\frac{k}{n}\right)$$

and

$$B(a, b) = \int_0^1 \tau^{a-1} (1-\tau)^{b-1}, \quad a, b > 0.$$

Proof. From (3), we can write

$$\begin{aligned} K_n^{(\alpha)}(t^m; \tau) &= \sum_{k=0}^{\infty} \alpha s_{n,k}(\tau) \int_0^1 \left(\frac{k+t}{n}\right)^m (1-t)^{\alpha-1} dt \\ &= \sum_{k=0}^{\infty} \alpha s_{n,k}(\tau) \sum_{j=0}^m \binom{m}{j} \frac{k^j}{n^m} \int_0^1 t^{m-j} (1-t)^{\alpha-1} dt \\ &= \sum_{j=0}^m \binom{m}{j} \frac{\alpha n^j B(m-j+1, \alpha)}{n^m} \sum_{k=0}^{\infty} s_{n,k}(\tau) \frac{k^j}{n^j} \\ &= \sum_{j=0}^m \binom{m}{j} \frac{\alpha n^j B(m-j+1, \alpha)}{n^m} S_n(t^j; \tau). \quad \square \end{aligned}$$

The following Lemma for $K_n^{\alpha}(t^j; \tau)$ ($j = 0, 1, 2, 3, 4$) can be written immediately.

LEMMA 4. Let $\alpha > 0$ and $n \in \mathbb{N}$. Then for $\tau \geq 0$, we have

- (i) $K_n^{(\alpha)}(1; \tau) = 1,$
- (ii) $K_n^{(\alpha)}(t; \tau) = \tau + \frac{1}{n(\alpha+1)},$
- (iii) $K_n^{(\alpha)}(t^2; \tau) = \frac{2}{(\alpha+1)(\alpha+2)n^2} + \frac{(\alpha+3)}{(\alpha+1)n}\tau + \tau^2,$
- (iv) $K_n^{(\alpha)}(t^3; \tau) = \frac{6}{(\alpha+1)(\alpha+2)(\alpha+3)n^3} + \frac{6+(\alpha+2)(4\alpha+7)}{(\alpha+1)(\alpha+2)n^2}\tau$
 $+ \frac{3(2+\alpha)}{(\alpha+1)n}\tau^2 + \tau^3$
- (v) $K_n^{(\alpha)}(t^4; \tau) = \frac{24}{(\alpha+1)(\alpha+2)(\alpha+3)(\alpha+4)n^4}$
 $+ \frac{24+(\alpha+3)[12+(\alpha+2)(\alpha+5)]}{(\alpha+1)(\alpha+2)(\alpha+3)(\alpha+4)n^3}\tau$
 $+ \frac{12+(\alpha+2)(7\alpha+19)}{(\alpha+1)(\alpha+2)n^2}\tau^2 + \frac{(6\alpha+10)}{(\alpha+1)n}\tau + \tau^4$

Proof. Since they have the same proof technique, we only give for $K_n^{(\alpha)}(t^2; \tau)$. Using recurrence formula (4) and Lemma 1, we get

$$\begin{aligned} K_{n,k}^{(\alpha)}(t^2; \tau) &= \frac{\alpha B(3, \alpha)}{n^2} S_{n,k}(1; \tau) + \frac{2n\alpha B(2, \alpha)}{n^2} S_{n,k}(t; \tau) + \frac{n^2 \alpha B(1, \alpha)}{n^2} S_{n,k}(t^2; \tau) \\ &= \frac{2}{(\alpha+1)(\alpha+2)n^2} + \frac{2}{(\alpha+1)n}\tau + \left(\tau^2 + \frac{\tau}{n}\right) \\ &= \frac{2}{(\alpha+1)(\alpha+2)n^2} + \frac{(\alpha+3)}{(\alpha+1)n}\tau + \tau^2. \quad \square \end{aligned}$$

We are now ready to present the central moments of the operators $K_n^{(\alpha)}$.

LEMMA 5. Let $\alpha > 0$. For every $\tau \geq 0$, there holds

$$\begin{aligned} K_n^{(\alpha)}(t - \tau; \tau) &= \frac{1}{(\alpha+1)n}, \\ K_n^{(\alpha)}((t - \tau)^2; \tau) &= \frac{2}{(\alpha+1)(\alpha+2)n^2} + \frac{\tau}{n}, \\ K_n^{(\alpha)}((t - \tau)^4; \tau) &= \frac{24}{(\alpha+1)(\alpha+2)(\alpha+3)(\alpha+4)n^4} \\ &\quad + \frac{12+(\alpha+2)(\alpha+5)}{(\alpha+1)(\alpha+2)n^3}\tau + \frac{3}{n^2}\tau^2. \end{aligned}$$

Proof. Since they have the same proof technique, we only give for $K_n^{(\alpha)}((t-\tau)^2; \tau)$. From the linearity of $K_n^{(\alpha)}(t; \tau)$ and lemma 4, we get

$$\begin{aligned} K_n^{(\alpha)}((t-\tau)^2; \tau) &= K_n^{(\alpha)}(t^2; \tau) - 2\tau K_n^{(\alpha)}(t; \tau) + \tau^2 K_n^{(\alpha)}(1; \tau) \\ &= \frac{2}{(\alpha+1)(\alpha+2)n^2} + \frac{(\alpha+3)}{(\alpha+1)n}\tau + \tau^2 \\ &\quad - 2\tau \left(\tau + \frac{1}{n(\alpha+1)} \right) + \tau^2. \quad \square \end{aligned}$$

LEMMA 6. Let $\alpha > 0$ and $n \in \mathbb{N}$. For every $\tau \geq 0$, there holds

$$\lim_{n \rightarrow \infty} nK_{n,k}^{(\alpha)}((t-\tau); \tau) = \frac{1}{(\alpha+1)}, \quad (5)$$

$$\lim_{n \rightarrow \infty} nK_{n,k}^{(\alpha)}((t-\tau)^2; \tau) = \tau, \quad (6)$$

and

$$\lim_{n \rightarrow \infty} nK_{n,k}^{(\alpha)}((t-\tau)^4; \tau) = 0. \quad (7)$$

Proof. From Lemma 5, for every $\tau \geq 0$, we have

$$\lim_{n \rightarrow \infty} nK_{n,k}^{(\alpha)}((t-\tau); \tau) = \frac{1}{(\alpha+1)},$$

$$\lim_{n \rightarrow \infty} nK_{n,k}^{(\alpha)}((t-\tau)^2; \tau) = \lim_{n \rightarrow \infty} n \left(\frac{2}{(\alpha+1)(\alpha+2)n^2} + \frac{\tau}{n} \right) = \tau$$

and

$$\lim_{n \rightarrow \infty} nK_{n,k}^{(\alpha)}((t-\tau)^4; \tau) = 0. \quad \square$$

In [17], Becker proved the following explicit formula for the moments of S_n .

LEMMA 7. [17] For $m \in \mathbb{N}$, there holds

$$S_n(t^m; \tau) = \sum_{j=1}^m a_{m,j} \frac{\tau^j}{n^{m-j}} \quad (8)$$

with positive coefficients $a_{m,j}$. In particular $S_n(t^m; \tau)$ is a polynomial of degree m without a constant term.

For the next main results we need to define the following classes of functions:

1. $B_m[0, \infty) = \{\varphi : [0, \infty) \rightarrow \mathbb{R}; |\varphi(\tau)| \leq M_\varphi (1 + \tau^m)\}$, where M_φ is constant depending on the function $\varphi, m > 0$.
2. $C_m[0, \infty) = B_m[0, \infty) \cap C[0, \infty), m > 0$.

$$3. \quad C_m^*[0, \infty) = \left\{ \varphi : C_m[0, \infty) : \lim_{|\tau| \rightarrow \infty} \frac{|\varphi(\tau)|}{1+\tau^m} < \infty \right\}, \quad m > 0.$$

The norm on the space $C_m^*[0, \infty)$ is showed as $\|\varphi(\tau)\|_m = \sup_{\tau \in [0, \infty)} \frac{|\varphi(\tau)|}{1+\tau^m}$, where $m > 0$.

LEMMA 8. Let $m \in \mathbb{N} \cup \{0\}$ and $\alpha > 0$ be fixed. We have

$$\left\| K_n^{(\alpha)}(1+t^m; \tau) \right\|_m \leq C_{m,j}(\alpha), \quad n \in \mathbb{N}. \quad (9)$$

where $C_m(\alpha)$ is a positive constant. Moreover, we have

$$\left\| K_n^{(\alpha)}(\varphi; \tau) \right\|_m \leq C_{m,j}(\alpha) \|\varphi\|_m, \quad n \in \mathbb{N}, \quad (10)$$

where $\varphi \in C_m^*[0, \infty)$. Thus, for any $m \in \mathbb{N} \cup \{0\}$, $K_n^{(\alpha)} : C_m^*[0, \infty) \rightarrow C_m^*[0, \infty)$ is a linear positive operator.

Proof. For $m = 0$, inequality (9) is obvious.

For $m \geq 1$, combining Lemma (4) and inequality (8), we have

$$\begin{aligned} \frac{1}{\tau^m + 1} K_n^{(\alpha)}(1+t^m; \tau) &= \frac{1}{\tau^m + 1} + \frac{1}{\tau^m + 1} K_n^{(\alpha)}(t^m; \tau) \\ &= \frac{1}{\tau^m + 1} + \frac{1}{\tau^m + 1} \sum_{j=0}^m \binom{m}{j} \frac{\alpha n^j B(m-j+1, \alpha)}{n^m} \sum_{j_0=1}^j a_{j,j_0} \frac{\tau^{j_0}}{n^{j-j_0}} \\ &\leq 1 + k_{m,j}(\alpha) = C_{m,j}(\alpha). \end{aligned}$$

$C_{m,j}(\alpha)$ is a positive constant with depend on m, j and α . Moreover,

$$\left\| K_n^{(\alpha)}(\varphi; \tau) \right\|_m \leq \|\varphi\|_m \left\| K_n^{(\alpha)}(1+t^m; \tau) \right\|_m \quad (11)$$

for every $\varphi \in C_m^*[0, \infty)$. Therefore, from (9), we get

$$\left\| K_n^{(\alpha)}(\varphi; \tau) \right\|_m \leq C_{m,j}(\alpha) \|\varphi\|_m. \quad \square$$

3. Direct results

Let $C_B[0, \infty)$ denote the space of all real-valued continuous and bounded functions φ on $[0, \infty)$. The norm on the space $C_B[0, \infty)$ is showed as

$$\|\varphi\|_{C_B[0, \infty)} = \sup_{\tau \in [0, \infty)} |\varphi(\tau)|.$$

Then, the modulus of continuity of $\varphi \in C_B[0, \infty)$ is given by

$$w(\varphi, \delta) = \sup_{0 < h \leq \delta} \sup_{\tau \in [0, \infty)} |\varphi(\tau+h) - \varphi(\tau)|.$$

Further, Peetre's K -functional is defined by

$$K_2(\varphi; \delta) = \inf_{g \in w^2} \left\{ \|\varphi - g\| + \delta \|g''\| \right\} \quad \delta > 0,$$

where $w^2 := \left\{ g \in C_B[0, \infty) : g', g'' \in C_B[0, \infty) \right\}$. By Theorem 2.4 in [13], there exists an absolute constant $L > 0$ such that

$$K_2(\varphi; \delta) \leq L \omega_2(\varphi; \sqrt{\delta}). \quad (12)$$

where $\delta > 0$ are absolute constant.

Here, $\omega_2(\varphi; \delta)$ is the second order modulus of smoothness of $\varphi \in C_B[0, \infty)$ and defined as

$$\omega_2(\varphi; \delta) = \sup_{0 < h \leq \delta} \sup_{\tau \in [0, \infty)} |\varphi(\tau + 2h) - 2\varphi(\tau + h) + \varphi(\tau)|$$

LEMMA 9. Let $\varphi \in C_B[0, \infty)$ and $\alpha > 0$, Consider the operators

$${}^*K_n^{(\alpha)}(\varphi; \tau) = K_n^{(\alpha)}(\varphi; \tau) + \varphi(\tau) - \varphi\left(\tau + \frac{1}{(\alpha+1)n}\right) \quad (13)$$

Then, for all $g \in w^2$, we have

$$\left| {}^*K_n^{(\alpha)}(g; \tau) - g(\tau) \right| \leq \left(\frac{2}{(\alpha+1)(\alpha+2)n^2} + \frac{\tau}{n} + \left(\frac{1}{(\alpha+1)n} \right)^2 \right) \|g''\|. \quad (14)$$

Proof. From (13) we have

$$\begin{aligned} {}^*K_n^{(\alpha)}((t-\tau); \tau) &= K_n^{(\alpha)}((t-\tau); \tau) - \left(\tau + \frac{1}{(\alpha+1)n} - \tau \right) \\ &= K_n^{(\alpha)}(t; \tau) - \tau K_n^{(\alpha)}(1; \tau) - \left(\tau + \frac{1}{(\alpha+1)n} \right) + \tau = 0. \end{aligned} \quad (15)$$

Let $\tau \in [0, \infty)$ and $g \in w^2$. Using the Taylor's formula,

$$g(t) - g(\tau) = (t - \tau)g'(\tau) + \int_{\tau}^t (t - u)g''(u)du, \quad (16)$$

Applying ${}^*K_n^{(\alpha)}$ and using (15), we can get

$$\begin{aligned} &{}^*K_n^{(\alpha)}(g; \tau) - g(\tau) \\ &= {}^*K_n^{(\alpha)}\left((t-\tau)g'(\tau); \tau\right) + {}^*K_n^{(\alpha)}\left(\int_{\tau}^t (t-u)g''(u)du; \tau\right) \end{aligned}$$

$$\begin{aligned}
&= g'(\tau)^* K_n^{(\alpha)}((t-\tau); \tau) + K_n^{(\alpha)} \left(\int_{\tau}^t (t-u) g''(u) du; \tau \right) \\
&\quad - \int_{\tau}^{\tau + \frac{1}{(\alpha+1)n}} \left(\tau + \frac{1}{(\alpha+1)n} - u \right) g''(u) du \\
&= K_n^{(\alpha)} \left(\int_{\tau}^t (t-u) g''(u) du; \tau \right) - \int_{\tau}^{\tau + \frac{1}{(\alpha+1)n}} \left(\tau + \frac{1}{(\alpha+1)n} - u \right) g''(u) du.
\end{aligned}$$

On the other hand, since

$$\int_{\tau}^t |t-u| |g''(u)| du \leq \|g''\| \int_{\tau}^t |t-u| du \leq (t-\tau)^2 \|g''\|$$

and

$$\left| \int_{\tau}^{\tau + \frac{1}{(\alpha+1)n}} \left(\tau + \frac{1}{(\alpha+1)n} - u \right) g''(u) du \right| \leq \left(\frac{1}{(\alpha+1)n} \right)^2 \|g''\|,$$

we conclude that

$$\begin{aligned}
&\left| {}^*K_n^{(\alpha)}(g; \tau) - g(\tau) \right| \\
&= \left| K_n^{(\alpha)} \left(\int_{\tau}^t (t-u) g''(u) du; \zeta \right) - \int_{\tau}^{\tau + \frac{1}{(\alpha+1)n}} \left(\tau + \frac{1}{(\alpha+1)n} - u \right) g''(u) du \right| \\
&\leq \|g''\| K_n^{(\alpha)} \left((t-\tau)^2; \tau \right) + \left(\frac{1}{(\alpha+1)n} \right)^2 \|g''\|.
\end{aligned}$$

From Lemma 5, we can write

$$\left| {}^*K_n^{(\alpha)}(g; \tau) - g(\tau) \right| \leq \frac{2}{(\alpha+1)(\alpha+2)n^2} + \frac{\tau}{n} + \left(\frac{1}{(\alpha+1)n} \right)^2 \|g''\|. \quad \square$$

THEOREM 10. Let $\varphi \in C_B[0, \infty)$ and $\alpha > 0$. Then, for every $\tau \in [0, \infty)$, there exists a constant $M > 0$ such that

$$\left| K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| \leq M \omega_2 \left(\varphi; \sqrt{\delta_n(\tau)} \right) + \omega \left(\varphi; \beta_n^{(\alpha)}(\tau) \right)$$

where

$$\delta_n^{(\alpha)}(\tau) = \frac{2}{(\alpha+1)(\alpha+2)n^2} + \frac{\tau}{n} + \left(\frac{1}{(\alpha+1)n} \right)^2 \|g''\|$$

and

$$\beta_{n,k}^{(\alpha)}(\tau) = \left| \frac{1}{(\alpha+1)n} \right|$$

Proof. It follows from Lemma 9, that

$$\begin{aligned} \left| K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| &\leq \left| {}^*K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| + \left| \varphi(\tau) - \varphi\left(\tau + \frac{1}{(\alpha+1)n}\right) \right| \\ &\leq \left| {}^*K_n^{(\alpha)}(\varphi - g; \tau) - (\varphi - g)(\tau) \right| \\ &\quad + \left| \varphi(\tau) - \varphi\left(\tau + \frac{1}{(\alpha+1)n}\right) \right| + \left| {}^*K_n^{(\alpha)}(g; \tau) - g(\tau) \right| \\ &\leq \left| {}^*K_n^{(\alpha)}(\varphi - g; \tau) \right| + |(\varphi - g)(\tau)| \\ &\quad + \left| \varphi(\tau) - \varphi\left(\tau + \frac{1}{(\alpha+1)n}\right) \right| + \left| {}^*K_n^{(\alpha)}(g; \tau) - g(\tau) \right|. \end{aligned}$$

Now, considering the boundedness of the ${}^*K_n^{(\alpha)}$ and inequality (14), we get

$$\begin{aligned} \left| K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| &\leq 4\|\varphi - g\| + \left| \varphi(\tau) - \varphi\left(\tau + \frac{1}{(\alpha+1)n}\right) \right| \\ &\quad + \frac{2}{(\alpha+1)(\alpha+2)n^2} + \frac{\tau}{n} + \left(\frac{1}{(\alpha+1)n} \right)^2 \|g''\| \\ &\leq 4\|\varphi - g\| + \omega\left(\varphi; \left| \frac{1}{(\alpha+1)n} \right| \right) + \delta_n(\tau) \|g''\|. \end{aligned}$$

Now, taking infimum on the right hand side over all $g \in w^2$ and using the property of Peetre's K -functional (12), we can get

$$\begin{aligned} \left| K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| &\leq 4K_2\left(\varphi; \delta_n^{(\alpha)}(\tau)\right) + \omega\left(\varphi; \beta_n^{(\alpha)}(\tau)\right) \\ &\leq M\omega_2\left(\varphi; \sqrt{\delta_n^{(\alpha)}(\tau)}\right) + \omega\left(\varphi; \beta_n^{(\alpha)}(\tau)\right). \quad \square \end{aligned}$$

COROLLARY 11. Let $\alpha > 0$. For any $A > 0$ and $\varphi \in C_B[0, \infty)$, then $K_n^{(\alpha)}(\varphi; \tau) \rightrightarrows \varphi$ on $[0, A]$, where the symbol \rightrightarrows denotes the uniform convergence.

THEOREM 12. Let $K_n^{(\alpha)}$ be the operators defined by (3), $\alpha > 0$, $\gamma \in (0, 1]$ and D be any subset of the interval $[0, \infty)$. if $\varphi \in C_B[0, \infty)$ is locally $Lip(\gamma)$ on D , i.e., if φ satisfies the following inequality:

$$|\varphi(t) - \varphi(\tau)| \leq C_{\varphi, \rho} |t - \tau|^\rho, \quad t \in D \text{ and } \tau \in [0, \infty) \quad (17)$$

then for each $\tau \in [0, \infty)$, we have

$$\left| K_{n,k}^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| \leq C_{\varphi, \gamma} \left\{ \left(K_n^{(\alpha)}((t - \tau)^2; \tau) \right)^{\frac{\rho}{2}} + 2d^\rho(\tau, D) \right\}$$

where $C_{\varphi,\rho}$ is constant depending on φ and ρ and $d(\tau, D)$ is the distance between τ and D defined by

$$d(\tau, D) = \inf \{ |t - \tau| : t \in D \}.$$

Proof. Let \overline{D} be the closure of D . Using the properties of infimum, there is at least a point $t_0 \in \overline{D}$ such that $d(\tau, D) = |\tau - t_0|$. By the triangle inequality

$$|\varphi(t) - \varphi(\tau)| \leq |\varphi(t) - \varphi(t_0)| + |\varphi(\tau) - \varphi(t_0)|.$$

Applying $K_n^{(\alpha)}$ to the above inequality and using ((17)), we can get

$$\begin{aligned} |K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau)| &\leq K_n^{(\alpha)}(|\varphi(t) - \varphi(t_0)|; \tau) + K_n^{(\alpha)}(|\varphi(\tau) - \varphi(t_0)|; \tau) \\ &\leq C_{\varphi,\gamma} \left\{ K_n^{(\alpha)}(|t - t_0|^\rho; \tau) + |\tau - t_0|^\rho \right\} \\ &\leq C_{\varphi,\gamma} \left\{ K_n^{(\alpha)}(|t - \tau|^\rho + |\tau - t_0|^\rho; \tau) + |\tau - t_0|^\rho \right\} \\ &= C_{\varphi,\gamma} \left\{ K_n^{(\alpha)}(|t - \tau|^\rho; \tau) + 2|\tau - t_0|^\rho \right\}. \end{aligned}$$

Choosing $a_1 = \frac{2}{\rho}$ and $a_2 = \frac{2}{2-\rho}$ and applying Hölder inequality, we have:

$$\begin{aligned} |K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau)| &\leq C_{\varphi,\gamma} \left\{ \left(K_n^{(\alpha)}(|t - \tau|^{a_1}; \tau) \right)^{\frac{1}{a_1}} \left(K_n^{(\alpha)}(1^{a_2}; \tau) \right)^{\frac{1}{a_2}} + 2d^\rho(\tau, D) \right\} \\ &\leq C_{\varphi,\gamma} \left\{ \left(K_n^{(\alpha)}((t - \tau)^2; \tau) \right)^{\frac{\rho}{2}} + 2d^\rho(\tau, D) \right\}. \quad \square \end{aligned}$$

In [14], Lipschitz type maximal function of the order γ defined as

$$\phi_\rho(\varphi; \tau) = \sup_{\tau, t \in [0, \infty), t \neq \tau} \frac{|\varphi(t) - \varphi(\tau)|}{|t - \tau|^\rho} \quad (18)$$

where $\tau \in [0, \infty)$ and $\gamma \in (0, 1]$. In the next theorem we obtain local direct estimate of the operators $K_n^{(\alpha)}$ by using (18).

THEOREM 13. *Let $\varphi \in C_B[0, \infty)$, $\alpha > 0$ and $\rho \in (0, 1]$. Then, for all $\tau \in [0, \infty)$, we have*

$$|K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau)| \leq \phi_\rho(\varphi; \tau) \left(K_n^{(\alpha)}((t - \tau)^2; \tau) \right)^{\frac{\rho}{2}}.$$

Proof. From the equation (18), we have

$$|K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau)| \leq \phi_\rho(\varphi; \tau) K_n^{(\alpha)}(|t - \tau|^\rho; \tau)$$

Applying the Hölder inequality with $a_1 = \frac{2}{\rho}$ and $a_2 = \frac{2}{2-\rho}$, we get

$$|K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau)| \leq \phi_\rho(\varphi; \tau) \left(K_n^{(\alpha)}((t - \tau)^2; \tau) \right)^{\frac{\rho}{2}}. \quad \square$$

THEOREM 14. For $\alpha > 0$, $\varphi \in C_2[0, \infty)$, $w_{a+1}(\varphi; \delta)$ is the modulus of continuity of φ on the interval $[0, a+1] \subset [0, \infty)$, $a > 0$. Then, we have

$$\left| K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| \leq 4N_\varphi(1 + a^2) \delta_n(\tau) + 2w_{a+1}\left(\varphi; \sqrt{\delta_n(\tau)}\right).$$

where $\delta_n(\tau) = K_n^{(\alpha)}((t - \tau)^2; \tau)$ given by Lemma 5.

Proof. For $0 \leq \tau \leq a$ and $a+1 < t$, since $1 < t - \tau$, we have

$$\begin{aligned} |\varphi(t) - \varphi(\tau)| &\leq M_\varphi(\tau^2 + t^2 + 2) \\ &\leq M_\varphi(2(t - \tau)^2 + 2 + 3\tau^2) \\ &\leq M_\varphi(t - \tau)^2(4 + 3\tau^2) \\ &\leq 4M_\varphi(t - \tau)^2(1 + a^2) \end{aligned} \quad (19)$$

Also, for $0 \leq \tau \leq a$ and $a+1 \geq t$, we have

$$\begin{aligned} |\varphi(t) - \varphi(\tau)| &\leq w_{a+1}(\varphi; |t - \tau|) \\ &\leq \left(1 + \frac{|t - \tau|}{\delta}\right) w_{a+1}(\varphi; \delta) \end{aligned} \quad (20)$$

with $\delta > 0$.

For $0 \leq \tau \leq a$ and $t \geq 0$, combining (19) and (20) gives

$$|\varphi(t) - \varphi(\tau)| \leq 4M_\varphi(t - \tau)^2(1 + a^2) + \left(1 + \frac{|t - \tau|}{\delta}\right) w_{a+1}(\varphi; \delta), \quad (21)$$

From the above inequality (21) and Cauchy-Schwarz's inequality, we get

$$\begin{aligned} &\left| K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| \\ &\leq K_n^{(\alpha)}(|\varphi(t) - \varphi(\tau)|; \tau) \\ &\leq 4M_\varphi(1 + a^2) K_n^{(\alpha)}((t - \tau)^2; \tau) + \left(1 + \frac{\sqrt{K_n^{(\alpha)}((t - \tau)^2; \tau)}}{\delta}\right) w_{a+1}(\varphi; \delta) \\ &\leq 4M_\varphi(1 + a^2) K_n^{(\alpha)}((t - \tau)^2; \tau) + 2w_{a+1}(\varphi; \delta_n(\tau)) \end{aligned}$$

on choosing $\delta := \delta_n(\tau) = \sqrt{K_n^{(\alpha)}((t - \tau)^2; \tau)}$. \square

4. Weighted approximation

THEOREM 15. Let $\alpha > 0$. Then for each $\varphi \in C_2^*[0, \infty)$, we have:

$$\lim_{n \rightarrow \infty} \left\| K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right\|_2 = 0.$$

Proof. By the Korovkin type theorem on the weighted approximation([15]), we need to verify the following

$$\lim_{n \rightarrow \infty} \left\| K_n^{(\alpha)}(t^m; \tau) - \tau^m \right\|_2 = 0, \quad m = 0, 1, 2. \quad (22)$$

- Since $K_n^{(\alpha)}(1; \tau) = 1$, then inequality (22) holds for $m = 0$.
- From Lemma 4, we can write:

$$\begin{aligned} \lim_{n \rightarrow \infty} \left\| K_n^{(\alpha)}(t; \tau) - \tau \right\|_2 &= \sup_{\tau \geq 0} \frac{|K_n^{(\alpha)}(t; \tau) - \tau|}{1 + \tau^2} \\ &= \sup_{\tau \geq 0} \frac{1}{1 + \tau^2} \left| \frac{1}{(\alpha + 1)n} \right| \\ &\leq \frac{1}{(\alpha + 1)n} \sup_{\tau \geq 0} \frac{1}{1 + \tau^2} \\ &\leq \frac{1}{(\alpha + 1)n} \rightarrow 0, n \rightarrow \infty \end{aligned}$$

and

$$\begin{aligned} \lim_{n \rightarrow \infty} \left\| K_n^{(\alpha)}(t^2; \tau) - \tau^2 \right\|_2 &= \sup_{\tau \geq 0} \frac{|K_n^{(\alpha)}(t^2; \tau) - \tau^2|}{1 + \tau^2} \\ &= \sup_{\tau \geq 0} \frac{1}{1 + \tau^2} \left| \frac{(\alpha + 3)}{(\alpha + 1)n} \tau + \frac{2}{(\alpha + 1)(\alpha + 2)n^2} \right| \\ &\leq \frac{(\alpha + 3)}{(\alpha + 1)n} \sup_{\tau \geq 0} \frac{\tau}{1 + \tau^2} + \frac{2}{(\alpha + 1)(\alpha + 2)n^2} \sup_{\tau \geq 0} \frac{1}{1 + \tau^2} \\ &\leq \frac{(\alpha + 3)}{(\alpha + 1)n} + \frac{2}{(\alpha + 1)(\alpha + 2)n^2} \rightarrow 0, n \rightarrow \infty. \end{aligned}$$

which implies that

$$\lim_{n \rightarrow \infty} \left\| K_{n,k}^{(\alpha)}(t^m; \tau) - \tau^m \right\|_2 = 0, \quad m = 0, 1, 2. \quad \square$$

In the next theorem, we present a weighted approximation theorem for $\varphi \in C_2^*[0, \infty)$. In [16], Doğru studied for the classical Szász operators,

THEOREM 16. *Let $\alpha > 0$. For each $\varphi \in C_2^*[0, \infty)$ and $\beta > 0$, we have*

$$\lim_{n \rightarrow \infty} \sup_{\tau \geq 0} \frac{|K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau)|}{(1 + \tau^2)^{1+\beta}} = 0.$$

Proof. Let $\tau_0 \in [0, \infty)$ be arbitrary but fixed. Then

$$\begin{aligned} \sup_{\tau \in [0, \infty)} \frac{|K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau)|}{(1 + \tau^2)^{1+\beta}} &= \sup_{\tau \in [0, \tau_0]} \frac{|K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau)|}{(1 + \tau^2)^{1+\beta}} + \sup_{\tau \in (\tau_0, \infty)} \frac{|K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau)|}{(1 + \tau^2)^{1+\beta}} \\ &\leq \|K_n^{(\alpha)}(\varphi) - \varphi\|_{C[0, \tau_0]} + \|\varphi\|_2 \sup_{\tau \in (\tau_0, \infty)} \frac{|K_{n,k}^{(\alpha)}(1 + t^2; \tau)|}{(1 + \tau^2)^{1+\beta}} \\ &\quad + \sup_{\tau \in (\tau_0, \infty)} \frac{|\varphi(\tau)|}{(1 + \tau^2)^{1+\beta}} \\ &= H_1 + H_2 + H_3. \end{aligned}$$

Since $|\varphi(\tau)| \leq N_\varphi(1 + \tau^2)$, we have

$$H_3 = \sup_{\tau \in (\tau_0, \infty)} \frac{|\varphi(\tau)|}{(1 + \tau^2)^{1+\beta}} \leq \sup_{\tau \in (\tau_0, \infty)} \frac{N_\varphi}{(1 + \tau^2)^\beta} \leq \frac{N_\varphi}{(1 + \tau_0^2)^\beta}.$$

By Theorem 15, we can get

$$\begin{aligned} H_2 &= \|\varphi\|_2 \lim_{n \rightarrow \infty} \sup_{\tau \in (\tau_0, \infty)} \frac{|K_n^{(\alpha)}(1 + t^2; \tau)|}{(1 + \tau^2)^{1+\beta}} \\ &= \sup_{\tau \in (\tau_0, \infty)} \frac{(1 + \tau^2)}{(1 + \tau^2)^{1+\beta}} \|\varphi\|_2 \\ &= \sup_{\tau \in (\tau_0, \infty)} \frac{\|\varphi\|_2}{(1 + \tau^2)^\beta} \leq \frac{\|\varphi\|_2}{(1 + \tau_0^2)^\beta}. \end{aligned}$$

Using Theorem 14, we can see that

H_1 goes to zero as $n \rightarrow \infty$.

Moreover, if we choose $\tau_0 > 0$ large enough, we can see that

$H_2 \rightarrow 0$ and $H_3 \rightarrow 0$ as $n \rightarrow \infty$,

Thus, combining, H_1, H_2 and H_3 , we get desired result. \square

In the next theorem we obtain direct estimation in terms of weighted modulus of continuity. For every $\varphi \in C_m^*[0, \infty)$ the weighted modulus of continuity defined as follows

$$\Omega_m(\varphi, \delta) = \sup_{\tau \geq 0, 0 < h \leq \delta} \frac{|\varphi(\tau + h) - \varphi(\tau)|}{1 + (\tau + h)^m}, \quad (23)$$

LEMMA 17. [18] If $\varphi \in C_m^*[0, \infty)$, $m \in \mathbb{N}$, then

- (i) $\Omega_m(\varphi, \delta)$ is a monotone increasing function of δ ,
- (ii) $\lim_{\delta \rightarrow 0^+} \Omega_m(\varphi, \delta) = 0$,
- (iii) for any $\rho \in [0, \infty)$, $\Omega_m(\varphi, \rho\delta) \leq (1 + \rho)\Omega_m(\varphi, \delta)$.

In the next theorem, we express the approximation error of $K_n^{(\alpha)}$ by using Ω_m .

THEOREM 18. *Let $m \in \mathbb{N}$. For $\varphi \in C_m^* [0, \infty)$, we have*

$$\left\| K_n^{(\alpha)}(\varphi) - \varphi \right\|_{m+1} \leq N \Omega_m(\varphi, (1/\sqrt{n})),$$

where N is a constant independent of φ and n .

Proof. From (23) and Lemma 17, we can write

$$\begin{aligned} |\varphi(t) - \varphi(\tau)| &\leq (1 + (\tau + |t - \tau|)^m) \left(\frac{|t - \tau|}{\delta} + 1 \right) \Omega_m(\varphi, \delta) \\ &\leq (1 + (2\tau + t)^m) \left(\frac{|t - \tau|}{\delta} + 1 \right) \Omega_m(\varphi, \delta). \end{aligned}$$

Then

$$\begin{aligned} &\left| K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| \\ &\leq K_n^{(\alpha)} |(\varphi(t) - \varphi(\tau))| ; \tau \\ &\leq \Omega_m(\varphi, \delta) \left(K_n^{(\alpha)}((1 + (2\tau + t)^m); \tau) + K_n^{(\alpha)} \left((1 + (2\tau + t)^m) \frac{|t - \tau|}{\delta}; \tau \right) \right) . \\ &= \Omega_m(\varphi, \delta) \left(K_n^{(\alpha)}(1 + (2\tau + t)^m; \tau) + I_1 \right) . \end{aligned}$$

Applying Cauchy-Schwartz inequality to the I_1 , we get

$$I_1 \leq (K_n^{(\alpha)}((1 + (2\tau + t)^m)^2; \tau))^{1/2} \left(K_n^{(\alpha)} \left(\frac{|t - \tau|^2}{\delta^2}; \tau \right) \right)^{1/2} .$$

Therefore,

$$\begin{aligned} &\left| K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| \\ &\leq \Omega_m(\varphi, \delta) \left(K_n^{(\alpha)}((1 + (2\tau + t)^m); \tau) + (K_n^{(\alpha)}((1 + (2\tau + t)^m)^2; \tau))^{1/2} \right. \\ &\quad \times \left. \left(K_n^{(\alpha)} \left(\frac{|t - \tau|^2}{\delta^2}; \tau \right) \right)^{1/2} \right) . \end{aligned} \tag{24}$$

By Lemma 8 and 7,

$$\begin{aligned} K_n^{(\alpha)}(1 + (2\tau + t)^m; \tau) &\leq C_m(\alpha)(1 + \tau^m), \\ (K_n^{(\alpha)}((1 + (2\tau + t)^m)^2; \tau))^{1/2} &\leq C_m^1(\alpha)(1 + \tau^m) . \end{aligned} \tag{25}$$

and

$$\left(K_n^{(\alpha)} \left(\frac{|t-\tau|^2}{\delta^2}; \tau \right) \right)^{1/2} \leq \frac{1}{\delta} \sqrt{\frac{2}{(\alpha+1)(\alpha+2)n^2}} + \frac{\tau}{n} \leq \frac{(1+\tau)}{\delta\sqrt{n}}. \quad (26)$$

Combining (25), (25) and (26), we have

$$\begin{aligned} \left| K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right| &\leq \Omega_m(\varphi, \delta) \left(C_m(\alpha)(1+\tau^m) + C_m^1(\alpha) \frac{(1+\tau^m)(1+\tau)}{\delta\sqrt{n}} \right) \\ &= \Omega_m(\varphi, \delta) \left(C_m(\alpha)(1+\tau^m) + C_m^1(\alpha)C_1 \frac{(1+\tau^{m+1})}{\delta\sqrt{n}} \right), \end{aligned}$$

where

$$C_1 = \sup_{\tau \geq 0} \frac{(1+\tau^m + \tau + \tau^{m+1})}{1 + \tau^{m+1}}.$$

if we take $\delta = (1/\sqrt{n})$ in the above inequality, we obtain the desired result. \square

Next result is a Voronovskaja type formula for the operators $K_n^{(\alpha)}(\varphi; \tau)$.

5. Voronovskaja type results

THEOREM 19. Let $\alpha > 0$. For any $\varphi \in C_2^*[0, \infty)$ such that $\varphi', \varphi'' \in C_2^*[0, \infty)$ the following equality holds

$$\lim_{n \rightarrow \infty} n \left[K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right] = \frac{1}{(\alpha+1)} \varphi'(\tau) + \frac{1}{2} \tau \varphi''(\tau).$$

Proof. By the Taylor's formula, we can write

$$\varphi(t) = \varphi(\tau) + \varphi'(\tau)(t-\tau) + \frac{1}{2} \varphi''(\tau)(t-\tau)^2 + r(t, \tau)(t-\tau)^2 \quad (27)$$

where $r(t, \tau)$ is Peano form of remainder, $r(\cdot, \tau) \in C_2^*[0, \infty)$ and $\lim_{t \rightarrow \tau} r(t, \tau) = 0$.

Applying $K_n^{(\alpha)}$ to the both sides of (27), we get

$$\begin{aligned} &n \left(K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right) \\ &= \varphi'(\tau) n K_n^{(\alpha)}((t-\tau); \tau) + \frac{1}{2} \varphi''(\tau) n K_n^{(\alpha)}((t-\tau)^2; \tau) + n K_n^{(\alpha)}(r(t, \tau)(t-\tau)^2; \tau). \end{aligned}$$

By Cauchy-Schwarz inequality, we have

$$K_n^{(\alpha)}(r(t, \tau)(t-\tau)^2; \tau) \leq \sqrt{K_n^{(\alpha)}(r^2(t, \tau); \tau)} \sqrt{K_n^{(\alpha)}((t-\tau)^4; \tau)}. \quad (28)$$

Observe that $r^2(t, \tau) = 0$ and $r^2(\cdot, \tau) \in C_2^*[0, \infty)$.

Then, it follows from that Corollary 11,

$$\lim_{n \rightarrow \infty} n K_n^{(\alpha)}(r^2(t, \tau); \tau) = r^2(\tau, \tau) = 0. \quad (29)$$

Moreover, from (7), (28) and (29), we can obtain

$$\lim_{n \rightarrow \infty} K_n^{(\alpha)}(r(t, \tau)(t - \tau)^2; \tau) = 0 \quad (30)$$

Hence, combining (5), (6) and (30), we get

$$\lim_{n \rightarrow \infty} n \left[K_n^{(\alpha)}(\varphi; \tau) - \varphi(\tau) \right] = \frac{1}{(\alpha + 1)} \varphi'(\tau) + \frac{1}{2} \tau \varphi''(\tau). \quad \square$$

6. Bivariate Riemann-Liouville fractional integral type Szász-Mirakyan-Kantorovich operators

In this section, we introduce the bivariate Riemann-Liouville fractional integral type of $K_n^{(\alpha)}(\varphi; \tau)$ (3) as follows:

$$K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(\varphi; \tau, \gamma) = \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \alpha_1 \alpha_2 s_{n_1, k_1}(\tau) s_{n_2, k_2}(\gamma) \int_0^1 \int_0^1 \varphi\left(\frac{k_1+t_1}{n_1}, \frac{k_2+t_2}{n_2}\right) (1-t_1)^{\alpha_1-1} (1-t_2)^{\alpha_2-1} dt_1 dt_2$$

where $(\tau, \gamma) \in I^2 = [0, \infty) \times [0, \infty)$ and $\alpha_1, \alpha_2 > 0$.

Bivariate Riemann-Liouville fractional integral type Szász-Mirakyan-Kantorovich operators can be rewritten as

$$K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(\cdot; \tau, \gamma) = K_{n_1}^{\alpha_1}(\cdot; \tau) \times K_{n_2}^{\alpha_2}(\cdot; \gamma)$$

LEMMA 20. Let $e_{ij}(\tau, \gamma) = \tau^i \gamma^j$, $0 \leq i+j \leq 2$ and $\alpha_1, \alpha_2 > 0$. For $(\tau, \gamma) \in I^2 = [0, \infty) \times [0, \infty)$, we have

$$\begin{aligned} K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(e_{00}; \tau, \gamma) &= 1, \\ K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(e_{10}; \tau, \gamma) &= \tau + \frac{1}{n_1(\alpha_1 + 1)}, \\ K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(e_{10}; \tau, \gamma) &= \gamma + \frac{1}{n_2(\alpha_2 + 1)}, \\ K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(e_{20}; \tau, \gamma) &= \frac{2}{(\alpha_1 + 1)(\alpha_1 + 2)n_1^2} + \frac{(\alpha_1 + 3)}{(\alpha_1 + 1)n_1} \tau + \tau^2, \\ K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(e_{02}; \tau, \gamma) &= \frac{2}{(\alpha_2 + 1)(\alpha_2 + 2)n_2^2} + \frac{(\alpha_2 + 3)}{(\alpha_2 + 1)n_2} \tau + \tau^2. \end{aligned}$$

LEMMA 21. According to above Lemma 20, we get

$$\begin{aligned} K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(e_{10} - \tau; \tau, \gamma) &= \frac{1}{(\alpha_1 + 1)(n_1 + 1)}, \\ K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(e_{01} - \gamma; \tau, \gamma) &= \frac{1}{(\alpha_2 + 1)(n_2 + 1)}, \\ K_{n_1, n_2}^{(\alpha_1, \alpha_2)}((e_{10} - \tau)^2; \tau, \gamma) &= \frac{2}{(\alpha_1 + 1)(\alpha_1 + 2)n_1^2} + \frac{\tau}{n_1} = \delta_{n_1}^{(\alpha_1)}(\tau), \\ K_{n_1, n_2}^{(\alpha_1, \alpha_2)}((e_{01} - \gamma)^2; \tau, \gamma) &= \frac{2}{(\alpha_2 + 1)(\alpha_2 + 2)n_2^2} + \frac{\gamma}{n_2} = \delta_{n_2}^{(\alpha_2)}(\gamma). \end{aligned}$$

For bivariate real functions, modulus of continuity is given by

$$w(\varphi; \delta_n, \delta_m) = \sup \{ |\varphi(t, s) - \varphi(\tau, \gamma)| : (t, s), (\tau, \gamma) \in I^2, |t - \tau| \leq \delta_n, |s - \gamma| \leq \delta_m \}.$$

THEOREM 22. Let $\varphi \in C(I^2)$ and $\alpha_1, \alpha_2 > 0$. Then for all $(\tau, \gamma) \in I^2$, the inequality

$$\left| K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(\varphi; \tau, \gamma) - \varphi(\tau, \gamma) \right| \leq 4w\left(\varphi; \delta_{n_1}^{(\alpha_1)}(\tau), \delta_{n_2}^{(\alpha_2)}(\gamma)\right)$$

holds, where $\delta_{n_1}^{(\alpha_1)}(\tau), \delta_{n_2}^{(\alpha_2)}(\gamma)$ are as in Lemma 21.

Proof. By the positivity and linearity properties of the $K_{n_1, n_2}^{(\alpha_1, \alpha_2)}$, we can write

$$\begin{aligned} \left| K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(\varphi; \tau, \gamma) - \varphi(\tau, \gamma) \right| &\leq K_{n_1, n_2}^{(\alpha_1, \alpha_2)}(|\varphi(t, s) - \varphi(\tau, \gamma)|; \tau, \gamma) \\ &\leq w(\varphi; \delta_1, \delta_2) \left(K_{n_1}^{(\alpha_1)}(1; \tau) + \frac{1}{\delta_1} K_{n_2}^{(\alpha_2)}(|t - \tau|; \tau) \right) \\ &\quad \times \left(K_{n_2}^{(\alpha_2)}(1; \gamma) + \frac{1}{\delta_2} K_{n_2}^{(\alpha_2)}(|s - \gamma|; \gamma) \right) \end{aligned}$$

Applying Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} K_{n_1}^{(\alpha_1)}(|t - \tau|; \tau) &\leq K_{n_1}^{(\alpha_1)}((t - \tau)^2; \tau)^{\frac{1}{2}} \\ K_{n_2}^{(\alpha_2)}(|s - \gamma|; \gamma) &\leq K_{n_2}^{(\alpha_2)}((s - \gamma)^2; \gamma)^{\frac{1}{2}} \end{aligned}$$

Choosing $\delta_1 = \delta_{n_1}^{(\alpha_1)}(\tau)$ and $\delta_2 = \delta_{n_2}^{(\alpha_2)}(\gamma)$, we have desired result. \square

7. Graphical simulations

In this section, we present some graphical simulations and numerical results for $K_n^{(\alpha)}$ and $K_{n_1, n_2}^{(\alpha_1, \alpha_2)}$ obtained by using Matlab. This simulations agree with our theoretical results.

EXAMPLE 23. Consider $f(x) = \begin{cases} -6x^3 + 9x^2 - (\frac{66}{25})x, & x \in [0, 1] \\ \frac{9}{25}, & x > 1 \end{cases}$. Here we take the value of $n \in \{50, 100, 150\}$ and $\alpha = 10$ for $K_n^{(\alpha)}(f; x)$ with $x \in [0, 1]$. The Figure 1 demonstrate the convergence of operators $K_n^{(\alpha)}$ to $f(x)$ for fixed α and increasing values of n . Then, some numerical values of $E_n^\alpha(f; x) = |K_n^{(\alpha)}(f; x) - f(x)|$ at certain points on the interval $[0, 1]$ for $n \in \{50, 100, 150\}$ and $\alpha = 10$ are given in Table 1.




Figure 1: Approximation to f by $K_n^{(\alpha)}(f; x)$ for $\alpha = 10$, $f(x) = -6x^3 + 9x^2 - \frac{66}{25}x$ and $n = 50, 100, 150$.

x	$ K_{50}^{10}(f; x) - f(x) $	$ K_{100}^{10}(f; x) - f(x) $	$ K_{150}^{10}(f; x) - f(x) $
0	0.004745622	0.002386385	0.001593946
0.1	0.012283469	0.006207252	0.004152721
0.2	0.021458014	0.010873615	0.007281206
0.3	0.022778014	0.011612706	0.007791509
0.4	0.016243469	0.008424524	0.00568363
0.5	0.001854378	0.00130907	0.00095757
0.6	0.020389259	0.009733657	0.006386673
0.7	0.050487441	0.024703657	0.016349097
0.8	0.088440168	0.04360093	0.028929703
0.9	0.134247441	0.066425476	0.044128491
1	0.187909259	0.093177294	0.061945461

Table 1: Estimation of the absolute error function E_n^{10} with $f(x) = -6x^3 + 9x^2 - \frac{66}{25}x$ for some values of x in $[0, 1]$ and $n \in \{50, 100, 150\}$.

As we increase the value of n and fixed α , the approximation is good, i.e for the largest value of n and fixed α , the error is minimum.

EXAMPLE 24. Let $f(x) = \begin{cases} -6x^3 + 9x^2 - \left(\frac{66}{25}\right)x, & x \in [0, 0.5] \\ \frac{117}{100}, & x > 0.5 \end{cases}$. Here we take the value of $\alpha \in \{0.5, 1, 5\}$ and $n = 50$ with $x \in [0.2, 0.5]$. The Figure 2 demonstrate the convergence of operators $K_n^{(\alpha)}$ to $f(x)$ for increasing values of α and fixed n . Secondly, the absolute error function $E_n^\alpha(f; x) = |K_n^{(\alpha)}(f; x) - f(x)|$ is illustrated in figure 3. Finally, some numerical values of $E_n^\alpha(f; x)$ at certain points on the interval $[0.2, 0.5]$ for $\alpha \in \{0.5, 1, 5\}$ and $n = 50$ are given in Table 2.




Figure 2: Approximation to f by $K_n^{(\alpha)}(f; x)$ for $n = 50$, $f(x) = -6x^3 + 9x^2 - \frac{66}{25}x$ and $\alpha = 0.5, 1, 5$.

x	$ K_{50}^{0.5}(f; x) - f(x) $	$ K_{50}^1(f; x) - f(x) $	$ K_{50}^5(f; x) - f(x) $
0.2	0.024490057	0.023508	0.021782
0.3	0.035386057	0.031668	0.024387714
0.4	0.034282057	0.029028	0.018593429
0.5	0.021178057	0.015588	0.004399143

Table 2: Estimation of the absolute error function $E_{50}^{(\alpha)}$ with $f(x) = -6x^3 + 9x^2 - \frac{66}{25}x$ for some values of x in $[0.2, 0.5]$ and $\alpha \in \{0.5, 1, 5\}$.

Now, we are present some graphs and numerical results for the convergence of bivariate Riemann-Liouville integral type Szász-Mirakyan-Kantorovich operators $K_{n_1, n_2}^{(\alpha_1, \alpha_2)}$ by considering the function $f(x, y) = x + xy + 12y^2$.




Figure 3: Absolute error function $E_n^{(\alpha)}(f; x)$ for $\alpha = 10$, $f(x) = -6x^3 + 9x^2 - \frac{66}{25}x$ and $n = 50$.

EXAMPLE 25. Consider, $f(x, y) = \begin{cases} x + xy + 12y^2, & (x, y) \in [0, 1] \times [0, 1], \\ 14, & (x, y) \in (1, \infty) \times (1, \infty). \end{cases}$

Here we take the value of $n_1, n_2 \in \{10, 20\}$ and $\alpha_1 = \alpha_2 = 10$ with $(x, y) \in [0, 1] \times [0, 1]$. The Figure 4 explains the convergence of the operators $K_{n_1, n_2}^{(\alpha_1, \alpha_2)}$ towards the




Figure 4: Convergence of the operators $K_{n_1, n_2}^{(10, 10)}$ to the function $f(x, y) = x + xy + 12y^2$ for the increasing values of $n_1, n_2 \in \{10, 20\}$ on the interval $[0, 1] \times [0, 1]$.

function $f(x,y)$ for increasing values of n_1, n_2 and fixed α_1 and α_2 . Then, some numerical values of $E_n^\alpha(f;x) = |K_n^{(\alpha)}(f;x) - f(x)|$ at certain points on the interval $[0, 1] \times [0, 1]$ for $n_1, n_2 \in \{10, 20\}$ and $\alpha_1 = \alpha_2 = 10$ are given in Table 3.

τ	y	$ K_{10,10}^{10,10}(f;\tau) - f(\tau) $	$ K_{20,20}^{10,10}(f;\tau) - f(\tau) $
1	0.1	0.162809917	0.080929752
1	0.2	0.30553719	0.152293388
1	0.3	0.448264463	0.223657025
1	0.4	0.590991736	0.295020661
1	0.5	0.733719008	0.366384298
1	0.6	0.876446281	0.437747934
1	0.7	1.019173554	0.50911157
1	0.8	1.161900826	0.580475207
1	0.9	1.304628099	0.651838843
1	1	1.447355372	0.723202479

Table 3: Estimation of the absolute error function $E_{n_1, n_2}^{(10,10)}$ with $f(x,y) = x + xy + 12y^2$ for some values of (x,y) in $[0, 1] \times [0, 1]$ and $n_1, n_2 \in \{10, 20\}$.

As we increase the value of n_1 and n_2 and fixed α_1 and α_2 , the approximation is good, i.e for the largest value of n_1 and n_2 and fixed α_1 and α_2 , the error is minimum.




Figure 5: Convergence of the operators $K_{10,10}^{(\alpha_1,\alpha_2)}(f;x,y)$ to the function $f(x,y) = x + xy + 12y^2$ for the increasing values of $\alpha_1, \alpha_2 \in \{0.1, 1, 5\}$ on the interval $[0, 1] \times [0, 1]$.

EXAMPLE 26. Consider $f(x,y) = \begin{cases} x+xy+12y^2, & (x,y) \in [0,1] \times [0,1], \\ 14, & (x,y) \in (1,\infty) \times (1,\infty). \end{cases}$

Here we take the value of $\alpha_1, \alpha_2 \in \{0.1, 1, 5\}$ and $n_1 = n_2 = 10$ with $(x,y) \in [0,1] \times [0,1]$. The Figure 5 explains the convergence of the operators $K_{n_1,n_2}^{(\alpha_1,\alpha_2)}$ towards the function $f(x,y)$ for increasing values of $\alpha_1, \alpha_2 \in \{0.1, 1, 5\}$ and fixed n_1 and n_2 . Then, some numerical values of $E_{n_1,n_2}^{(\alpha_1,\alpha_2)}(f;x,y)$ at certain points on the interval $[0,1] \times [0,1]$ for $n_1 = n_2 = 10$ and $\alpha_1, \alpha_2 \in \{0.1, 1, 5\}$ are given in Table 4.

x	y	$ K_{10,10}^{0.1,0.1}(f;x,y) - f(x,y) $	$ K_{10,10}^{1,1}(f;x,y) - f(x,y) $	$ K_{10,10}^{5,5}(f;x,y) - f(x,y) $
1	0.1	0.641251476	0.3875	0.200992063
1	0.2	0.988524203	0.6325	0.36265873
1	0.3	1.33579693	0.8775	0.524325397
1	0.4	1.683069658	1.1225	0.685992063
1	0.5	2.030342385	1.3675	0.84765873
1	0.6	2.377615112	1.6125	1.009325397
1	0.7	2.724887839	1.8575	1.170992063
1	0.8	3.072160567	2.1025	1.33265873
1	0.9	3.419433294	2.3475	1.494325397
1	1	3.766706021	2.5925	1.655992063

Table 4: Estimation of the absolute error function $E_{10,10}^{(\alpha_1,\alpha_2)}$ with $f(x,y) = x+xy+12y^2$ for some values of (x,y) in $[0,1] \times [0,1]$ and $\alpha_1, \alpha_2 \in \{0.1, 1, 5\}$.

As we increase the value of α_1 and α_2 and fixed n_1 and n_2 , the approximation is good, i.e for the largest value of α_1 and α_2 and fixed n_1 and n_2 , the error is minumum.

REFERENCES

- [1] Z. DITZIAN, V. TOTIK, *Moduli of Smoothness*, Springer Series in Computational Mathematics, vol. 9, New-York: Springer, 1987.
- [2] O. DUMAN, M. A. ÖZARSLAN, B. DELLA VECCHIA, *Modified Szász-Mirakyan-Kantorovich operators preserving linear functions*, Turk J Math. 2009; 33: 151–158.
- [3] A. ARAL, D. INOAN, I. RASA, *On the generalized Szász-Mirakyan operators*, Results Math. 2014; 65: 441–452.
- [4] T. ACAR, A. ARAL, D. CÁRDENAS-MORALES, P. GARRANCHO, *Szász-Mirakyan type operators which fix exponentials*, Results Math. 2017; 72 (3): 1393–1404.
- [5] T. ACAR, A. ARAL, H. GONSKA, *On Szász-Mirakyan operators preserving e^{2ax} , $a > 0$* , Mediterr. J. Math. 2017; 14 (1): Art. 6, <https://doi.org/10.1007/s00009-016-0804-7>.
- [6] B. D. BOYANOV, V. M. VESELINOV, *A note on the approximation of functions in an infinite interval by linear positive operators*, Bull Math SocSci Math Roum. 1970; 14 (62): 9–13.
- [7] V. GUPTA, A. ARAL, *A note on Szász-Mirakyan-Kantorovich type operators preserving e^{-x} Positivity*, 22: 415–423, <https://doi.org/10.1007/s11117-017-0518-5>.
- [8] O. SZÁSZ, *Generalization of S. Bernstein's polynomials to the infinite interval*, J. Res. Nat. Bur. Standards 45 (1950), 239–245.

- [9] G. M. MIRAKJAN, *Approximation of continuous functions with the aid of polynomials*, In Dokl. Acad. Nauk SSSR 1941, Vol. 31, pp. 201–205.
- [10] P. L. BUTZER, *On the extensions of Bernstein polynomials to the infinite interval*, Proceedings of the American Mathematical Society 1954 Aug 1; 5 (4): 547–553.
- [11] P. N. AGRAWAL, H. S. KASANA, *On simultaneous approximation by Szasz–Mirakjan operators*, Bull. Inst. Math. Acad. Sinica 22 (2) (1994) 181–188.
- [12] V. TOTIK, *Uniform approximation by Szasz–Mirakjan type operators*, Acta Math. Hungar 41 (3–4) (1983) 291–307.
- [13] R. A. DEVORE, G. G. LORENTZ, *Constructive Approximation*, Grundlehren der Mathematischen Wissenschaften, Springer-Verlang, New York-London, (1993).
- [14] B. LENZE, *On Lipschitz type maximal functions and their smoothness spaces*, Nederl. Akad. Indag. Math., 50 (1988), 53–63.
- [15] A. D. GADJIEVA, *A problem on the convergence of a sequence of positive linear operators on unbounded sets, and theorems that are analogous to P. P. Korovkin's theorem*, Doklady Akademii Nauk SSSR, vol. 218, pp. 1001–1004, 1974 (Russian), Soviet Mathematics. Doklady, vol. 15, pp. 1433–1436, 1974.
- [16] OGÜN DOĞRU AND ELMIRA GADJIEVA, *Ağırlıklı uzaylarda Szász tipinde operatörler dizisinin sürekli fonksiyonlara yaklaşımı*, II, Kızılırmak Uluslararası Fen Bilimleri Kongresi Bildiri Kitabı, Kırıkkale, 29–37, 1998.
- [17] M. BECKER, *Global approximation theorems for Szász–Mirakjan and Baskakov operators in polynomial weight spaces*, Indiana Univ. Math. J., 27 (1) (1978), 127–142.
- [18] A.-J. LOPEZ-MORENO, *Weighted simultaneous approximation with Baskakov type operators*, Acta Mathematica Academiae Scientiarum Hungaricae, July 2004, doi:10.1023/B:AMHU.0000034368.81211.23.
- [19] V. I. VOLKOV, *On the convergence of sequences of linear positive operators in the space of continuous functions of two variables*, Dokl. Akad. Nauk SSSR, 115: 1 (1957), 17–19.
- [20] M. DHAMIJA, R. PRATAP, N. DEO, *Approximation by Kantorovich form of modified Szasz–Mirakyan operators*, Appl. Math. Comput. 317, 109–120 (2018).
- [21] V. GUPTA, A. M. ACU, *On Baskakov-Szász–Mirakyan-type operators preserving exponential type functions*, Posit. 22 (3), 919–929 (2018).
- [22] M. ISMAIL, *On a generalization of Szász operators*, Mathematica (Cluj) 39, 259–267 (1974).
- [23] A. KAJLA, A. M. ACU, P. N. AGRAWAL, et al., *Baskakov-Szász-type operators based on inverse Pólya Eggenberger distribution*, Ann. Funct. Anal. 8 (1), 106–123 (2017).
- [24] M. MURSALEEN, A. ALOTAIBI, K. J. ANSARI, *On a Kantorovich variant of Szász–Mirakjan operators*, J. Funct. Spaces, 2016, (2016).
- [25] B. WOOD, *Generalized Szasz operators for the approximation in the complex domain*, SIAM J. Appl. Math. 17 (4), 790–801 (1969).
- [26] T. ACAR, V. GUPTA AND A. ARAL, *Rate of convergence for generalized Szász operators*, Bull. Math. Sci. 1 (1) (2011), 99–113, doi:10.1007/s13373-011-0005-4.
- [27] P. N. AGRAWAL, V. GUPTA, A. S. KUMAR AND A. KAJLA, *Generalized Baskakov-Szász type operators*, Appl. Math. Comput. 236 (2014), 311–324, doi:10.1016/j.amc.2014.03.084.
- [28] A. ARAL, *A generalization of Szász–Mirakyan operators based on q -integers*, Math. Comput. Modelling 47 (9–10) (2008), 1052–1062, doi:10.1016/j.mcm.2007.06.018.
- [29] Z. FINTA, N. K. GOVIL AND V. GUPTA, *Some results on modified Szász–Mirakjan operators*, J. Math. Anal. Appl. 327 (2007), 1284–1296, doi:10.3906/mat-0801-2.
- [30] S. M. MAZHAR AND V. TOTIK, *Approximation by modified Szász operators*, Acta Sci. Math. 49 (1985), 257–269.
- [31] V. TOTIK, *Approximation by Szász–Mirakjan–Kantorovich operators in $Lp(p > 1)$* , Analysis Mathematica. 1983 Jun 1; 9 (2): 147–167.
- [32] Z. DAHMANI, L. TABHARIT AND S. TAF, *New generalizations of Grüss inequality using Riemann–Liouville type integrals*, Bull. Math. Anal. Appl., 2 (3) (2010), 93–99.
- [33] U. N. KATUGOMPOLA, *New Approach Generalized Fractional Integral*, Applied Math and Comp. 218 (2011), 860–865.
- [34] M. A. LATIF AND S. HUSSAIN, *New inequalities of Ostrowski type for co-ordinated convex functions via fractional integrals*, Journal of Fractional Calculus and Applications, Vol. 2., 2012, No. 9, pp. 1–15.

- [35] L. G. ROMERO, L. L. LUQUE, G. A. DORREGO AND R. A. CERUTTI, *On the k -Riemann Liouville Fractional Derivative*, Int. J. Contemp. Math. Sciences, 8 (1), 2013, 41–51.
- [36] M. TUNC, *On new inequalities for h -convex functions via Riemann-Liouville fractional integration*, Filomat 27: 4 (2013), 559–565.

(Received February 15, 2022)

Nazim Mahmudov
Department of Mathematics
Eastern Mediterranean University
Gazimagusa 99628, T.R. Northen Cyprus, Mersin 10, Turkey

Mustafa Kara
Department of Mathematics
Eastern Mediterranean University
Gazimagusa 99628, T.R. Northen Cyprus, Mersin 10, Turkey