

# HERMITE–HADAMARD TYPE INEQUALITIES FOR RIEMANN–LIOUVILLE FRACTIONAL INTEGRALS VIA STRONGLY $h$ -CONVEX FUNCTIONS

YI XING, CHAOQUN JIANG AND JIANMIAO RUAN\*

*(Communicated by J. Pečarić)*

**Abstract.** In this paper, the Hermite-Hadamard type inequalities for Riemann-Liouville fractional integrals via strongly  $h$ -convex functions are established. Furthermore, we obtain some identities related to the fractional integrals with  $n$ -times differentiable functions, and then gain midpoint type and trapezoid type error estimates connected with the Hermite-Hadamard type inequalities, which generalize some known results.

## 1. Introduction

Let  $I$  be an interval in  $\mathbb{R}$  and  $h : [0, 1] \rightarrow [0, \infty)$  be a given function. A function  $f : I \rightarrow \mathbb{R}$  is called  $h$ -convex provided that

$$f(tx + (1-t)y) \leq h(t)f(x) + h(1-t)f(y) \quad (1.1)$$

holds for all  $x, y \in I$  and  $t \in (0, 1)$ . This notation was introduced by Varošanec [33] and generalizes the classes of *convex functions*, *s-convex functions (in the second sense)*, *Godunova-Levin functions* and *P-functions*, which are obtained by taking in (1.1)  $h(t) = t$ ,  $h(t) = t^s$  ( $s \in (0, 1)$ ),  $h(t) = 1/t$  and  $h(t) = 1$ , respectively. Many properties of them can be found, for instance, in [10, 12, 16, 25, 26, 38].

A significant application of the convex function is the well-known Hermite-Hadamard inequality, if  $f : [a, b] \rightarrow \mathbb{R}$  is a convex function, then

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a) + f(b)}{2}. \quad (1.2)$$

This inequality was studied extensively and had been extended under various convex type functions. In 1995, Dragomir, Pečarić and Persson [11] established similar results for Godunova-Levin functions and *P-functions*. In 1999, Dragomir and Fitzpatrick [9]

---

*Mathematics subject classification (2020): Primary 26A33; Secondary 26A51, 26D15.*

*Keywords and phrases:* Hermite-Hadamard type inequalities, strongly  $h$ -convex functions, Riemann-Liouville fractional integrals.

The research was supported by the College Students' Science and Technology Innovation Project and Xinmiao Project of Zhejiang Province (No. 2021R419010) and the National Natural Science Foundation of China (No. 11771358).

\* Corresponding author.

obtained an analogous inequality for  $s$ -convex functions (in the second sence). In 2008, Sarikaya, Saglam and Yildirim [29] extended it to  $h$ -convex functions.

Following Polyak [23], a function  $f : I \rightarrow \mathbb{R}$  is said to be *strongly convex with modulus  $\beta > 0$*  if

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y) - \beta t(1-t)(x-y)^2, \quad (1.3)$$

for all  $x, y \in I$  and  $t \in (0, 1)$ . The function played an important role in optimization theory, mathematical economics and so on (see e.g. [17, 18, 19, 23, 24, 34, 35]). In 2011, Angulo, Gimenez, Moros and Nikodem [3] introduced the strongly  $h$ -convex function, which unified the classes of strongly convex functions and  $h$ -convex functions. And then they extended (1.2) to these new functions.

**DEFINITION 2.** [3] Let  $h : [0, 1] \rightarrow [0, \infty)$  be a given function and  $\beta$  be a positive constant. We say that  $f : I \rightarrow \mathbb{R}$  is strongly  $h$ -convex with modulus  $\beta$ , or  $f$  belongs to the class  $SX(h, \beta, I)$ , if

$$f(tx + (1-t)y) \leq h(t)f(x) + h(1-t)f(y) - \beta t(1-t)(x-y)^2, \quad (1.4)$$

for all  $x, y \in I$  and  $t \in (0, 1)$ .

Particularly, if  $f$  satisfies (1.4) with  $h(t) = t$ ,  $h(t) = t^s$  ( $s \in (0, 1)$ ),  $h(t) = 1/t$  and  $h(t) = 1$ , then  $f$  is said to be *strongly convex functions*, *strongly  $s$ -convex functions*, *strongly Godunova-Levin functions* and *strongly P-function*, respectively. Moreover, it is not difficult to check that  $h(1/2) > 0$  if  $f \in SX(h, \beta, I)$  and  $f \geq 0$ . As an application, the authors [3] established the following Hermite-Hadamard inequality.

**THEOREM A.** Let  $f \in SX(h, \beta, [a, b])$  and  $h$  be Lebesgue integrable on  $(0, 1)$  with  $h(1/2) > 0$ . If  $f$  is Lebesgue integrable on  $[a, b]$ , then

$$\begin{aligned} \frac{1}{2h(1/2)} \left[ f\left(\frac{a+b}{2}\right) + \frac{\beta}{12}(b-a)^2 \right] &\leq \frac{1}{b-a} \int_a^b f(x)dx \\ &\leq (f(a) + f(b)) \int_0^1 h(t)dt - \frac{\beta}{6}(b-a)^2. \end{aligned} \quad (1.5)$$

It is notable that Theorem A reduces to Theorem 6 in [29] with  $\beta \rightarrow 0$ .

Let  $\alpha > 0$  and  $f \in L([a, b])$ . The left-sided and the right-sided Riemann-Liouville fractional integrals  $J_{a+}^\alpha f$  and  $J_{b-}^\alpha f$  of order  $\alpha$  are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t)dt, \quad x \in (a, b], \quad (1.6)$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t)dt, \quad x \in [a, b), \quad (1.7)$$

respectively, where  $\Gamma(\alpha) = \int_0^\infty t^{\alpha-1} e^{-t} dt$  is the Gamma function.

In recent years, Hermite-Hadamard type inequalities via the fractional integrals are studied extensively, for instance, see [5, 8, 27, 30, 32, 36, 37] and the references therein.

In 2017, Sarikaya and Yildirim [28] first obtained a remarkable inequality of Hermite-Hadamard type involving the fractional integrals.

**THEOREM B.** [28] Let  $f : [a, b] \rightarrow \mathbb{R}$  be a convex function and  $f \in L([a, b])$ . Then

$$f\left(\frac{a+b}{2}\right) \leq \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{(\frac{a+b}{2})^-}^\alpha f(a) + J_{(\frac{a+b}{2})^+}^\alpha f(b) \right] \leq \frac{f(a) + f(b)}{2}.$$

In 2020, Budak, Ertugral and Sarikaya [6] extended it to more generalized fractional integrals. In 2021, Zhang, Farid and Akbar [39] obtained an analogue inequality as Theorem B for strongly  $(s, m)$ -convex functions.

In 2000, Pearce and Pečarić [25] proved an important equality connect with the left part of (1.2).

**LEMMA A.** [25] Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function and  $f' \in L([a, b])$ . Then

$$\begin{aligned} & \frac{1}{b-a} \int_a^b f(x) dx - f\left(\frac{a+b}{2}\right) \\ &= \frac{1}{b-a} \left[ \int_{(a+b)/2}^b (b-x)f'(x) dx - \int_a^{(a+b)/2} (x-a)f'(x) dx \right] \\ &= (b-a) \left[ \int_0^{1/2} t f'(ta + (1-t)b) dt - \int_{1/2}^1 (1-t) f'(ta + (1-t)b) dt \right]. \end{aligned}$$

By the lemma, the authors showed the following result.

**THEOREM C.** [25] Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function. If  $|f'|^q$  is convex on  $[a, b]$  with  $1 \leq q < \infty$ , then

$$\left| \frac{1}{b-a} \int_a^b f(x) dx - f\left(\frac{a+b}{2}\right) \right| \leq \frac{b-a}{4} \left( \frac{|f'(a)|^q + |f'(b)|^q}{2} \right)^{1/q}. \quad (1.8)$$

Furthermore, some estimates for concave functions are also achieved in [15] and [25]. In 2004, Kirmaci [14] rediscovered Lemma A and established some other estimates similar to Theorem C. In 2011, Alomori, Darus and Kirmaci [2] obtained analogue results for  $s$ -convex functions. And more results about the difference estimates connected with (1.2) can be referred, for instance, to [4, 13, 25].

In 2017, Sarikaya and Yildirim [28] found an important identity related to Riemann-Liouville integrals as follows.

**LEMMA B.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function and  $f' \in L([a, b])$ . Then

$$\begin{aligned} & \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{(\frac{a+b}{2})^-}^\alpha f(a) + J_{(\frac{a+b}{2})^+}^\alpha f(b) \right] - f\left(\frac{a+b}{2}\right) \\ &= \frac{b-a}{4} \int_0^1 (1-t)^\alpha \left[ f'\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) - f'\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) \right] dt. \end{aligned}$$

It is not difficult to check that Lemma B becomes Lemma A with  $\alpha = 1$ . As a consequence, they obtained the following midpoint type inequalities for differentiable functions.

**THEOREM D.** [28] Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable function. Suppose that  $|f'|^q$  is convex on  $[a, b]$  for  $1 \leq q < \infty$ .

(i) If  $1 \leq q < \infty$ , then

$$\begin{aligned} & \left| \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - f\left(\frac{a+b}{2}\right) \right| \\ & \leq \frac{b-a}{4(\alpha+1)} \left( \frac{1}{2(\alpha+2)} \right)^{1/q} \left\{ [(\alpha+1)|f'(a)|^q + (\alpha+3)|f'(b)|^q]^{1/q} \right. \\ & \quad \left. + [(\alpha+3)|f'(a)|^q + (\alpha+1)|f'(b)|^q]^{1/q} \right\}. \end{aligned}$$

(ii) If  $1 < q < \infty$ , then

$$\begin{aligned} & \left| \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - f\left(\frac{a+b}{2}\right) \right| \\ & \leq \frac{b-a}{4} \left( \frac{1}{\alpha p+1} \right)^{1/p} \left[ \left( \frac{|f'(a)|+3|f'(b)|}{4} \right)^{1/q} + \left( \frac{3|f'(a)|+4|f'(b)|}{4} \right)^{1/q} \right] \\ & \leq \frac{b-a}{4} \left( \frac{4}{\alpha p+1} \right)^{1/p} [|f'(a)| + |f'(b)|], \end{aligned}$$

where  $1/p + 1/q = 1$ .

It is notable that Theorem D reduces to the theorems in [14]. In 2017, Set, Sarikaya and Gözpınart [31] generalized the proceeding theorem to conformal fractional integrals. In 2020, the authors [6] extended them to more generalized fractional integrals. In 2021, the authors [39] gained similar inequalities for strongly  $(s, m)$ -convex functions. Moreover, Noor and Awan [20] proved an equality for twice differentiable functions.

**LEMMA C.** [20] Let  $f : [a, b] \rightarrow \mathbb{R}$  be a twice differentiable function and  $f'' \in L([a, b])$ . Then

$$\begin{aligned} & \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - f\left(\frac{a+b}{2}\right) \\ & = \frac{(b-a)^2}{8(\alpha+1)} \int_0^1 (1-t)^{\alpha+1} \left[ f''\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) + f''\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) \right] dt. \end{aligned}$$

Consequently, they established the following inequalities for  $s$ -convex functions.

**THEOREM E.** [20] Let  $f \in C^2([a, b])$  and  $f'' \in L([a, b])$ . Suppose that  $|f''|^q$  ( $1 \leq q < \infty$ ) is an  $s$ -convex function (in the second sense).

(i) If  $1 \leq q < \infty$ , then

$$\begin{aligned} & \left| \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - f\left(\frac{a+b}{2}\right) \right| \\ & \leq \frac{(b-a)^2}{2^{3+s/q}(\alpha+1)} \left( \frac{1}{\alpha+2} \right)^{1-1/q} \\ & \quad \times \left\{ \left( \int_0^1 (1-t)^{\alpha+1} (1+t)^s dt |f''(a)|^q + \frac{1}{s+\alpha+2} |f''(b)|^q \right)^{1/q} \right. \\ & \quad \left. + \left( \frac{1}{s+\alpha+2} |f''(a)|^q + \int_0^1 (1-t)^{\alpha+1} (1+t)^s dt |f''(b)|^q \right)^{1/q} \right\}. \end{aligned}$$

(ii) If  $1 < q < \infty$ , then

$$\begin{aligned} & \left| \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - f\left(\frac{a+b}{2}\right) \right| \\ & \leq \frac{(b-a)^2}{2^{3+s/q}(\alpha+1)} \left( \frac{1}{p(\alpha+1)+1} \right)^{1/p} \left( \frac{1}{s+1} \right)^{1/q} \\ & \quad \times \left\{ [(2^{s+1}-1) |f''(a)|^q + |f''(b)|^q]^{\frac{1}{q}} + [|f''(a)|^q + (2^{s+1}-1) |f''(b)|^q]^{1/q} \right\}, \end{aligned}$$

where  $1/p + 1/q = 1$ .

On the other hand, in 1998, Dragomir and Agarwal [7] established the following identity for the right hand side of (1.2), and then they gained error estimates related to it. Some more studies please refer to, for examples, [1, 13, 14, 22].

LEMMA D. [7] Let  $f \in C^1([a,b])$  and  $f' \in L([a,b])$ . Then

$$\frac{f(a)+f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx = \frac{b-a}{2} \int_0^1 (1-2t) f'(ta+(1-t)b) dt.$$

In 2016, Özdemir, M. Avci-Ardinç and H. Kavurmacı-Önalan [21] (Lemma 2 for  $x = (a+b)/2$ ) proved a trapezoid type equality for differentiable function via fractional integral.

LEMMA E. [21] Let  $f \in C^1([a,b])$  and  $f' \in L([a,b])$ . Then

$$\begin{aligned} & \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - \frac{f(a)+(b)}{2} \\ & = \frac{(b-a)}{4} \int_0^1 [1 - (1-t)^\alpha] \left[ f'\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) - f'\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) \right] dt. \end{aligned}$$

Thereafter, Budak [5] obtained it for generalized fractional integral in 2019 and Budak, Ertuğral and Sarikaya [6] extended it to other fractional integrals in 2020. As a consequence, the authors obtained the following results.

**THEOREM F. [5, 6]** Let  $f \in C^1([a, b])$  and  $f' \in L([a, b])$ . Suppose that  $|f'|^q$  is convex on  $[a, b]$  for  $1 \leq q < \infty$ .

(i) If  $1 \leq q < \infty$ , then

$$\begin{aligned} & \left| \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - \frac{f(a)+f(b)}{2} \right| \\ & \leq \frac{b-a}{2^{2+1/q}} \frac{\alpha}{\alpha+1} \left[ \left( \frac{\alpha+1}{2(\alpha+2)} |f'(a)|^q + \frac{3\alpha+7}{2(\alpha+2)} |f'(b)|^q \right)^{1/q} \right. \\ & \quad \left. + \left( \frac{3\alpha+7}{2(\alpha+2)} |f'(a)|^q + \frac{\alpha+1}{2(\alpha+2)} |f'(b)|^q \right)^{1/q} \right]. \end{aligned}$$

(ii) If  $1 < q < \infty$ , then

$$\begin{aligned} & \left| \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - \frac{f(a)+f(b)}{2} \right| \\ & \leq \frac{b-a}{4} \left( \frac{1}{\alpha p+1} \right)^{1/p} \left[ \left( \frac{|f'(a)|+3|f'(b)|}{4} \right)^{1/q} + \left( \frac{3|f'(a)|+4|f'(b)|}{4} \right)^{1/q} \right] \\ & \leq \frac{b-a}{4} \left( \frac{4}{\alpha p+1} \right)^{1/p} [|f'(a)|+|f'(b)|], \end{aligned}$$

where  $1/p+1/q=1$ .

In the sequel, we assume that the function  $h$  in the above definitions is always Lebesgue integrable on  $[0, 1]$ . Denote  $L(I)$  be the set of Lebesgue integrable functions on the interval  $I$  and let  $C^n(I)$  be the space of functions  $f$  with its derivatives  $f^{(k)}$  continuous on  $I$  for all  $0 \leq k \leq n$ . The aim of this paper is to extend the above results to strongly  $h$ -convex functions and obtain some error estimates related to these inequalities.

## 2. New Hermite-Hadamard inequality via Riemann-Liouville fractional integrals

In this section, we establish a similar results as Theorem B for strongly  $h$ -convex functions.

**THEOREM 1.** Let  $f \in L([a, b])$  and  $f \in SX(h, \beta, [a, b])$  with  $h(1/2) > 0$ . Then

$$\begin{aligned} & \frac{1}{2h(1/2)} \left[ f\left(\frac{a+b}{2}\right) + \frac{\beta(b-a)^2}{2(\alpha+1)(\alpha+2)} \right] \\ & \leq \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] \\ & \leq \frac{f(a)+f(b)}{2} \alpha \int_0^1 t^{\alpha-1} \left[ h\left(\frac{t}{2}\right) + h\left(1-\frac{t}{2}\right) \right] dt - \frac{\beta\alpha(\alpha+3)(b-a)^2}{4(\alpha+1)(\alpha+2)}. \end{aligned}$$

*Proof.* Since  $f$  is a strongly  $h$ -covex function with modulus  $\beta$ , for any  $t \in [0, 1]$  we have

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &= f\left[\frac{1}{2}\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) + \frac{1}{2}\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right)\right] \\ &\leq h(1/2)f\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) + h(1/2)f\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) - \frac{\beta}{4}t^2(b-a)^2, \end{aligned}$$

with means that

$$\begin{aligned} &\frac{1}{\alpha h(1/2)}f\left(\frac{a+b}{2}\right) \\ &\leq \int_0^1 (1-t)^{\alpha-1}f\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right)dt + \int_0^1 (1-t)^{\alpha-1}f\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right)dt \\ &\quad - \frac{\beta(b-a)^2}{4h(1/2)} \int_0^1 (1-t)^{\alpha-1}t^2 dt \\ &= \frac{2^\alpha}{(b-a)^\alpha} \int_{(a+b)/2}^b (b-u)^{\alpha-1}f(u)du + \frac{2^\alpha}{(b-a)^\alpha} \int_a^{(a+b)/2} (u-a)^{\alpha-1}f(u)du \quad (2.1) \end{aligned}$$

$$\begin{aligned} &\quad - \frac{\beta(b-a)^2}{4h(1/2)\alpha(\alpha+1)(\alpha+2)} \\ &= \frac{2^\alpha\Gamma(\alpha)}{(b-a)^\alpha} \left[ J_{(\frac{a+b}{2})^+}^\alpha f(b) + J_{(\frac{a+b}{2})^-}^\alpha f(a) \right] - \frac{\beta(b-a)^2}{4h(1/2)\alpha(\alpha+1)(\alpha+2)}. \quad (2.2) \end{aligned}$$

Therefore we finish the first inequality of the theorem.

On the other hand, it is easy to see that

$$\begin{aligned} f\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) &\leq h\left(\frac{1-t}{2}\right)f(a) + h\left(\frac{1+t}{2}\right)f(b) - \frac{\beta(1-t)(1+t)}{4}(b-a)^2, \\ f\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) &\leq h\left(\frac{1+t}{2}\right)f(a) + h\left(\frac{1-t}{2}\right)f(b) - \frac{\beta(1-t)(1+t)}{4}(b-a)^2, \end{aligned}$$

which, combining with (2.1) and (2.2), imply that

$$\begin{aligned} &\frac{2^\alpha\Gamma(\alpha)}{(b-a)^\alpha} \left[ J_{(\frac{a+b}{2})^-}^\alpha f(a) + J_{(\frac{a+b}{2})^+}^\alpha f(b) \right] \\ &= \int_0^1 (1-t)^{\alpha-1}f\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right)dt + \int_0^1 (1-t)^{\alpha-1}f\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right)dt \\ &\leq [f(a) + f(b)] \int_0^1 (1-t)^{\alpha-1} \left[ h\left(\frac{1-t}{2}\right) + h\left(\frac{1+t}{2}\right) \right] dt - \frac{\beta(b-a)^2}{2} \frac{\alpha+3}{(\alpha+1)(\alpha+2)}. \end{aligned}$$

Thus we finish the proof of Theorem 1.  $\square$

If taking  $h(t) = t$  and  $h(t) = t^s$ , then Theorem 1 reduces to Corollary 3 and Corollary 4 in [39], respectively. And, it is not difficult to see that the theorem is Theorem A for  $\alpha = 1$ . Letting  $\beta \rightarrow 0$  in Theorem 1, we have the following result.

COROLLARY 1. Let  $f : [a, b] \rightarrow \mathbb{R}$  be an  $h$ -covex function with  $h(1/2) > 0$  and  $f \in L([a, b])$ . Then

$$\begin{aligned} \frac{1}{2h(1/2)} f\left(\frac{a+b}{2}\right) &\leq \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] \\ &\leq \frac{f(a)+f(b)}{2} \alpha \int_0^1 t^{\alpha-1} \left[ h\left(\frac{t}{2}\right) + h\left(1-\frac{t}{2}\right) \right] dt. \end{aligned}$$

Especially, if  $h(t) = t$ , Corollary 1 becomes Theorem B.

### 3. Midpoint type inequalities for $n$ times differentiable functions

In this section, we will extend the midpoint type inequalities in Theorem D and Theorem E to the case of strongly  $h$ -convex functions with  $n$  order derivatives. For the sake of convenience, if  $f : [a, b] \rightarrow \mathbb{R}$  is an  $n$ -times differentiable function, we denote

$$\begin{aligned} \mathfrak{L}\left(f, \frac{a+b}{2}\right) &= \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - f\left(\frac{a+b}{2}\right) \\ &\quad - \sum_{j=1}^{n-1} \frac{[1+(-1)^j](b-a)^j}{2^{j+1} \prod_{k=1}^j (\alpha+k)} f^{(j)}\left(\frac{a+b}{2}\right). \end{aligned} \quad (3.1)$$

It is easy to see that if  $n = 1$  or  $2$ ,  $\mathfrak{L}(f, \frac{a+b}{2})$  have the same concise form:

$$\mathfrak{L}\left(f, \frac{a+b}{2}\right) = \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - f\left(\frac{a+b}{2}\right). \quad (3.2)$$

Now we introduce the following key lemma.

LEMMA 1. Let  $f \in C^n([a, b])$  and  $f^{(n)} \in L([a, b])$ ,  $n \in \mathbb{Z}^+$ . Then

$$\begin{aligned} \mathfrak{L}\left(f, \frac{a+b}{2}\right) &= \frac{(b-a)^n}{2^{n+1} \prod_{k=1}^{n-1} (\alpha+k)} \left[ (-1)^n \int_0^1 (1-t)^{\alpha+n-1} f^{(n)}\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) dt \right. \\ &\quad \left. + \int_0^1 (1-t)^{\alpha+n-1} f^{(n)}\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) dt \right], \end{aligned}$$

here we denote  $\prod_{k=1}^0 (\alpha+k) \equiv 1$ .

It is not difficult to check that Lemma 1 reduces to Lemma B and Lemma C for  $n = 1$  and  $n = 2$ , respectively.

*Proof.* Without loss of generality, we may assume  $n \geq 2$ . Integration by parts  $n$  times show that

$$\begin{aligned} &\int_0^1 (1-t)^{\alpha+n-1} f^{(n)}\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) dt \\ &= \frac{2}{b-a} f^{(n-1)}\left(\frac{a+b}{2}\right) - \frac{2(\alpha+n-1)}{b-a} \int_0^1 (1-t)^{\alpha+n-2} f^{(n-1)}\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) dt \end{aligned}$$

$$\begin{aligned}
&= \frac{2}{b-a} f^{(n-1)}\left(\frac{a+b}{2}\right) - \frac{2^2(\alpha+n-1)}{(b-a)^2} f^{(n-2)}\left(\frac{a+b}{2}\right) \\
&\quad + \frac{2^2(\alpha+n-1)(\alpha+n-2)}{(b-a)^2} \int_0^1 (1-t)^{\alpha+n-3} f^{(n-2)}\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) dt \\
&= \dots \\
&= \frac{2}{b-a} f^{(n-1)}\left(\frac{a+b}{2}\right) - \frac{2^2(\alpha+n-1)}{(b-a)^2} f^{(n-2)}\left(\frac{a+b}{2}\right) \\
&\quad + \frac{2^3(\alpha+n-1)(\alpha+n-2)}{(b-a)^3} f^{(n-3)}\left(\frac{a+b}{2}\right) \\
&\quad + \dots + \frac{(-1)^{n-1} 2^n (\alpha+n-1)(\alpha+n-2)\dots(\alpha+1)}{(b-a)^n} f\left(\frac{a+b}{2}\right) \\
&\quad + \frac{(-1)^n 2^n (\alpha+n-1)(\alpha+n-2)\dots(\alpha+1)\alpha}{(b-a)^{\alpha+n}} \int_0^1 (1-t)^{\alpha-1} f\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) dt \\
&= \sum_{j=0}^{n-1} \frac{(-1)^{n-j-1} 2^{n-j} \prod_{k=j+1}^{n-1} (\alpha+k)}{(b-a)^{n-j}} f^{(j)}\left(\frac{a+b}{2}\right) \\
&\quad + \frac{(-1)^n 2^{\alpha+n} \prod_{k=0}^{n-1} (\alpha+k) \Gamma(\alpha)}{(b-a)^{\alpha+n}} J_{(\frac{a+b}{2})^-}^\alpha f(a).
\end{aligned}$$

Multiplying both sides of the proceeding equality by  $\frac{(-1)^n (b-a)^n}{2^{n+1} \prod_{k=1}^{n-1} (\alpha+k)}$ , we obtain

$$\begin{aligned}
&\frac{(-1)^n (b-a)^n}{2^{n+1} \prod_{k=1}^{n-1} (\alpha+k)} \int_0^1 (1-t)^{\alpha+n-1} f^{(n)}\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) dt \tag{3.3} \\
&= \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha} J_{(\frac{a+b}{2})^-}^\alpha f(a) - \frac{1}{2} f\left(\frac{a+b}{2}\right) - \sum_{j=1}^{n-1} \frac{(-1)^j (b-a)^j}{2^{j+1} \prod_{k=1}^j (\alpha+k)} f^{(j)}\left(\frac{a+b}{2}\right).
\end{aligned}$$

Similarly, using integration by parts  $n$  times again,

$$\begin{aligned}
&\int_0^1 (1-t)^{\alpha+n-1} f^{(n)}\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) dt \\
&= - \sum_{j=0}^{n-1} \frac{2^{n-j} \prod_{k=j+1}^{n-1} (\alpha+k)}{(b-a)^{n-j}} f^{(j)}\left(\frac{a+b}{2}\right) + \frac{2^{\alpha+n} \prod_{k=0}^{n-1} (\alpha+k) \Gamma(\alpha)}{(b-a)^{\alpha+n}} J_{(\frac{a+b}{2})^+}^\alpha f(b),
\end{aligned}$$

which means that

$$\begin{aligned}
&\frac{(b-a)^n}{2^{n+1} \prod_{k=1}^{n-1} (\alpha+k)} \int_0^1 (1-t)^{\alpha+n-1} f^{(n)}\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) dt \tag{3.4} \\
&= \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha} J_{(\frac{a+b}{2})^+}^\alpha f(b) - \frac{1}{2} f\left(\frac{a+b}{2}\right) - \sum_{j=1}^{n-1} \frac{(b-a)^j}{2^{j+1} \prod_{k=1}^j (\alpha+k)} f^{(j)}\left(\frac{a+b}{2}\right).
\end{aligned}$$

Therefore we complete the proof of the lemma by (3.3) and (3.4).  $\square$

Using Lemma 1, we obtain the following fractional integral inequalities. For simplicity, we first denote

$$\mathcal{A} = \int_0^{1/2} t^{\alpha+n-1} h(t) dt, \quad \mathcal{B} = \int_{1/2}^1 (1-t)^{\alpha+n-1} h(t) dt.$$

**THEOREM 2.** Let  $f \in C^n([a, b])$  and  $f^{(n)} \in L([a, b]), n \in \mathbb{Z}^+$ . Suppose that  $|f^{(n)}|^q \in SX(h, \beta, [a, b]), 1 \leq q < \infty$ .

(i) If  $q = 1$ , then

$$\begin{aligned} & \left| \mathfrak{L}\left(f, \frac{a+b}{2}\right) \right| \\ & \leq \frac{2^{\alpha-1}(b-a)^n}{\prod_{k=1}^{n-1}(\alpha+k)} \left\{ (\mathcal{A} + \mathcal{B}) \left( |f^{(n)}(a)| + |f^{(n)}(b)| \right) - \frac{\beta(\alpha+n+3)(b-a)^2}{2^{n+\alpha+1}(\alpha+n+1)(\alpha+n+2)} \right\}. \end{aligned}$$

(ii) If  $1 < q < \infty$ , then

$$\begin{aligned} & \left| \mathfrak{L}\left(f, \frac{a+b}{2}\right) \right| \\ & \leq \frac{2^{(\alpha+n)/q-n-1}(\alpha+n)^{1/q}(b-a)^n}{\prod_{k=1}^n(\alpha+k)} \\ & \quad \times \left\{ \left[ \mathcal{B} |f^{(n)}(a)|^q + \mathcal{A} |f^{(n)}(b)|^q - \frac{\beta(\alpha+n+3)(b-a)^2}{2^{\alpha+n+2}(\alpha+n+1)(\alpha+n+2)} \right]^{1/q} \right. \\ & \quad \left. + \left[ \mathcal{B} |f^{(n)}(b)|^q + \mathcal{A} |f^{(n)}(a)|^q - \frac{\beta(\alpha+n+3)(b-a)^2}{2^{\alpha+n+2}(\alpha+n+1)(\alpha+n+2)} \right]^{1/q} \right\} \\ & \leq \frac{2^{(\alpha+n)/q-n-1}(\alpha+n)^{1/q}(b-a)^n}{\prod_{k=1}^n(\alpha+k)} \\ & \quad \times \left\{ \left( \mathcal{A}^{1/q} + \mathcal{B}^{1/q} \right) \left( |f^{(n)}(a)| + |f^{(n)}(b)| \right) + 2 \left[ \frac{\beta(\alpha+n+3)(b-a)^2}{2^{\alpha+n+2}(\alpha+n+1)(\alpha+n+2)} \right]^{1/q} \right\}. \end{aligned}$$

(iii) If  $1 < q < \infty$ , then

$$\begin{aligned} & \left| \mathfrak{L}\left(f, \frac{a+b}{2}\right) \right| \\ & \leq \frac{(b-a)^n}{2^{n+1/p} \prod_{k=1}^{n-1}(\alpha+k)} \left( \frac{1}{p(\alpha+n-1)+1} \right)^{1/p} \\ & \quad \times \left\{ \left( \int_{1/2}^1 h(t) dt |f^{(n)}(a)|^q + \int_0^{1/2} h(t) dt |f^{(n)}(b)|^q - \frac{\beta(b-a)^2}{12} \right)^{1/q} \right. \\ & \quad \left. + \left( \int_{1/2}^1 h(t) dt |f^{(n)}(b)|^q + \int_0^{1/2} h(t) dt |f^{(n)}(a)|^q - \frac{\beta(b-a)^2}{12} \right)^{1/q} \right\} \end{aligned}$$

$$\leq \frac{(b-a)^n}{2^{n+1/p} \prod_{k=1}^{n-1} (\alpha+k)} \left( \frac{1}{p(\alpha+n-1)+1} \right)^{1/p} \\ \times \left\{ \left[ \left( \int_0^{1/2} h(t) dt \right)^{1/q} + \left( \int_{1/2}^1 h(t) dt \right)^{1/q} \right] \left( |f^{(n)}(a)| + |f^{(n)}(b)| \right) \right. \\ \left. + 2 \left[ \frac{\beta(b-a)^2}{12} \right]^{1/q} \right\},$$

where  $\frac{1}{p} + \frac{1}{q} = 1$ .

*Proof.* (1) If  $q = 1$ , then it follows from the fact of  $|f| \in SX(h, \beta, [a, b])$  that

$$\left| \int_0^1 (1-t)^{\alpha+n-1} f^{(n)} \left( \frac{1+t}{2} a + \frac{1-t}{2} b \right) dt \right| \\ \leq \int_0^1 (1-t)^{\alpha+n-1} \left[ h \left( \frac{1+t}{2} \right) |f^{(n)}(a)| + h \left( \frac{1-t}{2} \right) |f^{(n)}(b)| \right. \\ \left. - \beta \frac{(1-t)(1+t)}{4} (b-a)^2 \right] dt \\ = 2^{\alpha+n} \int_{1/2}^1 (1-t)^{\alpha+n-1} h(t) dt |f^{(n)}(a)| + 2^{\alpha+n} \int_0^{1/2} t^{\alpha+n-1} h(t) dt |f^{(n)}(b)| \\ - \frac{\beta(\alpha+n+3)(b-a)^2}{4(\alpha+n+1)(\alpha+n+2)} \\ = 2^{\alpha+n} \mathcal{B} |f^{(n)}(a)| + 2^{\alpha+n} \mathcal{A} |f^{(n)}(b)| - \frac{\beta(\alpha+n+3)(b-a)^2}{4(\alpha+n+1)(\alpha+n+2)}.$$

Similarly,

$$\left| \int_0^1 (1-t)^{\alpha+n-1} f^{(n)} \left( \frac{1-t}{2} a + \frac{1+t}{2} b \right) dt \right| \\ \leq 2^{\alpha+n} \mathcal{A} |f^{(n)}(a)| + 2^{\alpha+n} \mathcal{B} |f^{(n)}(b)| - \frac{\beta(\alpha+n+3)(b-a)^2}{4(\alpha+n+1)(\alpha+n+2)}.$$

Then we complete the proof of (i) by the proceeding two inequalities and Lemma 1.

(2) If  $1 < q < \infty$ , then power-mean inequality and the fact of  $|f|^q \in SX(h, \beta, [a, b])$  show that

$$\left| \int_0^1 (1-t)^{\alpha+n-1} f^{(n)} \left( \frac{1+t}{2} a + \frac{1-t}{2} b \right) dt \right| \\ \leq \left( \int_0^1 (1-t)^{\alpha+n-1} dt \right)^{1-1/q} \left( \int_0^1 (1-t)^{\alpha+n-1} \left| f^{(n)} \left( \frac{1+t}{2} a + \frac{1-t}{2} b \right) \right|^q dt \right)^{1/q}$$

$$\begin{aligned}
&\leq \left( \frac{1}{\alpha+n} \right)^{1-1/q} \left[ 2^{\alpha+n} \int_{1/2}^1 (1-t)^{\alpha+n-1} h(t) dt \left| f^{(n)}(a) \right|^q \right. \\
&\quad \left. + 2^{\alpha+n} \int_0^{1/2} t^{\alpha+n-1} h(t) dt \left| f^{(n)}(b) \right|^q - \frac{\beta(\alpha+n+3)(b-a)^2}{4(\alpha+n+1)(\alpha+n+2)} \right]^{1/q} \\
&= \frac{2^{(\alpha+n)/q}}{(\alpha+n)^{1-1/q}} \left[ \mathcal{B} \left| f^{(n)}(a) \right|^q + \mathcal{A} \left| f^{(n)}(b) \right|^q - \frac{\beta(\alpha+n+3)(b-a)^2}{2^{\alpha+n+2}(\alpha+n+1)(\alpha+n+2)} \right]^{1/q},
\end{aligned}$$

and

$$\begin{aligned}
&\left| \int_0^1 (1-t)^{\alpha+n-1} f^{(n)} \left( \frac{1-t}{2}a + \frac{1+t}{2}b \right) dt \right| \\
&\leq \frac{2^{(\alpha+n)/q}}{(\alpha+n)^{1-1/q}} \left[ \mathcal{B} \left| f^{(n)}(b) \right|^q + \mathcal{A} \left| f^{(n)}(a) \right|^q - \frac{\beta(\alpha+n+3)(b-a)^2}{2^{\alpha+n+2}(\alpha+n+1)(\alpha+n+2)} \right]^{1/q},
\end{aligned}$$

which finish the proof of the first inequality in (ii) by Lemma 1 again.

For the proof of the second inequality, let

$$\begin{aligned}
a_1 &= \int_{1/2}^1 (1-t)^{\alpha+n-1} h(t) dt \left| f^{(n)}(a) \right|^q, \quad b_1 = 2 \int_0^{1/2} t^{\alpha+n-1} h(t) dt \left| f^{(n)}(b) \right|^q, \\
a_2 &= \int_{1/2}^1 (1-t)^{\alpha+n-1} h(t) dt \left| f^{(n)}(b) \right|^q, \quad b_2 = \int_0^{1/2} t^{\alpha+n-1} h(t) dt \left| f^{(n)}(a) \right|^q, \\
c_1 = c_2 &= -\frac{\beta(\alpha+n+3)}{2^{\alpha+n+2}(\alpha+n+1)(\alpha+n+2)} (b-a)^2.
\end{aligned}$$

According to the fact that

$$\sum_{k=1}^m (|a_k| + |b_k| + |c_k|)^s \leq \sum_{k=1}^m |a_k|^s + \sum_{k=1}^m |b_k|^s + \sum_{k=1}^m |c_k|^s, \quad 0 \leq s < 1,$$

then the desired result can be obtained easily.

(3) If  $1 < q < \infty$ , then the Hölder inequality and the fact of  $|f|^q \in SX(h, \beta, [a, b])$  tell us that

$$\begin{aligned}
&\left| \int_0^1 (1-t)^{\alpha+n-1} f^{(n)} \left( \frac{1+t}{2}a + \frac{1-t}{2}b \right) dt \right| \\
&\leq \left( \int_0^1 (1-t)^{p(\alpha+n-1)} dt \right)^{1/p} \left( \int_0^1 \left| f^{(n)} \left( \frac{1+t}{2}a + \frac{1-t}{2}b \right) \right|^q dt \right)^{q/p} \\
&\leq \left( \frac{1}{p(\alpha+n-1)+1} \right)^{1/p} \left\{ \int_0^1 \left[ h \left( \frac{1+t}{2} \right) \left| f^{(n)}(a) \right|^q + h \left( \frac{1-t}{2} \right) \left| f^{(n)}(b) \right|^q \right. \right. \\
&\quad \left. \left. - \beta \frac{(1-t)(1+t)}{4} (b-a)^2 \right] dt \right\}^{1/q}
\end{aligned}$$

$$= \left( \frac{1}{p(\alpha+n-1)+1} \right)^{1/p} \left( 2 \int_{1/2}^1 h(t) dt \left| f^{(n)}(a) \right|^q + 2 \int_0^{1/2} h(t) dt \left| f^{(n)}(b) \right|^q - \frac{\beta(b-a)^2}{6} \right)^{1/q}.$$

By the same way, we have

$$\begin{aligned} & \left| \int_0^1 (1-t)^{\alpha+n-1} f^{(n)} \left( \frac{1-t}{2} a + \frac{1+t}{2} b \right) dt \right| \\ & \leq \left( \frac{1}{p(\alpha+n-1)+1} \right)^{1/p} \left( 2 \int_{1/2}^1 h(t) dt \left| f^{(n)}(b) \right|^q + 2 \int_0^{1/2} h(t) dt \left| f^{(n)}(a) \right|^q - \frac{\beta(b-a)^2}{6} \right)^{1/q}. \end{aligned}$$

Then we complete the proof of the first inequality in (ii) by the above two inequalities and Lemma 1.

The second inequality is proved by a similar way as (2), we leave the details to readers.  $\square$

Letting  $\beta \rightarrow 0$ . We have the following results.

**COROLLARY 2.** Let  $f \in C^n([a,b])$  and  $f^{(n)} \in L([a,b])$ ,  $n \in \mathbb{Z}^+$ . Suppose that  $|f^{(n)}|^q$  is an  $h$ -convex function with  $1 \leq q < \infty$ .

(i) If  $1 \leq q < \infty$ , then

$$\begin{aligned} & \left| \mathfrak{L} \left( f, \frac{a+b}{2} \right) \right| \\ & \leq \frac{2^{(\alpha+n)/q-n-1} (\alpha+n)^{1/q} (b-a)^n}{\prod_{k=1}^n (\alpha+k)} \\ & \quad \times \left\{ [\mathcal{B} |f^{(n)}(a)|^q + \mathcal{A} |f^{(n)}(b)|^q]^{1/q} + [\mathcal{B} |f^{(n)}(b)|^q + \mathcal{A} |f^{(n)}(a)|^q]^{1/q} \right\} \\ & \leq \frac{2^{(\alpha+n)/q-n-1} (\alpha+n)^{1/q} (b-a)^n}{\prod_{k=1}^n (\alpha+k)} \left( \mathcal{A}^{1/q} + \mathcal{B}^{1/q} \right) \left( |f^{(n)}(a)| + |f^{(n)}(b)| \right). \end{aligned}$$

(ii) If  $1 < q < \infty$ , then

$$\begin{aligned} & \left| \mathfrak{L} \left( f, \frac{a+b}{2} \right) \right| \leq \frac{(b-a)^n}{2^{n+1/p} \prod_{k=1}^{n-1} (\alpha+k)} \left( \frac{1}{p(\alpha+n-1)+1} \right)^{1/p} \\ & \quad \times \left\{ \left( \int_{1/2}^1 h(t) dt \left| f^{(n)}(a) \right|^q + \int_0^{1/2} h(t) dt \left| f^{(n)}(b) \right|^q \right)^{1/q} \right. \\ & \quad \left. + \left( \int_{1/2}^1 h(t) dt \left| f^{(n)}(b) \right|^q + \int_0^{1/2} h(t) dt \left| f^{(n)}(a) \right|^q \right)^{1/q} \right\} \end{aligned}$$

$$\leq \frac{(b-a)^n}{2^{n+1/p} \prod_{k=1}^{n-1} (\alpha+k)} \left( \frac{1}{p(\alpha+n-1)+1} \right)^{1/p} \\ \times \left[ \left( \int_0^{1/2} h(t) dt \right)^{1/q} + \left( \int_{1/2}^1 h(t) dt \right)^{1/q} \right] \left( |f^{(n)}(a)| + |f^{(n)}(b)| \right),$$

where  $\frac{1}{p} + \frac{1}{q} = 1$ .

REMARK. By changing of variable, it is not difficult to check that Corollary 2 extends Theorem D and Theorem E for  $n = 1$ ,  $h(t) = t$  and  $n = 2$ ,  $h(t) = t^s$ , respectively.

#### 4. Trapezoid type inequalities for $n$ times differentiable functions

In this section, we will prove some similar results as Theorem F for strongly  $h$ -convex functions with  $n$  order derivatives. For simplicity, if  $f \in C^n([a, b])$ , we denote

$$\mathfrak{R}\left(f, \frac{a+b}{2}\right) = \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - \frac{f(a) + f(b)}{2} \\ + \sum_{j=1}^{n-1} \frac{[1 + (-1)^j]}{2^{j+1}} \frac{(b-a)^j}{j! \prod_{k=1}^j (\alpha+k)} \prod_{k=1}^j (\alpha+k) - j! f^{(j)}\left(\frac{a+b}{2}\right). \quad (4.1)$$

It is easy to check that if  $n = 1$  or  $2$ ,  $\mathfrak{R}\left(f, \frac{a+b}{2}\right)$  has the simplified form:

$$\mathfrak{R}\left(f, \frac{a+b}{2}\right) = \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] - \frac{f(a) + f(b)}{2}. \quad (4.2)$$

Now we introduce the following key lemma.

LEMMA 2. Let  $f \in C^n([a, b])$  and  $f^{(n)} \in L([a, b])$ ,  $n \in \mathbb{Z}^+$ . Then

$$\mathfrak{R}\left(f, \frac{a+b}{2}\right) = -\frac{(b-a)^n}{2^{n+1} \prod_{k=1}^{n-1} (\alpha+k)} \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] \\ \times \left[ (-1)^n f^{(n)}\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) + f^{(n)}\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) \right] dt.$$

It is not difficult to check that Lemma 2 reduces to Lemma E by (4.2) for  $n = 1$ .

*Proof.* Without loss of generality, we assume that  $n \geq 2$ . Integrating by parts  $n$  times show that

$$\int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] f^{(n)}\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) dt \\ = \frac{2}{b-a} \left( \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} - 1 \right) f^{(n-1)}\left(\frac{a+b}{2}\right) \\ - \frac{2(\alpha+n-1)}{b-a} \int_0^1 \left[ \frac{\prod_{k=1}^{n-2} (\alpha+k)}{(n-2)!} (1-t)^{n-2} - (1-t)^{\alpha+n-2} \right] f^{(n-1)}\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) dt$$

$$\begin{aligned}
&= \frac{2}{b-a} \left( \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} - 1 \right) f^{(n-1)} \left( \frac{a+b}{2} \right) \\
&\quad - \frac{2^2(\alpha+n-1)}{(b-a)^2} \left( \frac{\prod_{k=1}^{n-2} (\alpha+k)}{(n-2)!} - 1 \right) f^{(n-2)} \left( \frac{a+b}{2} \right) \\
&\quad + \frac{2^2(\alpha+n-1)(\alpha+n-2)}{(b-a)^2} \\
&\quad \times \int_0^1 \left[ \frac{\prod_{k=1}^{n-3} (\alpha+k)}{(n-3)!} (1-t)^{n-3} - (1-t)^{\alpha+n-3} \right] f^{(n-2)} \left( \frac{1+t}{2}a + \frac{1-t}{2}b \right) dt \\
&= \dots \\
&= \sum_{j=1}^{n-1} \frac{(-1)^{n-j+1} 2^{n-j} \prod_{i=j+1}^{n-1} (\alpha+i)}{(b-a)^{n-j}} \left( \frac{\prod_{k=1}^j (\alpha+k)}{j!} - 1 \right) f^{(j)} \left( \frac{a+b}{2} \right) \\
&\quad + \frac{(-1)^n 2^{n-1} \prod_{k=1}^{n-1} (\alpha+k)}{(b-a)^{n-1}} \int_0^1 [1 - (1-t)^\alpha] f' \left( \frac{1+t}{2}a + \frac{1-t}{2}b \right) dt \\
&= \sum_{j=1}^{n-1} \frac{(-1)^{n-j+1} 2^{n-j} \prod_{i=j+1}^{n-1} (\alpha+i)}{(b-a)^{n-j}} \left( \frac{\prod_{k=1}^j (\alpha+k)}{j!} - 1 \right) f^{(j)} \left( \frac{a+b}{2} \right) \\
&\quad + \frac{(-1)^n 2^n \prod_{k=1}^{n-1} (\alpha+k)}{(b-a)^n} f(a) + \frac{(-1)^{n+1} 2^n \prod_{k=0}^{n-1} (\alpha+k)}{(b-a)^n} \\
&\quad \times \int_0^1 (1-t)^{\alpha-1} f \left( \frac{1+t}{2}a + \frac{1-t}{2}b \right) dt \\
&= \sum_{j=1}^{n-1} \frac{(-1)^{n-j+1} 2^{n-j} \prod_{i=j+1}^{n-1} (\alpha+i)}{(b-a)^{n-j}} \left( \frac{\prod_{k=1}^j (\alpha+k)}{j!} - 1 \right) f^{(j)} \left( \frac{a+b}{2} \right) \\
&\quad + \frac{(-1)^n 2^n \prod_{k=1}^{n-1} (\alpha+k)}{(b-a)^n} f(a) + \frac{(-1)^{n+1} 2^{n+\alpha} \Gamma(\alpha) \prod_{k=0}^{n-1} (\alpha+k)}{(b-a)^{n+\alpha}} J_{(\frac{a+b}{2})^-}^\alpha f(a),
\end{aligned}$$

which means that

$$\begin{aligned}
&\frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha} J_{(\frac{a+b}{2})^-}^\alpha f(a) - \frac{f(a)}{2} \\
&\quad + \sum_{j=1}^{n-1} \frac{(-1)^j (b-a)^j \prod_{k=1}^j (\alpha+k) - j!}{2^{j+1} j! \prod_{k=1}^j (\alpha+k)} f^{(j)} \left( \frac{a+b}{2} \right) \\
&= \frac{(-1)^{n+1} (b-a)^n}{2^{n+1} \prod_{k=1}^{n-1} (\alpha+k)} \\
&\quad \times \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] f^{(n)} \left( \frac{1+t}{2}a + \frac{1-t}{2}b \right) dt. \tag{4.3}
\end{aligned}$$

Similarly, integrating by parts  $n$  times again tell us that

$$\begin{aligned}
& \int_0^1 \left[ \frac{\prod_{k=1}^{n-1}(\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] f^{(n)} \left( \frac{1-t}{2}a + \frac{1+t}{2}b \right) dt \\
&= -\frac{2}{b-a} \left( \frac{\prod_{k=1}^{n-1}(\alpha+k)}{(n-1)!} - 1 \right) f^{(n-1)} \left( \frac{a+b}{2} \right) + \frac{2(\alpha+n-1)}{b-a} \\
&\quad \times \int_0^1 \left[ \frac{\prod_{k=1}^{n-2}(\alpha+k)}{(n-2)!} (1-t)^{n-2} - (1-t)^{\alpha+n-2} \right] f^{(n-1)} \left( \frac{1-t}{2}a + \frac{1+t}{2}b \right) dt \\
&= \dots \\
&= -\sum_{j=1}^{n-1} \frac{2^{n-j} \prod_{i=j+1}^{n-1}(\alpha+i)}{(b-a)^{n-j}} \left( \frac{\prod_{k=1}^j(\alpha+k)}{j!} - 1 \right) f^{(j)} \left( \frac{a+b}{2} \right) \\
&\quad + \frac{2^{n-1} \prod_{k=1}^{n-1}(\alpha+k)}{(b-a)^{n-1}} \int_0^1 [1 - (1-t)^\alpha] f' \left( \frac{1-t}{2}a + \frac{1+t}{2}b \right) dt \\
&= -\sum_{j=1}^{n-1} \frac{2^{n-j} \prod_{i=j+1}^{n-1}(\alpha+i)}{(b-a)^{n-j}} \left( \frac{\prod_{k=1}^j(\alpha+k)}{j!} - 1 \right) f^{(j)} \left( \frac{a+b}{2} \right) \\
&\quad + \frac{2^n \prod_{k=1}^{n-1}(\alpha+k)}{(b-a)^n} f(b) - \frac{2^{n+\alpha} \Gamma(\alpha) \prod_{k=0}^{n-1}(\alpha+k)}{(b-a)^{n+\alpha}} J_{(\frac{a+b}{2})^+}^\alpha f(b),
\end{aligned}$$

which implies that

$$\begin{aligned}
& \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha} J_{(\frac{a+b}{2})^-}^\alpha f(a) - \frac{f(b)}{2} \\
&+ \sum_{j=1}^{n-1} \frac{(b-a)^j}{2^{j+1}} \frac{\prod_{k=1}^j(\alpha+k) - j!}{j! \prod_{k=1}^j(\alpha+k)} f^{(j)} \left( \frac{a+b}{2} \right) \\
&= -\frac{(b-a)^n}{2^{n+1} \prod_{k=1}^{n-1}(\alpha+k)} \\
&\quad \times \int_0^1 \left[ \frac{\prod_{k=1}^{n-1}(\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] f^{(n)} \left( \frac{1-t}{2}a + \frac{1+t}{2}b \right) dt.
\end{aligned} \tag{4.4}$$

Then we complete the proof by (4.3) and (4.4).  $\square$

Using Lemma 2, we obtain the following error estimates. For convenience, we first set

$$\mathcal{C} = \int_0^{1/2} \left( \frac{\prod_{k=1}^{n-1}(\alpha+k)}{(n-1)!} t^{n-1} - 2^\alpha t^{\alpha+n-1} \right) h(t) dt,$$

$$\mathcal{D} = \int_{1/2}^1 \left( \frac{\prod_{k=1}^{n-1}(\alpha+k)}{(n-1)!} (1-t)^{n-1} - 2^\alpha (1-t)^{\alpha+n-1} \right) h(t) dt.$$

**THEOREM 3.** Let  $f \in C^n([a, b])$  and  $f^{(n)} \in L([a, b]), n \in \mathbb{Z}^+$ . Suppose that  $|f^{(n)}|^q \in SX(h, \beta, [a, b]), 1 \leq q < \infty$ .

(i) If  $q = 1$ , then

$$\left| \Re \left( f, \frac{a+b}{2} \right) \right| \leq \frac{(b-a)^n}{2 \prod_{k=1}^{n-1} (\alpha+k)} \left[ (\mathcal{C} + \mathcal{D}) \left( |f^{(n)}(a)| + |f^{(n)}(b)| \right) - \beta \frac{n(n+3) \prod_{k=1}^{n+2} (\alpha+k) - (n+2)!(\alpha+n+3)}{2^{n+1}(n+2)!(\alpha+n+1)(\alpha+n+2)} (b-a)^2 \right].$$

(ii) If  $1 < q < \infty$ , then

$$\begin{aligned} & \left| \Re \left( f, \frac{a+b}{2} \right) \right| \\ & \leq \frac{(b-a)^n}{2^{n(1-1/q)+1} \prod_{k=1}^{n-1} (\alpha+k)} \left( \frac{\prod_{k=1}^{n-1} (\alpha+k)}{n!} - \frac{1}{\alpha+n} \right)^{1-1/q} \\ & \quad \times \left\{ \left[ \mathcal{C} |f^{(n)}(a)|^q + \mathcal{D} |f^{(n)}(b)|^q \right. \right. \\ & \quad \left. \left. - \beta \frac{n(n+3) \prod_{k=1}^{n+2} (\alpha+k) - (n+2)!(\alpha+n+3)}{2^{n+2}(n+2)!(\alpha+n+1)(\alpha+n+2)} (b-a)^2 \right]^{1/q} \right. \\ & \quad \left. + \left[ \mathcal{C} |f^{(n)}(b)|^q + \mathcal{D} |f^{(n)}(a)|^q \right. \right. \\ & \quad \left. \left. - \beta \frac{n(n+3) \prod_{k=1}^{n+2} (\alpha+k) - (n+2)!(\alpha+n+3)}{2^{n+2}(n+2)!(\alpha+n+1)(\alpha+n+2)} (b-a)^2 \right]^{1/q} \right\} \\ & \leq \frac{(b-a)^n}{2^{n(1-1/q)+1} \prod_{k=1}^{n-1} (\alpha+k)} \left( \frac{\prod_{k=1}^{n-1} (\alpha+k)}{n!} - \frac{1}{\alpha+n} \right)^{1-1/q} \\ & \quad \times \left\{ \left( \mathcal{C}^{1/q} + \mathcal{D}^{1/q} \right) \left( |f^{(n)}(a)| + |f^{(n)}(b)| \right) \right. \\ & \quad \left. + 2 \left[ \beta \frac{n(n+3) \prod_{k=1}^{n+2} (\alpha+k) - (n+2)!(\alpha+n+3)}{2^{n+2}(n+2)!(\alpha+n+1)(\alpha+n+2)} (b-a)^2 \right]^{1/q} \right\}. \end{aligned}$$

(iii) If  $1 < q < \infty$ , then

$$\begin{aligned} & \left| \Re \left( f, \frac{a+b}{2} \right) \right| \\ & \leq \frac{(b-a)^n}{2^{n+1-1/q} \prod_{k=1}^{n-1} (\alpha+k)} \left( \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} t^{n-1} - t^{\alpha+n-1} \right]^p dt \right)^{1/p} \end{aligned}$$

$$\begin{aligned}
& \times \left\{ \left[ \int_{1/2}^1 h(t) dt \left| f^{(n)}(b) \right|^q + \int_0^{1/2} h(t) dt \left| f^{(n)}(a) \right|^q - \frac{\beta(b-a)^2}{12} \right]^{1/q} \right. \\
& \quad \left. + \left[ \int_{1/2}^1 h(t) dt \left| f^{(n)}(a) \right|^q + \int_0^{1/2} h(t) dt \left| f^{(n)}(b) \right|^q - \frac{\beta(b-a)^2}{12} \right]^{1/q} \right\} \\
& \leqslant \frac{(b-a)^n}{2^{n+1-1/q} \prod_{k=1}^{n-1} (\alpha+k)} \left( \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} t^{n-1} - t^{\alpha+n-1} \right]^p dt \right)^{1/p} \\
& \quad \times \left\{ \left[ \left( \int_0^{1/2} h(t) dt \right)^{1/q} + \left( \int_{1/2}^1 h(t) dt \right)^{1/q} \right] \left( \left| f^{(n)}(a) \right| + \left| f^{(n)}(b) \right| \right) \right. \\
& \quad \left. + 2 \left[ \frac{\beta(b-a)^2}{12} \right]^{1/q} \right\},
\end{aligned}$$

where  $\frac{1}{p} + \frac{1}{q} = 1$ .

*Proof.* (1) If  $q = 1$ , then it follows from the fact of  $|f| \in SX(h, \beta, [a, b])$  that

$$\begin{aligned}
& \left| \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] f^{(n)} \left( \frac{1+t}{2} a + \frac{1-t}{2} b \right) dt \right| \\
& \leqslant \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] h \left( \frac{1+t}{2} \right) dt \left| f^{(n)}(a) \right| \\
& \quad + \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] h \left( \frac{1-t}{2} \right) dt \left| f^{(n)}(b) \right| \\
& \quad - \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] \beta \frac{(1-t)(1+t)}{4} (b-a)^2 dt \\
& = 2^n \int_{1/2}^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - 2^\alpha (1-t)^{\alpha+n-1} \right] h(t) dt \left| f^{(n)}(a) \right| \\
& \quad + 2^n \int_0^{1/2} \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} t^{n-1} - 2^\alpha t^{\alpha+n-1} \right] h(t) dt \left| f^{(n)}(b) \right| \\
& \quad - \frac{n(n+3) \prod_{k=1}^{n+2} (\alpha+k) - (n+2)!(\alpha+n+3)}{4(n+2)!(\alpha+n+1)(\alpha+n+2)} \beta (b-a)^2 \\
& = 2^n \left( \mathcal{C} \left| f^{(n)}(a) \right| + \mathcal{D} \left| f^{(n)}(b) \right| \right) \\
& \quad - \frac{n(n+3) \prod_{k=1}^{n+2} (\alpha+k) - (n+2)!(\alpha+n+3)}{4(n+2)!(\alpha+n+1)(\alpha+n+2)} \beta (b-a)^2.
\end{aligned}$$

Similarly,

$$\begin{aligned} & \left| \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] f^{(n)} \left( \frac{1-t}{2}a + \frac{1+t}{2}b \right) dt \right| \\ & \leq 2^n \left( \mathcal{D} |f^{(n)}(a)| + \mathcal{C} |f^{(n)}(b)| \right) \\ & \quad - \frac{n(n+3) \prod_{k=1}^{n+2} (\alpha+k) - (n+2)!(\alpha+n+3)}{4(n+2)!(\alpha+n+1)(\alpha+n+2)} \beta(b-a)^2. \end{aligned}$$

Then we finish the proof of (i) by the above inequalities and Lemma 2.

(2) If  $1 < q < \infty$ , then power-mean inequality and the fact of  $|f|^q \in SX(h, \beta, [a, b])$  show that

$$\begin{aligned} & \left| \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] f^{(n)} \left( \frac{1+t}{2}a + \frac{1-t}{2}b \right) dt \right| \\ & \leq \left( \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] dt \right)^{1-1/q} \\ & \quad \times \left( \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] \left| f^{(n)} \left( \frac{1+t}{2}a + \frac{1-t}{2}b \right) \right|^q dt \right)^{1/q} \\ & \leq 2^{n/q} \left( \frac{\prod_{k=1}^{n-1} (\alpha+k)}{n!} - \frac{1}{\alpha+n} \right)^{1-1/q} \left\{ \mathcal{D} |f^{(n)}(a)|^q + \mathcal{C} |f^{(n)}(b)|^q \right. \\ & \quad \left. - \frac{n(n+3) \prod_{k=1}^{n+2} (\alpha+k) - (n+2)!(\alpha+n+3)}{2^{n+2}(n+2)!(\alpha+n+1)(\alpha+n+2)} \beta(b-a)^2 \right\}^{1/q}, \end{aligned}$$

and

$$\begin{aligned} & \left| \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] f^{(n)} \left( \frac{1-t}{2}a + \frac{1+t}{2}b \right) dt \right| \\ & \leq 2^{n/q} \left( \frac{\prod_{k=1}^{n-1} (\alpha+k)}{n!} - \frac{1}{\alpha+n} \right)^{1-1/q} \left\{ \mathcal{D} |f^{(n)}(b)|^q + \mathcal{C} |f^{(n)}(a)|^q \right. \\ & \quad \left. - \frac{n(n+3) \prod_{k=1}^{n+2} (\alpha+k) - (n+2)!(\alpha+n+3)}{2^{n+2}(n+2)!(\alpha+n+1)(\alpha+n+2)} \beta(b-a)^2 \right\}^{1/q}, \end{aligned}$$

which finish the proof of the first inequality in (ii) by Lemma 2 again.

The proof of the second inequality can be obtained by a similar method as in Theorem 2 (ii), we omit the details.

(3) If  $1 < q < \infty$ , then Hölder's inequality and  $|f|^q \in SX(h, \beta, [a, b])$  imply that

$$\begin{aligned} & \left| \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] f^{(n)} \left( \frac{1+t}{2}a + \frac{1-t}{2}b \right) dt \right| \\ & \leq \left( \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right]^p dt \right)^{1/p} \\ & \quad \times \left( \int_0^1 \left| f^{(n)} \left( \frac{1+t}{2}a + \frac{1-t}{2}b \right) \right|^q dt \right)^{1/q} \\ & \leq 2^{1/q} \left( \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} t^{n-1} - t^{\alpha+n-1} \right]^p dt \right)^{1/p} \\ & \quad \times \left[ \int_{1/2}^1 h(t) dt \left| f^{(n)}(a) \right|^q + \int_0^{1/2} h(t) dt \left| f^{(n)}(b) \right|^q - \frac{\beta(b-a)^2}{12} \right]^{1/q}, \end{aligned}$$

and

$$\begin{aligned} & \left| \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} (1-t)^{n-1} - (1-t)^{\alpha+n-1} \right] f^{(n)} \left( \frac{1-t}{2}a + \frac{1+t}{2}b \right) dt \right| \\ & \leq 2^{1/q} \left( \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} t^{n-1} - t^{\alpha+n-1} \right]^p dt \right)^{1/p} \\ & \quad \times \left[ \int_{1/2}^1 h(t) dt \left| f^{(n)}(b) \right|^q + \int_0^{1/2} h(t) dt \left| f^{(n)}(a) \right|^q - \frac{\beta(b-a)^2}{12} \right]^{1/q}. \end{aligned}$$

Therefore, we obtained the first inequality of (iii) by the above two inequalities and Lemma 2.

The second inequality is achieved by the same way in Theorem 2 (iii), we leave it to readers.  $\square$

Letting  $\beta \rightarrow 0$ , we take the following conclusion.

**COROLLARY 3.** Let  $f \in C^n([a, b])$  and  $f^{(n)} \in L([a, b]), n \in \mathbb{Z}^+$ . Suppose that  $|f^{(n)}|^q$  is an  $h$ -convex function with  $1 \leq q < \infty$ .

(i) If  $1 \leq q < \infty$ , then

$$\begin{aligned} & \left| \Re \left( f, \frac{a+b}{2} \right) \right| \\ & \leq \frac{(b-a)^n}{2^{n(1-1/q)+1} \prod_{k=1}^{n-1} (\alpha+k)} \left( \frac{\prod_{k=1}^{n-1} (\alpha+k)}{n!} - \frac{1}{\alpha+n} \right)^{1-1/q} \\ & \quad \times \left[ \left( \mathcal{C} \left| f^{(n)}(a) \right|^q + \mathcal{D} \left| f^{(n)}(b) \right|^q \right)^{1/q} + \left( \mathcal{D} \left| f^{(n)}(a) \right|^q + \mathcal{C} \left| f^{(n)}(b) \right|^q \right)^{1/q} \right] \end{aligned}$$

$$\leq \frac{(b-a)^n}{2^{n(1-1/q)+1} \prod_{k=1}^{n-1} (\alpha+k)} \left( \frac{\prod_{k=1}^{n-1} (\alpha+k)}{n!} - \frac{1}{\alpha+n} \right)^{1-1/q} \\ \times \left( \mathcal{C}^{1/q} + \mathcal{D}^{1/q} \right) \left( |f^{(n)}(a)| + |f^{(n)}(b)| \right).$$

(ii) If  $1 < q < \infty$ , then

$$\begin{aligned} & \left| \mathfrak{R} \left( f, \frac{a+b}{2} \right) \right| \\ & \leq \frac{(b-a)^n}{2^{n+1-1/q} \prod_{k=1}^{n-1} (\alpha+k)} \left( \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} t^{n-1} - t^{\alpha+n-1} \right]^p dt \right)^{1/p} \\ & \quad \times \left\{ \left[ \int_{1/2}^1 h(t) dt |f^{(n)}(b)|^q + \int_0^{1/2} h(t) dt |f^{(n)}(a)|^q \right]^{1/q} \right. \\ & \quad \left. + \left[ \int_{1/2}^1 h(t) dt |f^{(n)}(a)|^q + \int_0^{1/2} h(t) dt |f^{(n)}(b)|^q \right]^{1/q} \right\} \\ & \leq \frac{(b-a)^n}{2^{n+1-1/q} \prod_{k=1}^{n-1} (\alpha+k)} \left( \int_0^1 \left[ \frac{\prod_{k=1}^{n-1} (\alpha+k)}{(n-1)!} t^{n-1} - t^{\alpha+n-1} \right]^p dt \right)^{1/p} \\ & \quad \times \left[ \left( \int_0^{1/2} h(t) dt \right)^{1/q} + \left( \int_{1/2}^1 h(t) dt \right)^{1/q} \right] \left( |f^{(n)}(a)| + |f^{(n)}(b)| \right), \end{aligned}$$

where  $\frac{1}{p} + \frac{1}{q} = 1$ .

**REMARK.** If  $n = 1$  and  $h(t) = t$ , then Corollary 3 reduces to Theorem F. As a special case of Corollary 3, we have the following results.

**COROLLARY 4.** Let  $f \in C^2([a, b])$  and  $f'' \in L([a, b])$ . Suppose that  $|f''|^q$  is a convex function with  $1 \leq q < \infty$ .

(i) If  $1 \leq q < \infty$ , then

$$\begin{aligned} & \left| \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{\left(\frac{a+b}{2}\right)}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)}^\alpha f(b) \right] - \frac{f(a) + f(b)}{2} \right| \\ & \leq \frac{(b-a)^2}{2^{3+1/q} (\alpha+1)} \left( \frac{\alpha+1}{2} - \frac{1}{\alpha+2} \right)^{1-1/q} \\ & \quad \times \left\{ \left[ \left( \frac{\alpha+1}{3} - \frac{1}{\alpha+3} \right) |f''(a)|^q + \left( \frac{2(\alpha+1)}{3} - \frac{2}{\alpha+2} + \frac{1}{\alpha+3} \right) |f''(b)|^q \right]^{1/q} \right. \\ & \quad \left. + \left[ \left( \frac{\alpha+1}{3} - \frac{1}{\alpha+3} \right) |f''(b)|^q + \left( \frac{2(\alpha+1)}{3} - \frac{2}{\alpha+2} + \frac{1}{\alpha+3} \right) |f''(a)|^q \right]^{1/q} \right\} \end{aligned}$$

$$\leq \frac{(b-a)^2}{2^{3+1/q}(\alpha+1)} \left( \frac{\alpha+1}{2} - \frac{1}{\alpha+2} \right)^{1-1/q} \\ \times \left[ \left( \frac{\alpha+1}{3} - \frac{1}{\alpha+3} \right)^{1/q} + \left( \frac{2(\alpha+1)}{3} - \frac{2}{\alpha+2} + \frac{1}{\alpha+3} \right)^{1/q} \right] (|f''(a)| + |f''(b)|).$$

(ii) If  $1 < q < \infty$ , then

$$\left| \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{(\frac{a+b}{2})^-}^\alpha f(a) + J_{(\frac{a+b}{2})^+}^\alpha f(b) \right] - \frac{f(a)+f(b)}{2} \right| \\ \leq \frac{(b-a)^2}{2^{3+2/q}(\alpha+1)} \left( \int_0^1 [(\alpha+1)t - t^{\alpha+1}]^p dt \right)^{1/p} \\ \times \left[ (3|f''(b)|^q + |f''(a)|^q)^{1/q} + (3|f''(a)|^q + |f''(b)|^q)^{1/q} \right] \\ \leq \frac{(1+3^{1/q})(b-a)^2}{2^{3+2/q}(\alpha+1)} \left( \int_0^1 [(\alpha+1)t - t^{\alpha+1}]^p dt \right)^{1/p} (|f''(a)| + |f''(b)|) \\ \leq \frac{(b-a)^2}{2^{1+2/q}(\alpha+1)} \left( \int_0^1 [(\alpha+1)t - t^{\alpha+1}]^p dt \right)^{1/p} (|f''(a)| + |f''(b)|),$$

where  $\frac{1}{p} + \frac{1}{q} = 1$ .

REMARK. Especially, if taking  $q = 1$  in Corollary 4 (i), we have

$$\left| \frac{2^{\alpha-1}\Gamma(\alpha+1)}{(b-a)^\alpha} \left[ J_{(\frac{a+b}{2})^-}^\alpha f(a) + J_{(\frac{a+b}{2})^+}^\alpha f(b) \right] - \frac{f(a)+f(b)}{2} \right| \\ \leq \frac{(b-a)^2}{16} \left[ 1 - \frac{2}{(\alpha+1)(\alpha+2)} \right] (|f''(a)| + |f''(b)|).$$

*Acknowledgements.* The authors would like to express their deep thanks to the referees for many helpful comments and suggestions.

## REFERENCES

- [1] M. ALOMARI, M. DARUS AND S. DRAGOMIR, *New inequalities of Hermite-Hadamard type for functions whose second derivatives absolute values are Quasi-convex*, Tamkang J. Math., **41** (4) (2010), 353–359.
- [2] M. ALOMARI, M. DARUS AND U. KIRMACI, *Some inequalities of Hermite-Hadamard type for s-convex functions*, Acta Math. Scientia, **31** B (4) (2011), 1643–1652.
- [3] H. ANGULO, J. GIMENEZ, A. MOROS AND K. NIKODEM, *On strongly h-convex functions*, Ann. Funct. Anal., **2** (2) (2011), 85–91.
- [4] M. AVCI, H. KAVURMACI AND M. ÖZDEMİR, *New inequalities of Hermite-Hadamard type via s-convex functions in the second sense with applications*, Appl. Math. Comput., **217** (2011), 5171–5176.
- [5] H. BUDAK, *New Hermite-Hadamard type inequalities for convex mappings utilizing generalized fractional integrals*, Filomat, **33** (8) (2019), 2329–2344.
- [6] H. BUDAK, F. ERTUĞRAL AND M. SARIKAYA, *New generalization of Hermite-Hadamard type inequalities via generalized fractional integrals*, Ann. University Craiova, Math. Comput. Sci. Ser., **47** (2) (2020), 369–386.

- [7] S. DRAGOMIR AND R. AGARWAL, *Two inequalities for differentiable mappings and applications to special means of real numbers and trapezoidal formula*, Appl. Math. Lett., **11** (5) (1998), 91–95.
- [8] S. DRAGOMIR, M. BHATTI, M. IQBAL AND M. MUDDASSAR, *Some new Hermite-Hadamard's type fractional integral inequalities*, J. Comput. Anal. Appl., **18** (4) (2015), 655–661.
- [9] S. DRAGOMIR AND S. FITZPATRICK, *The Hadamard's inequality for  $s$ -convex functions in the second sense*, Demonstratio Math., **32** (4) (1999), 687–696.
- [10] S. DRAGOMIR AND C. PEARCE, *Selected topics on Hermite-Hadamard inequalities and Applications*, RGMIA Monographs, Victoria University, 2000, (online: <http://rgmia.vu.edu.au/monographs/>).
- [11] S. DRAGOMIR, J. PEČARIĆ AND L. PERSSON, *Some inequalities of Hadamard type*, Soochow J. Math., **21** (1995), 335–241.
- [12] A. HAZY, *Bernstein-Doetsch type results for  $h$ -convex functions*, Math. Inequal. Appl., **14** (3) (2011), 499–508.
- [13] H. KAVURMACI, M. AVCI AND M. ÖZDEMİR, *New inequalities of Hermite-Hadamard type for convex functions with applications*, J. Inequal. Appl., **2011** (86) (2011), 1–11.
- [14] U. KIRMAKİ, *Inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula*, Appl. Math. Comput., **147** (2004), 137–146.
- [15] U. KIRMAKİ AND M. ÖZDEMİR, *On some inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula*, Appl. Math. Comput., **153** (2004), 361–368.
- [16] M. LATIF, *On some inequalities for  $h$ -convex functions*, Int. J. Math. Anal., **4** (30) (2010), 1473–1482.
- [17] N. MERENTES AND K. NIKODEM, *Remarks on strongly convex functions*, Aequationes Math., **80** (2010), 193–199.
- [18] K. NIKODEM, *Strongly convex functions and related classes of functions*, in: Th. M. Rassias (Ed.) *Handbook of Functional Equations. Functional Inequalities*, Springer Optimizations and Its Applications, **95** (2015), 365–405.
- [19] K. NIKODEM, J. SÁNCHEZ AND L. SÁNCHEZ, *Jensen and Hermite-Hadamard inequalities for strongly convex set-valued maps*, Math. Aeterna, **4**, (2014), 979–987.
- [20] M. NOOR AND M. AWAN, *Some integral inequalities for two kinds of convexities via fractional integrals*, TJMM, **5** (2) (2013), 129–136.
- [21] M. ÖZDEMİR, M. AVCI-ARDINÇ AND H. KAVURMACI-ÖNALAN, *Hermite-Hadamard type inequalities for  $s$ -convex and  $s$ -concave functions via fractional integrals*, Turkish J. Science, **1** (1) (2016), 28–40.
- [22] M. ÖZDEMİR, M. AVCI AND E. SET, *On some inequalities of Hermite-Hadamard type via  $m$ -convexity*, Appl. Math. Letters, **23** (2010), 1065–1070.
- [23] B. POLYAK, *Existence theorems and convergence of minimizing sequences in extremum problems with restrictions*, Soviet Math. Dokl., **2** (7) (1966), 72–75.
- [24] E. POLOVINKIN, *Strongly convex analysis*, Sbornik Math., **187** (2) (1996), 103–130.
- [25] C. PEARCE AND J. PEČARIĆ, *Inequalities for differentiable mappings with application to special means and quadrature formula*, Appl. Math. Lett., **13** (2000), 51–55.
- [26] C. PEARCE AND A. RUBINOV,  *$P$ -functions, quasi-convex functions and Hadamard-type inequalities*, J. Math. Anal. Appl., **240** (1) (1999), 92–104.
- [27] S. QAISAR, F. AHMAD, S. DRAGOMIR AND M. IQBAL, *New Hermite-Hadamard inequalities via fractional integrals, whose absolute values of second derivatives is  $P$ -convex*, J. Math. Inequal. **12** (3) (2018), 655–664.
- [28] M. SARIKAYA AND H. YILDIRIM, *On Hermite-Hadamard type inequalities for Riemann-Liouville fractional integrals*, Miskolc Math. Notes, **17** (2) (2017), 1049–1059.
- [29] M. SARIKAYA, A. SAGLAM AND H. YILDIRIM, *On some Hadamard-type inequalities for  $h$ -convex functions*, J. Math. Inequal., **2** (3) (2008), 335–341.
- [30] M. SARIKAYA, E. SET, H. YALDIZ AND N. BAŞAK, *Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities*, Math. Comput. Modelling, **57** (2013), 2403–2407.
- [31] E. SET, M. SARIKAYA AND A. GÖZPINART, *Some Hermite-Hadamard type inequalities for convex functions via conformable fractional integrals and related inequalities*, Creat. Math. Inform., **2** (26) (2017), 221–229.
- [32] S. TURHAN, I. ISCAN AND M. KUNT, *Hermite-Hadamard type inequalities for  $n$ -times differentiable convex functions via Riemann-Liouville fractional integrals*, Filomat, **32** (16) (2018), 5611–5622.
- [33] S. VAROŠANEC, *On  $h$ -convexity*, J. Math. Anal. Appl., **326** (1) (2007), 303–311.

- [34] J. VIAL, *Strong convexity of sets and functions*, J. Math. Econom., **9** (1982), 187–205.
- [35] J. VIAL, *Strong and weak convexity of sets and functions*, Math. Oper. Res., **8** (2) (1983), 231–259.
- [36] J. WANG, X. LI, M. FEČKAN AND Y. ZHOU, *Hermite-Hadamard-type inequalities for Riemann-Liouville fractional integrals via two kind of convexity*, Appl. Anal., **92** (11) (2013), 2241–2253.
- [37] J. WANG, X. LI AND Y. ZHOU, *Hermite-Hadamard-type inequalities involving Riemann-Liouville fractional integrals via  $s$ -convex functions and applications to special means*, Filomat, **30** (2016), 1143–1150.
- [38] X. WANG, J. RUAN AND X. MA, *On the Hermite-Hadamard inequalities for  $h$ -convex functions on balls and ellipsoids*, Filomat, **33** (18) (2019), 5817–5886.
- [39] F. ZHANG, G. FARID AND S. AKBAR, *Inequalities for Riemann-Liouville fractional integrals of strongly  $(s,m)$ -convex functions*, J. Math., (2021), Article ID 5577203, 14 pages, 2021.

(Received March 5, 2022)

Yi Xing

Department of Mathematics  
Zhejiang International Studies University  
Hangzhou 310014, China  
e-mail: 947228976@qq.com

Chaoqun Jiang

Department of Mathematics  
Zhejiang International Studies University  
Hangzhou 310014, China  
e-mail: 932963926@qq.com

Jianmiao Ruan

Department of Mathematics  
Zhejiang International Studies University  
Hangzhou 310014, China  
e-mail: rjmath@163.com