

POLYNOMIAL DIFFERENTIATION COMPOSITION OPERATORS FROM H^p SPACES TO WEIGHTED-TYPE SPACES ON THE UNIT BALL

STEVO STEVIĆ* AND SEI-ICHIRO UEKI

(Communicated by L. Mihoković)

Abstract. We characterize the boundedness, compactness, and estimate essential norm of a polynomial differentiation composition operator from the Hardy space H^p to the weighted-type spaces of holomorphic functions on the unit ball.

1. Introduction

Let \mathbb{N}_0 be the set of nonnegative integers. If $k, l \in \mathbb{N}_0$, $k \leq l$, then the notation $j = \overline{k, l}$ is an abbreviation for the notation $j = k, k+1, \dots, l$. Let $\mathbb{B} = \mathbb{B}^n \subset \mathbb{C}^n$ be the open unit ball, $\mathbb{S} = \partial\mathbb{B}$ its boundary, $d\sigma$ the normalized Lebesgue measure on \mathbb{S} , $\mathbb{D} = \mathbb{B}^1$, $\langle z, w \rangle$ the inner product in \mathbb{C}^n , $|z| = \langle z, z \rangle^{1/2}$, D_j the partial derivative operator

$$D_j f(z) = \frac{\partial f}{\partial z_j}(z), \quad j \in \{1, 2, \dots, n\},$$

$S(\Omega)$ the family of holomorphic self-maps of a domain Ω , $H(\Omega)$ the space of holomorphic functions on Ω ([22, 23, 48]), and $H^p(\mathbb{B}) = H^p$, $p > 0$, the Hardy space consisting of all $f \in H(\mathbb{B})$ such that

$$\|f\|_{H^p} = \sup_{0 \leq r < 1} \left(\int_{\mathbb{S}} |f(r\zeta)|^p d\sigma(\zeta) \right)^{1/p} < +\infty,$$

(see, e.g., [23, 48]). For $p \geq 1$ it is a Banach space.

By $W(\Omega)$ we denote the family of positive and continuous functions on Ω and call them weights. Let $\mu \in W(\mathbb{B})$. The weighted-type space $H_\mu^\infty(\mathbb{B}) = H_\mu^\infty$ is defined as follows

$$H_\mu^\infty(\mathbb{B}) := \left\{ f \in H(\mathbb{B}) : \|f\|_{H_\mu^\infty} := \sup_{z \in \mathbb{B}} \mu(z) |f(z)| < +\infty \right\}.$$

Mathematics subject classification (2020): Primary 47B38; Secondary 47B33.

Keywords and phrases: Polynomial differentiation composition operator, Hardy space, weighted-type space, boundedness, compactness, essential norm.

* Corresponding author.

For $\mu(z) \equiv 1$ we get the space of bounded holomorphic functions $H^\infty(\mathbb{B}) = H^\infty$ with the supremum norm $\|\cdot\|_\infty$. The little weighted-type space $H_{\mu,0}^\infty(\mathbb{B}) = H_{\mu,0}^\infty$ is a closed subspace of H_μ^∞ consisting of $f \in H(\mathbb{B})$ such that

$$\lim_{|z| \rightarrow 1} \mu(z)|f(z)| = 0.$$

The spaces and operators on them and their generalizations have been studied a lot (see, for instance, [2, 10, 16, 21, 29, 30, 33, 36, 37, 38, 39, 43, 45, 50] and the related references therein).

Beside the differentiation operator $Df = f'$, some attention to researchers attracted the composition operator $C_\varphi f = f \circ \varphi$, where $\varphi \in S(\Omega)$, the multiplication operator $M_u f = uf$, where $u \in H(\Omega)$, as well as their products. Among the products containing differentiation operators, the operators DC_φ and $C_\varphi D$ have been studied among the first ones (see, e.g., [7, 14, 15, 19] and the references therein).

The following extension of the operator $C_\varphi D$ attracted also some attention

$$D_{\varphi,u}^m := M_u C_\varphi D^m \tag{1}$$

on subspaces of $H(\mathbb{D})$ (see, e.g., [8, 13, 16, 29, 32, 33, 44, 45, 46, 49, 50, 51, 52, 53, 54, 55]).

The following n -dimensional variant of operator (1)

$$\mathfrak{R}_{\varphi,u}^m := M_u C_\varphi \mathfrak{R}^m, \tag{2}$$

where \mathfrak{R} is the radial differentiation operator was introduced in [34]. The investigation was continued in [35, 38, 39].

Investigations of sums of the operators in (1) was initiated by Stević and Sharma. The first published results can be found in [40] and [41]. An extension of the sum in [40] and [41] appeared in [42]. The investigation was continued, for instance, in [1, 5, 6, 9, 17, 47]. For some other product type operators consult, e.g., [10, 11, 12, 20, 27, 28, 26, 31, 43] and the related references therein.

Investigations of sums of the operators in (2) was suggested by Stević soon after finishing [42], but the first published results can be found in recent paper [37]. Beside the sums he also suggested studying the polynomial differentiation composition operator of the form

$$P_{D,\varphi}^m f := \sum_{j=0}^m u_j C_\varphi D_{l_j} \cdots D_{l_1} f, \quad f \in H(\mathbb{B}), \tag{3}$$

where $m \in \mathbb{N}_0$, $u_j \in H(\mathbb{B})$, $j = \overline{0, m}$, and $\varphi \in S(\mathbb{B})$ (see [36]).

Let X and Y be two normed spaces. A linear operator $T : X \rightarrow Y$ is called bounded if there is $M \geq 0$ such that $\|Tf\|_Y \leq M\|f\|_X$ for every $f \in X$. If it maps bounded sets in X into relatively compact ones, then it is called compact [4, 24]. The essential norm of the operator $T : X \rightarrow Y$ is defined as follows

$$\|T\|_{e,X \rightarrow Y} = \inf\{\|T + K\|_{X \rightarrow Y} : K \text{ is compact from } X \text{ to } Y\},$$

where $\|\cdot\|_{X \rightarrow Y}$ denote the operator norm. The operator T is compact if and only if $\|T\|_{e, X \rightarrow Y} = 0$. We denote the unit ball in X by B_X .

There has been a huge recent interest in investigating the boundedness, compactness, and estimating essential norms of concrete operators on spaces of holomorphic functions (see, e.g., [3, 5, 6, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 18, 20, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 40, 41, 42, 37, 38, 39, 43, 44, 45, 46, 47, 49, 50, 51, 52, 53, 54, 55] and the references therein).

In this article we characterize the boundedness and compactness of the operator $P_{D, \varphi}^m : H^p \rightarrow H_\mu^\infty$ (or $H_{\mu, 0}^\infty$), for $p \geq 1$, and estimate the essential norm of the operator in the case $p > 1$.

Let C denote unspecified nonnegative constants. They can change from line to line. The notation $a \lesssim b$ (resp. $a \gtrsim b$) means that there is $C > 0$ such that $a \leq Cb$ (resp. $a \geq Cb$). If $a \lesssim b$ and $b \lesssim a$, then we use the notation $a \asymp b$.

2. Auxiliary results

Our first auxiliary result is a characterization for the compactness. It is proved in a standard way [25], because of which we omit the proof.

LEMMA 1. *Let $p \geq 1$, $u_j \in H(\mathbb{B})$, $j = \overline{0, m}$, $\varphi \in S(\mathbb{B})$, $\mu \in W(\mathbb{B})$ and $Y \in \{H_\mu^\infty(\mathbb{B}), H_{\mu, 0}^\infty(\mathbb{B})\}$. Then the bounded operator $P_{D, \varphi}^m : H^p(\mathbb{B}) \rightarrow Y$ is compact if and only if for any bounded sequence $(f_k)_{k \in \mathbb{N}} \subset H^p(\mathbb{B})$ such that $f_k \rightarrow 0$ uniformly on compacts of \mathbb{B} as $k \rightarrow +\infty$, we have*

$$\lim_{k \rightarrow +\infty} \|P_{D, \varphi}^m f_k\|_{H_\mu^\infty} = 0.$$

The following folklore lemma is a consequence of Cauchy’s estimate for derivatives and a known estimate for the point evaluation functional on $H^p(\mathbb{B})$ ([23, 48]).

LEMMA 2. *Let $p > 0$ and $N \in \mathbb{N}_0$. Then for every multi-index $\vec{l} = (l_1, l_2, \dots, l_j)$ such that $|\vec{l}| = N$, there is $C_{\vec{l}} > 0$ such that*

$$\left| \frac{\partial^N f(z)}{\partial z_{k_1}^{l_1} \partial z_{k_2}^{l_2} \dots \partial z_{k_j}^{l_j}} \right| \leq \frac{C_{\vec{l}} \|f\|_{H^p}}{(1 - |z|^2)^{\frac{N}{p} + N}},$$

for every $f \in H^p(\mathbb{B})$ and $z \in \mathbb{B}$.

The following result, which is a consequence of [23, Proposition 1.4.10] and monotonicity of the integral means, gives a known family of test functions in H^p space.

LEMMA 3. *Let $p > 0$, $a \geq 0$ and $w \in \mathbb{B}$. Then the function*

$$f_{w, a}(z) = \frac{(1 - |w|^2)^{\frac{a}{p} + a}}{(1 - \langle z, w \rangle)^{\frac{2a}{p} + a}}, \tag{4}$$

belongs to $H^p(\mathbb{B})$.

Moreover, we have

$$\sup_{w \in \mathbb{B}} \|f_{w,a}\|_{HP} \lesssim 1. \tag{5}$$

The following lemma is a known generalization of Lemma 1 in [18].

LEMMA 4. A closed set K in $H_{\mu,0}^\infty(\mathbb{B})$ is compact if and only if it is bounded and

$$\lim_{|z| \rightarrow 1} \sup_{f \in K} \mu(z) |f(z)| = 0.$$

The following lemma gives a useful family of test functions.

LEMMA 5. Let $p > 0$, $m \in \mathbb{N}$ and $w \in \mathbb{B}$. Then for each $s \in \{0, 1, \dots, m\}$ there are $c_k^{(s)}$, $k = \overline{0, m}$, such that the function

$$h_w^{(s)}(z) = \sum_{k=0}^m c_k^{(s)} f_{w,k}(z)$$

where $f_{w,a}$ is defined in (4), satisfies

$$D_{l_s} \cdots D_{l_1} h_w^{(s)}(w) = \frac{\overline{w}_{l_1} \overline{w}_{l_2} \cdots \overline{w}_{l_s}}{(1 - |w|^2)^{\frac{n}{p} + s}} \tag{6}$$

and

$$D_t \cdots D_{l_1} h_w^{(s)}(w) = 0, \tag{7}$$

for every $t \in \{0, 1, \dots, m\} \setminus \{s\}$.

We also have

$$\sup_{w \in \mathbb{B}} \|h_w^{(s)}\|_{HP} \lesssim 1. \tag{8}$$

Proof. Let

$$h_w(z) = \sum_{k=0}^m c_k f_{w,k}(z).$$

and $d_k = \frac{2n}{p} + k$, $k \in \mathbb{N}_0$. Then

$$D_{l_t} \cdots D_{l_1} h_w(z) = \sum_{k=0}^m c_k \frac{d_k d_{k+1} \cdots d_{k+t-1} \overline{w}_{l_1} \overline{w}_{l_2} \cdots \overline{w}_{l_t} (1 - |w|^2)^{\frac{n}{p} + k}}{(1 - \langle z, w \rangle)^{d_{k+t}}},$$

for $t \in \mathbb{N}_0$, and consequently

$$D_{l_t} \cdots D_{l_1} h_w(w) = \frac{\overline{w}_{l_1} \overline{w}_{l_2} \cdots \overline{w}_{l_t}}{(1 - |w|^2)^{\frac{n}{p} + t}} \sum_{k=0}^m c_k \prod_{l=0}^{t-1} d_{k+l},$$

for $t \in \mathbb{N}_0$.

Since the determinant of the system

$$\begin{bmatrix} 1 & 1 & \cdots & 1 \\ d_0 & d_1 & \cdots & d_m \\ \vdots & \vdots & \ddots & \vdots \\ \prod_{k=0}^{s-1} d_k & \prod_{k=0}^{s-1} d_{k+1} & \cdots & \prod_{k=0}^{s-1} d_{k+m} \\ \vdots & \vdots & \ddots & \vdots \\ \prod_{k=0}^{m-1} d_k & \prod_{k=0}^{m-1} d_{k+1} & \cdots & \prod_{k=0}^{m-1} d_{k+m} \end{bmatrix} \begin{bmatrix} c_0 \\ c_1 \\ \vdots \\ c_s \\ \vdots \\ c_m \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \tag{9}$$

is not equal to zero ([30, Lemma 3]), we have that for any $s \in \{0, 1, \dots, m\}$, it has a unique solution $c_k := c_k^{(s)}$, $k = \overline{0, m}$. It is easy to see that the function satisfying (6) and (7) is given by $h_w^{(s)}(z) := \sum_{k=0}^m c_k^{(s)} f_{w,k}(z)$, and that (5) implies (8). \square

3. Main results

The main results in the paper are presented in this section.

THEOREM 1. *Let $p \geq 1$, $m \in \mathbb{N}$, $\mu \in W(\mathbb{B})$, $u_j \in H(\mathbb{B})$, $j = \overline{0, m}$, $\varphi = (\varphi_1, \dots, \varphi_n) \in S(\mathbb{B})$,*

$$\min_{j=1, n} \inf_{z \in \mathbb{B}} |\varphi_j(z)| \geq \delta > 0. \tag{10}$$

Then $P_{D, \varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ is bounded if and only if

$$L_j := \sup_{z \in \mathbb{B}} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p} + j}} < +\infty, \quad j = \overline{0, m}. \tag{11}$$

Moreover, if the operator is bounded, then we have

$$\|P_{D, \varphi}^m\|_{H^p \rightarrow H_\mu^\infty} \asymp \sum_{j=0}^m L_j. \tag{12}$$

Proof. Suppose that $P_{D, \varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ is bounded. Lemma 5 implies that for each $s \in \{0, 1, \dots, m\}$ and $\varphi(w) \in \mathbb{B}$, there is $h_{\varphi(w)}^{(s)} \in H^p(\mathbb{B})$ such that

$$D_{l_s} \cdots D_{l_1} h_{\varphi(w)}^{(s)}(\varphi(w)) = \frac{\overline{\varphi_{l_1}(w)} \overline{\varphi_{l_2}(w)} \cdots \overline{\varphi_{l_s}(w)}}{(1 - |\varphi(w)|^2)^{\frac{n}{p} + s}}, \tag{13}$$

$$D_{l_t} \cdots D_{l_1} h_{\varphi(w)}^{(s)}(\varphi(w)) = 0, \tag{14}$$

for every $t \in \{0, 1, \dots, m\} \setminus \{s\}$. We also have $\sup_{w \in \mathbb{B}} \|h_{\varphi(w)}^{(s)}\|_{H^p} < +\infty$.

This together with the boundedness, (13), (14), as well as (10), implies

$$\begin{aligned} \|P_{D,\varphi}^m\|_{H^p \rightarrow H_\mu^\infty} &\gtrsim \|P_{D,\varphi}^m h_{\varphi(w)}^{(s)}\|_{H_\mu^\infty} \\ &= \sup_{z \in \mathbb{B}} \mu(z) \left| \sum_{j=0}^m u_j(z) D_{l_j} \cdots D_{l_1} h_{\varphi(w)}^{(s)}(\varphi(z)) \right| \\ &\geq \mu(w) \left| \sum_{j=0}^m u_j(w) D_{l_j} \cdots D_{l_1} h_{\varphi(w)}^{(s)}(\varphi(w)) \right| \\ &= \mu(w) |u_s(w)| \frac{|\overline{\varphi_{l_1}(w)}| \cdots |\overline{\varphi_{l_s}(w)}|}{(1 - |\varphi(w)|^2)^{\frac{s}{p} + s}} \\ &\geq \delta^s \frac{\mu(w) |u_s(w)|}{(1 - |\varphi(w)|^2)^{\frac{s}{p} + s}}, \end{aligned} \tag{15}$$

for every $w \in \mathbb{B}$, from which it easily follows that $L_s < +\infty$, $s \in \{0, 1, \dots, m\}$, and

$$L_s \lesssim \|P_{D,\varphi}^m\|_{H^p \rightarrow H_\mu^\infty}, \quad s = \overline{0, m},$$

and consequently

$$\sum_{j=0}^m L_j \lesssim \|P_{D,\varphi}^m\|_{H^p \rightarrow H_\mu^\infty}. \tag{16}$$

If (11) holds, then Lemma 2 implies

$$\begin{aligned} \mu(z) |P_{D,\varphi}^m f(z)| &= \mu(z) \left| \sum_{j=0}^m u_j(z) D_{l_j} \cdots D_{l_1} f(\varphi(z)) \right| \\ &\leq C \sum_{j=0}^m \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{j}{p} + j}} \|f\|_{H^p}, \end{aligned} \tag{17}$$

from which along with (11), the boundedness of $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ follows, as well as the asymptotic estimate

$$\|P_{D,\varphi}^m\|_{H^p \rightarrow H_\mu^\infty} \lesssim \sum_{j=0}^m L_j. \tag{18}$$

Asymptotic estimates (16) and (18) imply (12). \square

THEOREM 2. *Let $p \geq 1$, $m \in \mathbb{N}$, $u_j \in H(\mathbb{B})$, $j = \overline{0, m}$, $\varphi \in S(\mathbb{B})$, and $\mu \in W(\mathbb{B})$. Then $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu,0}^\infty(\mathbb{B})$ is bounded if and only if $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ is bounded and*

$$\lim_{|z| \rightarrow 1} \mu(z) |u_j(z)| = 0, \quad j = \overline{0, m}. \tag{19}$$

Proof. If $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ is bounded and (19) holds, then for any polynomial p we have

$$\begin{aligned} \mu(z) \left| \sum_{j=0}^m u_j(z) D_{l_j} \cdots D_{l_1} p(\varphi(z)) \right| &\leq \sum_{j=0}^m \mu(z) |u_j(z)| \|D_{l_j} \cdots D_{l_1} p(\varphi(z))\| \\ &\leq \sum_{j=0}^m \mu(z) |u_j(z)| \|D_{l_j} \cdots D_{l_1} p\|_\infty, \end{aligned}$$

from which together with (19) it easily follows that $P_{D,\varphi}^m p \in H_{\mu,0}^\infty(\mathbb{B})$.

Since for every $f \in H^p(\mathbb{B})$ there is a sequence of polynomials $(p_k)_{k \in \mathbb{N}}$ such that

$$\lim_{k \rightarrow +\infty} \|f - p_k\|_{H^p} = 0,$$

and the following inequality holds

$$\|P_{D,\varphi}^m f - P_{D,\varphi}^m p_k\|_{H_\mu^\infty} \leq \|P_{D,\varphi}^m\|_{H^p \rightarrow H_\mu^\infty} \|f - p_k\|_{H^p},$$

by letting $k \rightarrow +\infty$, and using the fact that $\overline{H_{\mu,0}^\infty(\mathbb{B})} = H_\mu^\infty(\mathbb{B})$, we have $P_{D,\varphi}^m f \in H_{\mu,0}^\infty(\mathbb{B})$, from which the boundedness of $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu,0}^\infty(\mathbb{B})$ follows.

Suppose that $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu,0}^\infty(\mathbb{B})$ is bounded. Then $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ is also such. Since $f_0(z) \equiv 1 \in H^p(\mathbb{B})$, we have $P_{D,\varphi}^m(f_0) \in H_{\mu,0}^\infty(\mathbb{B})$, that is

$$\lim_{|z| \rightarrow 1} \mu(z) |P_{D,\varphi}^m(f_0)(z)| = \lim_{|z| \rightarrow 1} \mu(z) |u_0(z)| = 0. \tag{20}$$

Hence (19) holds for $j = 0$.

Suppose that for some $s \in \{1, 2, \dots, m - 1\}$, (19) holds for $0 \leq j \leq s$. Let

$$f_{s+1}(z) = z_{l_1} z_{l_2} \cdots z_{l_{s+1}}.$$

Since $f_{s+1} \in H^p(\mathbb{B})$, we have $P_{D,\varphi}^m(f_{s+1}) \in H_{\mu,0}^\infty(\mathbb{B})$. Note that

$$f_{s+1}(z) = z_1^{\alpha_1} \cdots z_n^{\alpha_n},$$

where $\alpha_j \in \mathbb{N}_0$, $j = \overline{1, n}$, are such that $\sum_{j=1}^n \alpha_j = s + 1$. It is easy to see that for each $t \in \mathbb{N}_0$, $0 \leq t \leq s + 1$

$$D_{j_t} \cdots D_{j_1} f_{s+1}(z) = \gamma_t z_1^{\alpha_1 - k_1(t)} \cdots z_n^{\alpha_n - k_n(t)},$$

for some $\gamma_t \in \mathbb{N}$, where $k_i(t)$ is the number of operators D_i in the product $D_{j_t} \cdots D_{j_1}$. Note that $\sum_{j=1}^n k_i(t) = t$ and

$$D_{j_{s+1}} \cdots D_{j_1} f_{s+1}(z) = \gamma_{s+1}, \tag{21}$$

for some $\gamma_{s+1} \in \mathbb{N}$. Hence

$$\lim_{|z| \rightarrow 1} \mu(z) |P_{D,\varphi}^m f_{s+1}(z)| = \lim_{|z| \rightarrow 1} \mu(z) \left| \sum_{j=0}^{s+1} u_j(z) \gamma_j \prod_{i=1}^n (\varphi_i(z))^{\alpha_i - k_i(j)} \right| = 0,$$

from which, along with $|\varphi_i(z)| < 1$, $i = \overline{1, n}$, $\alpha_i \geq k_i(j)$, for $i = \overline{1, n}$, $j = \overline{0, s+1}$, the hypothesis $u_j \in H_{\mu, 0}^\infty(\mathbb{B})$, $j = \overline{0, s}$, (21) and $\gamma_{s+1} \neq 0$, we obtain

$$\lim_{|z| \rightarrow 1} \mu(z) |u_{s+1}(z)| = 0.$$

Hence (19) holds for $j = \overline{0, m}$. \square

THEOREM 3. *Let $p \geq 1$, $m \in \mathbb{N}$, $u_j \in H(\mathbb{B})$, $j = \overline{0, m}$, $\varphi \in S(\mathbb{B})$, $\mu \in W(\mathbb{B})$, and (10) holds. Then the operator $P_{D, \varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ is compact if and only if the operator is bounded and the following condition holds*

$$\lim_{|\varphi(z)| \rightarrow 1} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p} + j}} = 0, \tag{22}$$

for $j \in \{0, 1, \dots, m\}$.

Proof. Suppose $P_{D, \varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ is bounded and (22) holds. Then for every $\varepsilon > 0$ there is $\delta \in (0, 1)$ such that for $|\varphi(z)| > \delta$

$$\frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p} + j}} < \varepsilon, \quad j = \overline{0, m}. \tag{23}$$

Suppose that $\sup_{k \in \mathbb{N}} \|f_k\|_{H^p} \leq M$ and

$$f_k \rightarrow 0 \tag{24}$$

uniformly on compacts of \mathbb{B} . Let $K_\delta = \{z \in \mathbb{B} : |\varphi(z)| > \delta\}$. Then Lemma 2 and (23) imply

$$\begin{aligned} \|P_{D, \varphi}^m f_k\|_{H_\mu^\infty} &= \sup_{z \in \mathbb{B}} \mu(z) \left| \sum_{j=0}^m u_j(z) D_{l_j} \cdots D_{l_1} f_k(\varphi(z)) \right| \\ &\leq \sup_{z \in K_\delta} \mu(z) \left| \sum_{j=0}^m u_j(z) D_{l_j} \cdots D_{l_1} f_k(\varphi(z)) \right| \\ &\quad + \sup_{z \in \mathbb{B} \setminus K_\delta} \mu(z) \left| \sum_{j=0}^m u_j(z) D_{l_j} \cdots D_{l_1} f_k(\varphi(z)) \right| \\ &\leq C \sum_{j=0}^m \sup_{z \in K_\delta} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p} + j}} \|f_k\|_{H^p} \\ &\quad + C \sum_{j=0}^m \sup_{z \in \mathbb{B} \setminus K_\delta} \mu(z) |u_j(z)| |D_{l_j} \cdots D_{l_1} f_k(\varphi(z))| \\ &\leq (m+1)MC\varepsilon + C \sum_{j=0}^m \sup_{z \in \mathbb{B} \setminus K_\delta} \mu(z) |u_j(z)| \sup_{|\varphi(z)| \leq \delta} |D_{l_j} \cdots D_{l_1} f_k(\varphi(z))| \\ &\leq (m+1)MC\varepsilon + C \sum_{j=0}^m \|u_j\|_{H_\mu^\infty} \sup_{|w| \leq \delta} |D_{l_j} \cdots D_{l_1} f_k(w)|. \end{aligned} \tag{25}$$

Condition (24) together with Cauchy’s estimate imply

$$D_{l_j} \cdots D_{l_1} f_k \rightarrow 0, \tag{26}$$

uniformly on compacts of \mathbb{B} as $k \rightarrow +\infty$, for $j = \overline{0, m}$.

Let

$$f_s(z) = \prod_{j=1}^s z_{l_j}, \quad s = \overline{0, m}.$$

Arguing as in the proof of Theorem 2 we get $u_j \in H_\mu^\infty$, $j = \overline{0, m}$, from which along with (26), the compactness of $|w| \leq \delta$, and (25), we easily obtain

$$\lim_{k \rightarrow +\infty} \|P_{D, \varphi}^m f_k\|_{H_\mu^\infty} = 0.$$

This fact with Lemma 1 implies the compactness of $P_{D, \varphi}^m M_u : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$.

If $P_{D, \varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ is compact, then it is bounded. If $\|\varphi\|_\infty < 1$, then (22) holds.

Assume that $\|\varphi\|_\infty = 1$. Let $(z_k)_{k \in \mathbb{N}} \subset \mathbb{B}$ be a sequence such that $|\varphi(z_k)| \rightarrow 1$ as $k \rightarrow +\infty$, and

$$h_k^{(s)} := h_{\varphi(z_k)}^{(s)}, \quad s = \overline{0, m},$$

where $h_w^{(s)}$, $s = \overline{0, m}$, are as in Lemma 5. Then

$$\sup_{k \in \mathbb{N}} \|h_k^{(s)}\|_{H^p} < +\infty, \quad s = \overline{0, m}, \tag{27}$$

and $h_k^{(s)} \rightarrow 0$ uniformly on compacts of \mathbb{B} as $k \rightarrow +\infty$, for $s \in \{0, 1, \dots, m\}$. This along with Lemma 1 implies

$$\lim_{k \rightarrow +\infty} \|P_{D, \varphi}^m h_k^{(s)}\|_{H_\mu^\infty} = 0, \quad s = \overline{0, m}. \tag{28}$$

From (15) we have

$$\frac{\mu(z_k) |u_s(z_k)|}{(1 - |\varphi(z_k)|^2)^{\frac{n}{p} + s}} \lesssim \|P_{D, \varphi}^m h_k^{(s)}\|_{H_\mu^\infty}, \quad s = \overline{0, m}. \tag{29}$$

From (28) and (29), (22) easily follows. \square

When $p > 1$, we can estimate the essential norm of the bounded operator $P_{D, \varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ as follows.

THEOREM 4. *Let $p > 1$, $m \in \mathbb{N}$, $u_j \in H(\mathbb{B})$, $j = \overline{0, m}$, $\varphi \in S(\mathbb{B})$, $\mu \in W(\mathbb{B})$, and (10) holds. If the operator $P_{D, \varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$ is bounded then*

$$\|P_{D, \varphi}^m\|_{e, H^p \rightarrow H_\mu^\infty} \asymp \max_{j=1, \overline{m}} \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p} + j}}. \tag{30}$$

Proof. Let take a sequence $(z_k)_{k \in \mathbb{N}} \subset \mathbb{B}$ such that $|\varphi(z_k)| \rightarrow 1$ as $k \rightarrow +\infty$, and

$$h_k^{(s)} := h_{\varphi(z_k)}^{(s)}, \quad s = \overline{0, m},$$

where $h_w^{(s)}$, $s = \overline{0, m}$, are as in Lemma 5. Then (27) holds, and we have that $h_k^{(s)} \rightarrow 0$ uniformly on compacts of \mathbb{B} as $k \rightarrow +\infty$, for each $s \in \{0, 1, \dots, m\}$. Since the dual of $H^p(\mathbb{B})$ is known [48], it is easily verified that $h_k^{(s)} \rightarrow 0$ weakly in $H^p(\mathbb{B})$. Hence, we have

$$\lim_{k \rightarrow +\infty} \|Kh_k^{(s)}\|_{H_\mu^\infty} = 0$$

for any compact operator $K : H^p(\mathbb{B}) \rightarrow H_\mu^\infty(\mathbb{B})$.

From this, (13), (14) and (10) we have

$$\begin{aligned} \|P_{D,\varphi}^m\|_{e,H^p \rightarrow H_\mu^\infty} &\gtrsim \limsup_{k \rightarrow \infty} \left(\|P_{D,\varphi}^m h_k^{(s)}\|_{H_\mu^\infty} - \|Kh_k^{(s)}\|_{H_\mu^\infty} \right) \\ &\geq \limsup_{k \rightarrow \infty} \mu(z_k) |u_s(z_k)| \frac{|\varphi_{l_1}(z_k)| \cdots |\varphi_{l_s}(z_k)|}{(1 - |\varphi(z_k)|^2)^{\frac{n}{p} + s}} \\ &\geq \delta^s \limsup_{k \rightarrow \infty} \frac{\mu(z_k) |u_s(z_k)|}{(1 - |\varphi(z_k)|^2)^{\frac{n}{p} + s}} \end{aligned}$$

for each $s \in \{0, 1, \dots, m\}$. This implies that the lower estimate in (30) holds.

Next we prove the upper estimate in (30). For fixed t , $0 < t < 1$, put $C_t f(z) = f(tz)$. Since C_t is a compact operator on $H^p(\mathbb{B})$, $P_{D,\varphi}^m C_t$ is also compact from $H^p(\mathbb{B})$ into $H_\mu^\infty(\mathbb{B})$. Thus we have

$$\|P_{D,\varphi}^m\|_{e,H^p \rightarrow H_\mu^\infty} \leq \sup_{\|f\|_{H^p} \leq 1} \|P_{D,\varphi}^m f - P_{D,\varphi}^m C_t f\|_{H_\mu^\infty}. \tag{31}$$

Now we fix $f \in H^p(\mathbb{B})$ with $\|f\|_{H^p} \leq 1$ and R , $0 < R < 1$. Note that for each $z \in \mathbb{B}$ it holds that

$$\begin{aligned} &|P_{D,\varphi}^m f(z) - P_{D,\varphi}^m C_t f(z)| \\ &= \left| \sum_{j=0}^m u_j(z) \left\{ D_{l_j} \cdots D_{l_1} f(\varphi(z)) - t^j D_{l_j} \cdots D_{l_1} f(t\varphi(z)) \right\} \right|. \end{aligned}$$

By combining this with Lemma 2, we have

$$\begin{aligned} &\sup_{\|f\|_{H^p} \leq 1} \sup_{|\varphi(z)| > R} \mu(z) |P_{D,\varphi}^m f(z) - P_{D,\varphi}^m C_t f(z)| \\ &\lesssim \sum_{j=0}^m \sup_{|\varphi(z)| > R} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p} + j}} \\ &\lesssim \max_{j=0,m} \sup_{|\varphi(z)| > R} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p} + j}}. \end{aligned} \tag{32}$$

On the other hand, by using the mean value theorem and the Cauchy inequality, we obtain

$$\begin{aligned} & \sup_{|\varphi(z)| \leq R} \left| D_{l_j} \cdots D_{l_1} f(\varphi(z)) - D_{l_j} \cdots D_{l_1} f(t\varphi(z)) \right| \\ & \leq \sup_{|\varphi(z)| \leq R} (1-t) |\varphi(z)| \sup_{|w| \leq R} \left| \nabla(D_{l_j} \cdots D_{l_1} f)(w) \right| \\ & \lesssim \frac{R}{1-R} (1-t) \sup_{|w| \leq \frac{1+R}{2}} \left| D_{l_j} \cdots D_{l_1} f(w) \right|. \end{aligned}$$

Hence, Lemma 2 also shows that

$$\sup_{|\varphi(z)| \leq R} \left| D_{l_j} \cdots D_{l_1} f(\varphi(z)) - D_{l_j} \cdots D_{l_1} f(t\varphi(z)) \right| \lesssim \frac{R(1-t)}{(1-R)(1 - (\frac{1+R}{2})^2)^{\frac{n}{p}+j}} \tag{33}$$

for each $j \in \overline{0, m}$. Furthermore it follows from Lemma 2 that

$$\sup_{|\varphi(z)| \leq R} \left| D_{l_j} \cdots D_{l_1} f(t\varphi(z)) - t^j D_{l_j} \cdots D_{l_1} f(t\varphi(z)) \right| \lesssim \frac{1-t^j}{(1-R^2)^{\frac{n}{p}+j}} \tag{34}$$

for each $j \in \{0, 1, \dots, m\}$. Inequalities (33) and (34) give

$$\begin{aligned} & \sup_{\|f\|_{H^p} \leq 1} \sup_{|\varphi(z)| \leq R} \mu(z) \left| P_{D, \varphi}^m f(z) - P_{D, \varphi}^m C_t f(z) \right| \\ & \lesssim \sum_{j=0}^m \left\{ \frac{R(1-t)}{(1-R)(1 - (\frac{1+R}{2})^2)^{\frac{n}{p}+j}} + \frac{1-t^j}{(1-R^2)^{\frac{n}{p}+j}} \right\} \sup_{|\varphi(z)| \leq R} \mu(z) |u_j(z)| \\ & \rightarrow 0, \end{aligned} \tag{35}$$

as $t \rightarrow 1$. From (31), (32) and (35), we obtain

$$\|P_{D, \varphi}^m\|_{e, H^p \rightarrow H_{\mu}^{\infty}} \lesssim \max_{j \in \overline{0, m}} \sup_{|\varphi(z)| > R} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p}+j}}. \tag{36}$$

Letting $R \rightarrow 1^-$ in (36), we also obtain the upper estimate in (30). \square

THEOREM 5. *Let $p \geq 1$, $m \in \mathbb{N}$, $u_j \in H(\mathbb{B})$, $j \in \overline{0, m}$, $\varphi \in S(\mathbb{B})$, $\mu \in W(\mathbb{B})$, and condition (10) holds. Then the operator $P_{D, \varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu, 0}^{\infty}(\mathbb{B})$ is compact if and only if the operator is bounded and*

$$\lim_{|z| \rightarrow 1} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p}+j}} = 0, \quad j \in \overline{0, m}. \tag{37}$$

Proof. Assume (37) holds. Then (11) holds. From this and Theorem 1 the boundedness of $P_{D, \varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu}^{\infty}(\mathbb{B})$ easily follows. Letting $|z| \rightarrow 1$ in (17) and using (37), we have $P_{D, \varphi}^m f \in H_{\mu, 0}^{\infty}(\mathbb{B})$ for any $f \in H^p(\mathbb{B})$, from which the boundedness of

$P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu,0}^\infty(\mathbb{B})$ follows. Taking the supremum in (17) over \mathbb{B} and $B_{HP}(\mathbb{B})$, and using (11), we obtain

$$\sup_{f \in B_{HP}(\mathbb{B})} \sup_{z \in \mathbb{B}} \mu(z) |P_{D,\varphi}^m f(z)| \leq C \sum_{j=0}^m L_j < +\infty, \tag{38}$$

where $L_j, j = \overline{0, m}$, are the quantities in (11). So $\{P_{D,\varphi}^m f : f \in B_{HP}(\mathbb{B})\}$ is a bounded subset of $H_{\mu,0}^\infty(\mathbb{B})$. Taking the supremum in (17) over $B_{HP}(\mathbb{B})$ and letting $|z| \rightarrow 1$ we have

$$\lim_{|z| \rightarrow 1} \sup_{f \in B_{HP}(\mathbb{B})} \mu(z) |P_{D,\varphi}^m f(z)| = 0,$$

from which along with Lemma 4 the compactness of $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu,0}^\infty(\mathbb{B})$ follows.

If $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu,0}^\infty(\mathbb{B})$ is compact, then $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_\mu^p(\mathbb{B})$ is compact, from which and Theorem 3 we get (23). From Theorem 2 we get (19), so that there is $\eta \in (0, 1)$ such that

$$\mu(z) |u_j(z)| < \varepsilon (1 - \delta^2)^{\frac{\eta}{p} + j}, \quad j = \overline{0, m},$$

when $\eta < |z| < 1$, for ε chosen such that (23) holds, and consequently

$$\frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{\eta}{p} + j}} \leq \frac{\mu(z) |u_j(z)|}{(1 - \delta^2)^{\frac{\eta}{p} + j}} < \varepsilon, \quad j = \overline{0, m},$$

when $|\varphi(z)| \leq \delta$ and $\eta < |z| < 1$. This along with (23) imply (37). \square

In addition to Theorem 5, we also obtain the estimate for the essential norm of the operator $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu,0}^\infty(\mathbb{B})$, in the case $p > 1$.

THEOREM 6. *Let $p > 1, m \in \mathbb{N}, u_j \in H(\mathbb{B}), j = \overline{0, m}, \varphi \in S(\mathbb{B}), \mu \in W(\mathbb{B})$, and condition (10) holds. If the operator $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu,0}^\infty(\mathbb{B})$ is bounded then*

$$\|P_{D,\varphi}^m\|_{e, H^p \rightarrow H_{\mu,0}^\infty} \asymp \max_{j=\overline{1, m}} \limsup_{|z| \rightarrow 1} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{\eta}{p} + j}}. \tag{39}$$

Proof. By Theorem 2, the boundedness of $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu,0}^\infty(\mathbb{B})$ implies $u_j \in H_{\mu,0}^\infty(\mathbb{B})$ for $j = \overline{0, m}$. There are two cases to be considered.

Case $\|\varphi\|_\infty < 1$. In this case we see that $P_{D,\varphi}^m : H^p(\mathbb{B}) \rightarrow H_{\mu,0}^\infty(\mathbb{B})$ is compact, so $\|P_{D,\varphi}^m\|_{e, H^p \rightarrow H_{\mu,0}^\infty} = 0$. On the other hand, from $\|\varphi\|_\infty < 1$ and $u_j \in H_{\mu,0}^\infty(\mathbb{B})$ we have that the limit on the right-hand side of (39) equals to zero. Hence (39) holds in this case.

Case $\|\varphi\|_\infty = 1$. By Theorem 4, it is enough to prove that

$$\limsup_{|z| \rightarrow 1} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{\eta}{p} + j}} = \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(z) |u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{\eta}{p} + j}}, \tag{40}$$

for $j = \overline{0, m}$.

Note that

$$\limsup_{|z| \rightarrow 1} \frac{\mu(z)|u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p}+j}} \geq \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(z)|u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p}+j}}, \quad j = \overline{0, m}. \tag{41}$$

Assume that a sequence $(z_k)_{k \in \mathbb{N}} \subset \mathbb{B}$ satisfies

$$\limsup_{|z| \rightarrow 1} \frac{\mu(z)|u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p}+j}} = \lim_{k \rightarrow \infty} \frac{\mu(z_k)|u_j(z_k)|}{(1 - |\varphi(z_k)|^2)^{\frac{n}{p}+j}}, \quad j = \overline{0, m}.$$

If $\sup_{k \in \mathbb{N}} |\varphi(z_k)| < 1$, then since $u_j \in H_{\mu,0}^\infty(\mathbb{B})$, $j = \overline{0, m}$, we have that the first limit in (41) is zero and consequently the second one.

If $\sup_{k \in \mathbb{N}} |\varphi(z_k)| = 1$, then there is a subsequence $(\varphi(z_{k_l}))_{l \in \mathbb{N}}$ such that $|\varphi(z_{k_l})| \rightarrow 1$ as $l \rightarrow \infty$. Hence we obtain

$$\begin{aligned} \limsup_{|z| \rightarrow 1} \frac{\mu(z)|u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p}+j}} &= \lim_{l \rightarrow \infty} \frac{\mu(z_{k_l})|u_j(z_{k_l})|}{(1 - |\varphi(z_{k_l})|^2)^{\frac{n}{p}+j}} \\ &\leq \limsup_{|\varphi(z)| \rightarrow 1} \frac{\mu(z)|u_j(z)|}{(1 - |\varphi(z)|^2)^{\frac{n}{p}+j}}. \end{aligned} \tag{42}$$

From (41) and (42), (40) follows, finishing the proof of the theorem. \square

Acknowledgement. This research is partly supported by JSPS KAKENHI Grants-in-Aid for Scientific Research (C), Grant Number 21K03301.

Authors contributions. Stevo Stević proposed the investigation in the paper. The investigation was conducted by both authors who contributed equally to the writing of this paper. The authors read and approved the manuscript.

REFERENCES

- [1] M. S. AL GHAFRI, J. S. MANHAS, *On Stević-Sharma operators from weighted Bergman spaces to weighted-type spaces*, Math. Inequal. Appl. **23** (3) (2020), 1051–1077.
- [2] K. D. BIERSTEDT, W. H. SUMMERS, *Biduals of weighted Banach spaces of analytic functions*, J. Austral. Math. Soc. A **54** (1993), 70–79.
- [3] D. C. CHANG, S. LI, S. STEVIĆ, *On some integral operators on the unit polydisk and the unit ball*, Taiwanese J. Math. **11** (5) (2007), 1251–1286.
- [4] N. DUNFORD, J. T. SCHWARTZ, *Linear Operators I*, Interscience Publishers, Jon Willey and Sons, New York, 1958.
- [5] Z. GUO, L. LIU, Y. SHU, *On Stević-Sharma operators from the mixed-norm spaces to Zygmund-type spaces*, Math. Inequal. Appl. **24** (2) (2021), 445–461.
- [6] Z. GUO, Y. SHU, *On Stević-Sharma operators from Hardy spaces to Stević weighted spaces*, Math. Inequal. Appl. **23** (1) (2020), 217–229.
- [7] R. A. HIBSCHWEILER, N. PORTNOY, *Composition followed by differentiation between Bergman and Hardy spaces*, Rocky Mountain J. Math. **35** (3) (2005), 843–855.
- [8] Q. HU, X. ZHU, *Compact generalized weighted composition operators on the Bergman space*, Opuscula Math. **37** (2) (2017), 303–312.

- [9] Z. J. JIANG, *On Stević-Sharma operator from the Zygmund space to the Bloch-Orlicz space*, Adv. Differ. Equ. Vol. 2015, Article no. 228 (2015), 12 pp.
- [10] H. LI, Z. GUO, *Note on a Li-Stević integral-type operator from mixed-norm spaces to n th weighted spaces*, J. Math. Inequal. **11** (1) (2017), 77–85.
- [11] H. LI, Z. GUO, *On some product-type operators from area Nevanlinna spaces to Zygmund-type spaces*, Filomat **31** (3) (2017), 681–698.
- [12] S. LI, *Volterra composition operators between weighted Bergman spaces and Bloch type spaces*, J. Korean Math. Soc. **45** (1) (2008), 229–248.
- [13] S. LI, *Differences of generalized composition operators on the Bloch space*, J. Math. Anal. Appl. **394** (2012), 706–711.
- [14] S. LI, S. STEVIĆ, *Composition followed by differentiation from mixed-norm spaces to α -Bloch spaces*, Sb. Math. **199** (12) (2008), 1847–1857.
- [15] S. LI, S. STEVIĆ, *Products of composition and differentiation operators from Zygmund spaces to Bloch spaces and Bers spaces*, Appl. Math. Comput. **217** (2010), 3144–3154.
- [16] S. LI, S. STEVIĆ, *Weighted differentiation composition operators from the logarithmic Bloch space to the weighted-type space*, An. Stiint. Univ. “Ovidius” Constanta Ser. Mat. **24** (3) (2016), 223–240.
- [17] Y. LIU, Y. YU, *On a Stević-Sharma operator from Hardy spaces to the logarithmic Bloch spaces*, J. Inequal. Appl. **2015**, Article no. 22, (2015).
- [18] K. MADIGAN, A. MATHESON, *Compact composition operators on the Bloch space*, Trans. Amer. Math. Soc. **347** (7) (1995), 2679–2687.
- [19] S. OHNO, *Products of composition and differentiation on Bloch spaces*, Bull. Korean Math. Soc. **46** (6) (2009), 1135–1140.
- [20] C. PAN, *On an integral-type operator from $Q_k(p, q)$ spaces to α -Bloch spaces*, Filomat **25** (3) (2011), 163–173.
- [21] L. A. RUBEL, A. L. SHIELDS, *The second duals of certain spaces of analytic functions*, J. Austral. Math. Soc. **11** (1970), 276–280.
- [22] W. RUDIN, *Function Theory in Polydiscs*, New York, Amsterdam, WA Benjamin, 1969.
- [23] W. RUDIN, *Function Theory in the Unit Ball of \mathbb{C}^n* , Grundlehren der Mathematischen Wissenschaften, New York, Berlin, Springer-Verlag, 1980.
- [24] W. RUDIN, *Functional Analysis*, 2nd Ed., McGraw-Hill, Inc., New York, 1991.
- [25] H. J. SCHWARTZ, *Composition operators on H^p* , Thesis, University of Toledo, 1969.
- [26] B. SEHBA, S. STEVIĆ, *On some product-type operators from Hardy-Orlicz and Bergman-Orlicz spaces to weighted-type spaces*, Appl. Math. Comput. **233C** (2014), 565–581.
- [27] A. K. SHARMA, *Products of multiplication, composition and differentiation between weighted Bergman-Nevanlinna and Bloch-type spaces*, Turkish. J. Math. **35** (2011), 275–291.
- [28] S. STEVIĆ, *Products of integral-type operators and composition operators from the mixed norm space to Bloch-type spaces*, Siberian Math. J. **50** (4) (2009), 726–736.
- [29] S. STEVIĆ, *Weighted differentiation composition operators from mixed-norm spaces to weighted-type spaces*, Appl. Math. Comput. **211** (2009), 222–233.
- [30] S. STEVIĆ, *Composition followed by differentiation from H^∞ and the Bloch space to n th weighted-type spaces on the unit disk*, Appl. Math. Comput. **216** (2010), 3450–3458.
- [31] S. STEVIĆ, *On operator P_{ϕ}^g from the logarithmic Bloch-type space to the mixed-norm space on unit ball*, Appl. Math. Comput. **215** (2010), 4248–4255.
- [32] S. STEVIĆ, *Weighted differentiation composition operators from H^∞ and Bloch spaces to n th weighted-type spaces on the unit disk*, Appl. Math. Comput. **216** (2010), 3634–3641.
- [33] S. STEVIĆ, *Weighted differentiation composition operators from the mixed-norm space to the n th weighted-type space on the unit disk*, Abstr. Appl. Anal. Vol. 2010, Article ID 246287, (2010), 15 pages.
- [34] S. STEVIĆ, *Weighted iterated radial composition operators between some spaces of holomorphic functions on the unit ball*, Abstr. Appl. Anal. Vol. 2010, Article ID 801264, (2010), 14 pages.
- [35] S. STEVIĆ, *Weighted iterated radial operators between different weighted Bergman spaces on the unit ball*, Appl. Math. Comput. **218** (2012), 8288–8294.
- [36] S. STEVIĆ, *On a new product-type operator on the unit ball*, J. Math. Inequal. **16** (4) (2022), 1675–1692.
- [37] S. STEVIĆ, C. S. HUANG, Z. J. JIANG, *Sum of some product-type operators from Hardy spaces to weighted-type spaces on the unit ball*, Math. Methods Appl. Sci. **45** (17) (2022), 11581–11600.

- [38] S. STEVIĆ, Z. J. JIANG, *Weighted iterated radial composition operators from weighted Bergman-Orlicz spaces to weighted-type spaces on the unit ball*, *Math. Methods Appl. Sci.* **44** (11) (2021), 8684–8696.
- [39] S. STEVIĆ, Z. J. JIANG, *Weighted iterated radial composition operators from logarithmic Bloch spaces to weighted-type spaces on the unit ball*, *Math. Methods Appl. Sci.* **45** (2022), 3083–3097.
- [40] S. STEVIĆ, A. K. SHARMA, A. BHAT, *Products of multiplication composition and differentiation operators on weighted Bergman spaces*, *Appl. Math. Comput.* **217** (2011), 8115–8125.
- [41] S. STEVIĆ, A. K. SHARMA, A. BHAT, *Essential norm of products of multiplication composition and differentiation operators on weighted Bergman spaces*, *Appl. Math. Comput.* **218** (2011), 2386–2397.
- [42] S. STEVIĆ, A. K. SHARMA, R. KRISHAN, *Boundedness and compactness of a new product-type operator from a general space to Bloch-type spaces*, *J. Inequal. Appl.* Vol. 2016, Article No. 219, (2016), 32 pages.
- [43] S. STEVIĆ, S. I. UEKI, *Integral-type operators acting between weighted-type spaces on the unit ball*, *Appl. Math. Comput.* **215** (2009), 2464–2471.
- [44] W. YANG, *Generalized weighted composition operators from the $F(p, q, s)$ space to the Bloch-type space*, *Appl. Math. Comput.* **218** (2012), 4967–4972.
- [45] W. YANG, W. YAN, *Generalized weighted composition operators from area Nevanlinna spaces to weighted-type spaces*, *Bull. Korean Math. Soc.* **48** (6) (2011), 1195–1205.
- [46] W. YANG, X. ZHU, *Generalized weighted composition operators from area Nevanlinna spaces to Bloch-type spaces*, *Taiwan. J. Math.* **16** (3) (2012), 869–883.
- [47] F. ZHANG, Y. LIU, *On the compactness of the Stević-Sharma operator on the logarithmic Bloch spaces*, *Math. Inequal. Appl.* **19** (2) (2016), 625–642.
- [48] K. ZHU, *Spaces of Holomorphic Functions in the Unit Ball*, Graduate Texts in Mathematics, Springer, New York, 2005.
- [49] X. ZHU, *Generalized weighted composition operators on weighted Bergman spaces*, *Numer. Funct. Anal. Opt.* **30** (2009), 881–893.
- [50] X. ZHU, *Generalized weighted composition operators from Bloch spaces into Bers-type spaces*, *Filomat* **26** (2012), 1163–1169.
- [51] X. ZHU, *Generalized weighted composition operators from Bers-type spaces to Bloch-type spaces*, *Math. Ineq. Appl.* **17** (2014), 187–195.
- [52] X. ZHU, *A new characterization of the generalized weighted composition operator from H^∞ into the Zygmund space*, *Math. Inequal. Appl.* **18** (3) (2015), 1135–1142.
- [53] X. ZHU, *Generalized weighted composition operators on Bloch-type spaces*, *J. Ineq. Appl.* **2015** (2015), 59–68.
- [54] X. ZHU, *Essential norm of generalized weighted composition operators on Bloch-type spaces*, *Appl. Math. Comput.* **274** (2016), 133–142.
- [55] X. ZHU, *Generalized weighted composition operators on weighted Bergman spaces, II*, *Math. Ineq. Appl.* **22** (2019), 1055–1066.

(Received November 23, 2022)

Stevo Stević
 Mathematical Institute of the Serbian Academy of Sciences
 Knez Mihailova 36/III, 11000 Beograd, Serbia
 and
 Department of Medical Research
 China Medical University Hospital, China Medical University
 Taichung 40402, Taiwan, Republic of China
 e-mail: sscitel@gmail.com
 sstevic@ptt.rs

Sei-ichiro Ueki
 Department of Mathematical Science
 Faculty of Engineering, Yokohama National University
 Hodogaya, Yokohama, 240–8501 Japan
 e-mail: ueki-seiichiro-zg@ynu.ac.jp