

ABSOLUTE MONOTONICITY OF FOUR FUNCTIONS INVOLVING THE SECOND KIND OF COMPLETE ELLIPTIC INTEGRALS

FEI WANG AND FENG QI*

(Communicated by G. Nemes)

Abstract. In the study, the authors present absolute monotonicity of four functions involving the inverse hyperbolic tangent function and the second kind of complete elliptic integrals, derive four double inequalities for bounding the second kind of complete elliptic integrals, and acquire an upper bound of the Hersch–Pfluger distortion function. These inequalities improve several known ones. Moreover, the authors connect two of the four functions with normalized remainders of the Maclaurin series of two functions involving the inverse hyperbolic tangent function and the second kind of complete elliptic integrals.

1. Introduction and main results

For given complex numbers $a, b, c \in \mathbb{C}$ with $c \neq 0, -1, -2, \dots$, the Gauss hypergeometric function ${}_2F_1$ is defined by

$${}_2F_1(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!}, \quad |z| < 1, \quad (1)$$

where $(z)_n$ for $z \in \mathbb{C}$ is the Pochhammer symbol, also known as the rising factorial and the shifted factorial, which is defined by

$$(z)_n = \prod_{k=0}^{n-1} (z+k) = \frac{\Gamma(n+z)}{\Gamma(z)} = \begin{cases} z(z+1) \cdots (z+n-1), & n \in \mathbb{N}; \\ 1, & n = 0, \end{cases} \quad (2)$$

where

$$\Gamma(z) = \lim_{n \rightarrow \infty} \frac{n! n^z}{\prod_{k=0}^n (z+k)}, \quad z \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$$

Mathematics subject classification (2020): Primary 33E05; Secondary 26A48, 26A51, 26D05, 26D07, 26D15, 33B10, 33C05, 33C75, 41A80.

Keywords and phrases: Absolute monotonicity, inverse hyperbolic tangent function, complete elliptic integral, inequality, Hersch–Pfluger distortion function, normalized remainder, Gauss hypergeometric function.

The first author was partially supported by the Visiting Scholar Foundation of Zhejiang Higher Education (Grant No. FX2023103), by the Project for Combination of Education and Research Training at Zhejiang Polytechnic University of Mechanical and Electrical Engineering (Grant No. A027123212), and by the Project for Technology Innovation Team of Zhejiang Polytechnic University of Mechanical and Electrical Engineering (Grant No. A027421008). The second author was partially supported by the Youth Project of Hulunbuir City for Basic Research and Applied Basic Research (Grant No. GH2024020) and by the Natural Science Foundation of Inner Mongolia Autonomous Region of China (Grant No. 2025QN01041).

* Corresponding author.

is the classical Euler gamma function, whose reciprocal $\frac{1}{\Gamma(z)}$ is an entire function. For more information on the Gauss hypergeometric functions ${}_2F_1$, please refer to [1, Chapter 15], [6, Chapter 2], and the articles [22, 29, 40].

The complete elliptic integrals are the most important quasi-conformal mappings, they can be represented in terms of the Gauss hypergeometric functions ${}_2F_1$, and they have important applications in the theory of quasi-conformal mappings, the theory of geometric functions, and engineering. For more information, please refer to [1, 6, 51, 53] and closely related references therein.

For given $r \in (0, 1)$, the complete elliptic integrals of the first and second kinds $\mathcal{K}(r)$ and $\mathcal{E}(r)$ can be represented [43, 46] respectively by

$$\begin{cases} \mathcal{K}(r) = \frac{\pi}{2} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; r^2\right); \\ \mathcal{K}(0) = \frac{\pi}{2}; \\ \mathcal{K}(1) = \infty \end{cases}$$

and

$$\begin{cases} \mathcal{E}(r) = \frac{\pi}{2} {}_2F_1\left(-\frac{1}{2}, \frac{1}{2}; 1; r^2\right); \\ \mathcal{E}(0) = \frac{\pi}{2}; \\ \mathcal{E}(1) = 1. \end{cases} \quad (3)$$

For more information on some properties and applications of $\mathcal{K}(r)$ and $\mathcal{E}(r)$, please refer to [2, 3, 4, 5, 10, 27, 41, 44, 45, 52, 55, 56, 60, 61, 62, 64, 65, 70] and closely related references therein.

For $r \in (0, 1)$, the conformal modular function of the Grötzsch extremum ring $\mathbb{B}^2 \setminus [0, r]$ can be represented by

$$\mu(r) = \frac{\pi}{2} \frac{\mathcal{K}(\sqrt{1-r^2})}{\mathcal{K}(r)}. \quad (4)$$

The classical Ramanujan modular equation is defined by

$$\frac{\mathcal{K}(\sqrt{1-s^2})}{\mathcal{K}(s)} = p \frac{\mathcal{K}(\sqrt{1-r^2})}{\mathcal{K}(r)}, \quad p > 0. \quad (5)$$

By virtue of (4), we can rewritten (5) as

$$\mu(s) = p\mu(r), \quad p > 0. \quad (6)$$

Accordingly, the solution to the Ramanujan modular equation (6) is

$$s = \varphi_{\kappa}(r) = \mu^{-1}(p\mu(r)), \quad p = \frac{1}{\kappa}$$

and we call $\varphi_\kappa(r)$ the Hersch–Pfluger distortion function. It is well known that the Hersch–Pfluger distortion function $\varphi_\kappa(r)$ has important applications in quasi-conformal Schwarz lemma; see [21].

In 1970, Hübner [20] obtained the inequality

$$\varphi_\kappa(r) < r^{1/\kappa} \exp \left\{ \left(1 - \frac{1}{\kappa} \right) [m(r) + \ln r] \right\}$$

for $r \in (0, 1)$ and $\kappa \in (1, \infty)$, where

$$m(r) = \frac{2}{\pi} (1 - r^2) \mathcal{K}(r) \mathcal{K}(\sqrt{1 - r^2})$$

and we call $m(r) + \ln r$ the Hübner function. In 2008, Wang and her coauthors [58] established the inequality

$$\varphi_\kappa(r) < r^{1/\kappa} \exp \left\{ \frac{4 \ln 2}{\pi - 2} \left(1 - \frac{1}{\kappa} \right) [\mathcal{E}(r) - 1] \right\}, \quad r \in (0, 1). \quad (7)$$

In 2011, Guo and Qi [13] acquired the double inequality

$$\frac{\pi}{2} - \frac{1}{2} \ln \frac{(1+r)^{r-1}}{(1-r)^{r+1}} < \mathcal{E}(r) < \frac{\pi-1}{2} + \frac{1-r^2}{4r} \ln \frac{1+r}{1-r}, \quad r \in (0, 1). \quad (8)$$

In [54], Wang and his coauthors obtained two double inequalities

$$\frac{\pi}{2} - \alpha_1 \left[1 - (1-r^2) \frac{\operatorname{arctanh} r}{r} \right] < \mathcal{E}(r) < \frac{\pi}{2} - \beta_1 \left[1 - (1-r^2) \frac{\operatorname{arctanh} r}{r} \right] \quad (9)$$

and

$$\frac{\pi}{2} - r \operatorname{arctanh} r - \frac{\lambda_2}{2} \ln(1-r^2) < \mathcal{E}(r) < \frac{\pi}{2} - r \operatorname{arctanh} r - \frac{\mu_2}{2} \ln(1-r^2) \quad (10)$$

for $r \in (0, 1)$, where

$$\begin{aligned} \alpha_1 &= \frac{3\pi}{16} = 0.589\ldots, & \beta_1 &= \frac{\pi}{2} - 1 = 0.570\ldots, \\ \lambda_2 &= 1, & \mu_2 &= 2 - \frac{\pi}{4} = 1.214\ldots \end{aligned}$$

are the best possible constants. For more information on this kind of elementary inequalities, please refer to the articles [39, 41, 63] and closely related references therein.

Let $I \subseteq \mathbb{R}$ be an infinite or finite interval. A real infinitely differentiable function $f(x)$ defined on I is said to be absolutely monotonic in $x \in I$ if and only if all of its derivatives satisfy $f^{(k)}(x) \geq 0$ for $k \in \mathbb{N}_0$ and $x \in I$. A real infinitely differentiable function $f(x)$ defined on I is said to be completely monotonic in $x \in I$ if and only if all of its derivatives satisfy $(-1)^k f^{(k)}(x) \geq 0$ for $k \in \mathbb{N}_0$ and $x \in I$. When $I = (0, \infty)$ or $I = [0, \infty)$, there have been plenty of classical investigations on absolutely (or

completely, respectively) monotonic functions in [30, Chapter XIII], [59, Chapter IV], and the monograph [48]. By the way, in the papers [8, 12, 38], the authors invented the notions of logarithmically absolutely (or logarithmically completely, respectively) monotonic functions.

In the study, we will present absolute monotonicity of four functions involving the inverse hyperbolic tangent function $\operatorname{arctanh} r$ and the second kind of complete elliptic integrals $\mathcal{E}(r)$, derive four double inequalities for bounding the second kind of complete elliptic integrals $\mathcal{E}(r)$, and acquire an upper bound of the Hersch–Pfluger distortion function $\varphi_\kappa(r)$; see (45). These inequalities improve (7), (8), (9), and (10). Moreover, we will connect two of the four functions with normalized remainders of the Maclaurin series of two functions involving the inverse hyperbolic tangent function $\operatorname{arctanh} r$ and the second kind of complete elliptic integrals $\mathcal{E}(r)$; see Remarks 2 and 4 below.

The main results of this paper can be stated in the following theorems.

THEOREM 1. For $r \in (0, 1)$ and

$$a_0 = \frac{3}{8}, \quad a_1 = -\frac{3}{640}, \quad a_2 = -\frac{171}{89600}, \quad a_3 = -\frac{30161}{28672000}, \quad (11)$$

define

$$H(r) = \frac{1 - {}_2F_1\left(-\frac{1}{2}, \frac{1}{2}; 1; r\right)}{1 - (1-r){}_2F_1\left(\frac{1}{2}, 1; \frac{3}{2}; r\right)} = \frac{1 - \frac{2}{\pi}\mathcal{E}(\sqrt{r})}{1 + (r-1)\frac{\operatorname{arctanh}\sqrt{r}}{\sqrt{r}}} = \sum_{n=0}^{\infty} a_n r^n.$$

Then the following conclusions are true:

1. The limits $H(0^+) = \frac{3}{8}$ and $H(1^-) = 1 - \frac{2}{\pi}$ are valid.
2. For $n \in \mathbb{N}$, the coefficients a_n satisfy the recursive relation and the negativity

$$a_n = 3 \left\{ \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}}{4[(n+1)!]^2} - \sum_{k=0}^{n-1} \frac{a_k}{[2(n-k)+1][2(n-k)+3]} \right\} < 0. \quad (12)$$

3. The functions $-H'(r)$ and $H_1(r) = \frac{1}{r} \left[\frac{3}{8} - H(r) \right]$ are absolutely monotonic on $(0, 1)$, with the limits $H_1(0^+) = \frac{3}{640}$ and $H_1(1^-) = \frac{2}{\pi} - \frac{5}{8}$.
4. For $r \in (0, 1)$,

$$\begin{aligned} \mathcal{E}(r) = \frac{\pi}{2} - \frac{\pi}{2} \left[1 - (1-r^2) \frac{\operatorname{arctanh} r}{r} \right] \\ \times \left(\frac{3}{8} - \frac{3}{640} r^2 - \frac{171}{89600} r^4 - \frac{30161}{28672000} r^6 - \dots \right). \quad (13) \end{aligned}$$

THEOREM 2. For $r \in (0, 1)$ and

$$b_0 = 1 - \frac{\pi}{8}, \quad b_1 = \frac{15\pi - 64}{384}, \quad b_2 = \frac{95\pi - 384}{7680}, \quad b_3 = \frac{69195\pi - 274432}{1032190}, \quad (14)$$

define

$$\begin{aligned} G(r) &= \frac{\frac{\pi}{2} [{}_2F_1(-\frac{1}{2}, \frac{1}{2}; 1; r) - 1] + r {}_2F_1(\frac{1}{2}, 1; \frac{3}{2}; r)}{r {}_2F_1(1, 1; 2; r)} \\ &= \frac{\frac{\pi}{2} - \sqrt{r} \operatorname{arctanh} \sqrt{r} - \mathcal{E}(\sqrt{r})}{\ln(1-r)} \\ &= \sum_{n=0}^{\infty} b_n r^n. \end{aligned}$$

Then the following conclusions are true:

1. The limits $G(0^+) = 1 - \frac{\pi}{8}$ and $G(1^-) = \frac{1}{2}$ are valid.
2. For $n \in \mathbb{N}$, the coefficients b_n satisfy the recursive relation and the negativity

$$b_n = \frac{\pi}{2} \frac{(-\frac{1}{2})_{n+1} (\frac{1}{2})_{n+1}}{[(n+1)!]^2} + \frac{1}{2n+1} - \sum_{k=0}^{n-1} \frac{b_k}{n-k+1} < 0. \quad (15)$$

3. The functions $-G'(r)$ and $G_1(r) = \frac{1}{r} [1 - \frac{\pi}{8} - G(r)]$ are absolutely monotonic on $(0, 1)$, with the limits $G_1(0^+) = \frac{64-15\pi}{384}$ and $G_1(1^-) = \frac{4-\pi}{8}$.
4. For $r \in (0, 1)$,

$$\begin{aligned} \mathcal{E}(r) &= \frac{\pi}{2} - r \operatorname{arctanh} r - \left(1 - \frac{\pi}{8} + \frac{15\pi - 64}{384} r^2 \right. \\ &\quad \left. + \frac{95\pi - 384}{7680} r^4 + \frac{69195\pi - 274432}{1032190} r^6 + \dots \right) \ln(1-r^2). \quad (16) \end{aligned}$$

We will prove these main conclusions in Section 3 below.

2. A lemma

In the proofs of our main results, we will need the formulas

$${}_2F_1(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}, \quad c \neq 0, -1, -2, \dots, \quad \Re(c-a-b) > 0 \quad (17)$$

and

$$z^{b-a} \frac{\Gamma(z+a)}{\Gamma(z+b)} \sim 1 + \frac{(a-b)(a+b-1)}{2z} + \dots, \quad z \rightarrow \infty. \quad (18)$$

These two formulas can be found in [1, pp. 257 and 556, Entries 6.1.47 and 15.1.20] and [49, pp. 66–68, Section 3.6.2], respectively.

LEMMA 1. For $n \in \mathbb{N}$, the sequence

$$C_n = \frac{(2n+1)(2n+3)(4n+5)}{n+2} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}}{[(n+1)!]^2}$$

is increasing in $n \in \mathbb{N}$, with the limit $\lim_{n \rightarrow \infty} C_n = \frac{16}{\pi}$. Consequently, for $n \in \mathbb{N}$,

$$\frac{135}{32} \frac{n+2}{(2n+1)(2n+3)(4n+5)} \leq \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}}{[(n+1)!]^2} < \frac{16}{\pi} \frac{n+2}{(2n+1)(2n+3)(4n+5)}.$$

Proof. Due to $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$ and the asymptotic formula (18), we obtain the limit

$$\begin{aligned} \lim_{n \rightarrow \infty} C_n &= \lim_{n \rightarrow \infty} \frac{(2n+1)(2n+3)(4n+5) \left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}}{(n+2)[(n+1)!]^2} \\ &= \lim_{n \rightarrow \infty} \frac{(2n+1)(2n+3)(4n+5) \Gamma\left(n+\frac{1}{2}\right) \Gamma\left(n+\frac{3}{2}\right)}{(n+2) \left[\Gamma\left(\frac{1}{2}\right)\right]^2 [\Gamma(n+2)]^2} \\ &= \lim_{n \rightarrow \infty} \frac{(2n+1)(2n+3)(4n+5)}{(n+2) \left[\Gamma\left(\frac{1}{2}\right)\right]^2 n^2} \\ &= \frac{16}{\pi}. \end{aligned}$$

By virtue of the definition in (2), we arrive at

$$\begin{aligned} \frac{C_{n+1}}{C_n} &= \frac{(2n+3)(2n+5)(4n+9) \left(\frac{1}{2}\right)_{n+1} \left(\frac{1}{2}\right)_{n+2}}{(2n+1)(2n+3)(4n+5) \left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}} \frac{(n+2)[(n+1)!]^2}{(n+3)[(n+2)!]^2} \\ &= \frac{(2n+5)(2n+3)(4n+9)}{4(4n+5)(n+3)(n+2)} \\ &= \frac{16n^3 + 100n^2 + 204n + 135}{16n^3 + 100n^2 + 196n + 120} \\ &= 1 + \frac{8n + 15}{16n^3 + 100n^2 + 196n + 120} \\ &> 1. \end{aligned}$$

Accordingly, the sequence C_n is increasing in $n \in \mathbb{N}$.

In light of the increasing property of the sequence C_n in $n \in \mathbb{N}$ and in view of the limit $\lim_{n \rightarrow \infty} C_n = \frac{16}{\pi}$, we acquire $\frac{135}{32} = C_1 \leq C_n < \frac{16}{\pi}$. The proof of Lemma 1 is thus complete. \square

3. Proofs of main results

We are now in a position to prove Theorems 1 and 2.

3.1. Proof of Theorem 1

Let

$$h_1(r) = 1 - {}_2F_1\left(-\frac{1}{2}, \frac{1}{2}; 1; r\right)$$

and

$$h_2(r) = 1 - (1-r) {}_2F_1\left(\frac{1}{2}, 1; \frac{3}{2}; r\right).$$

Then $H(r) = \frac{h_1(r)}{h_2(r)} = \sum_{n=0}^{\infty} a_n r^n$. Making use of the definition in (1), we derive

$$h_1(r) = 1 - \sum_{n=0}^{\infty} \frac{\left(-\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_n}{[(n)!]^2} r^n = \frac{1}{2} \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}}{[(n+1)!]^2} r^{n+1}$$

and

$$h_2(r) = 1 - (1-r) \sum_{n=0}^{\infty} \frac{1}{2n+1} r^n = \sum_{n=0}^{\infty} \frac{2}{(2n+1)(2n+3)} r^{n+1}.$$

Accordingly, we deduce

$$H(r) = \frac{1}{4} \frac{\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}}{[(n+1)!]^2} r^n}{\sum_{n=0}^{\infty} \frac{1}{(2n+1)(2n+3)} r^n} = \sum_{n=0}^{\infty} a_n r^n,$$

which can be reformulated as

$$\frac{1}{4} \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}}{[(n+1)!]^2} r^n = \sum_{n=0}^{\infty} \frac{1}{(2n+1)(2n+3)} r^n \sum_{n=0}^{\infty} a_n r^n.$$

As a result, we arrive at

$$\begin{aligned} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}}{4[(n+1)!]^2} &= \sum_{k=0}^n \frac{a_k}{[2(n-k)+1][2(n-k)+3]} \\ &= \frac{a_n}{3} + \sum_{k=0}^{n-1} \frac{a_k}{[2(n-k)+1][2(n-k)+3]}, \end{aligned} \quad (19)$$

which is equivalent to the recursive relation in (12).

Let

$$A_n^{(1)} = \frac{(2n+1)(2n+3)}{4} \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}}{[(n+1)!]^2}, \quad n \in \mathbb{N}.$$

Then, by virtue of the definition in (2), we find

$$\frac{A_{n+1}^{(1)}}{A_n^{(1)}} = \frac{(2n+3)(2n+5)}{4(n+2)^2} = \frac{4n^2+16n+15}{4n^2+16n+16} < 1, \quad n \in \mathbb{N}.$$

Hence, the sequence $A_n^{(1)}$ decreases in $n \in \mathbb{N}$.

On the other hand, let

$$A_n^{(2)} = (2n+1)(2n+3) \sum_{k=0}^n \frac{a_k}{[2(n-k)+1][2(n-k)+3]}, \quad n \in \mathbb{N}.$$

Due to the last equation in (19), we see that $A_n^{(1)} = A_n^{(2)}$ for $n \in \mathbb{N}$. As a result, the sequence $A_n^{(2)}$ decreases in $n \in \mathbb{N}$. Furthermore, we have

$$\begin{aligned} \frac{A_{n+1}^{(2)} - A_n^{(2)}}{2n+3} &= (2n+5) \sum_{k=1}^{n+1} \frac{a_k}{[2(n-k)+3][2(n-k)+5]} \\ &\quad - (2n+1) \sum_{k=1}^n \frac{a_k}{[2(n-k)+1][2(n-k)+3]} \\ &= \frac{(2n+5)a_{n+1}}{3} + \sum_{k=1}^n \left(\frac{(2n+5)a_k}{[2(n-k)+3][2(n-k)+5]} \right. \\ &\quad \left. - \frac{(2n+1)a_k}{[2(n-k)+1][2(n-k)+3]} \right) \\ &= \frac{(2n+5)a_{n+1}}{3} - 8 \sum_{k=1}^n \frac{ka_k}{[2(n-k)+1][2(n-k)+3][2(n-k)+5]} \\ &< 0. \end{aligned}$$

Consequently, we conclude

$$a_{n+1} < \frac{24}{2n+5} \sum_{k=1}^n \frac{ka_k}{[2(n-k)+1][2(n-k)+3][2(n-k)+5]}, \quad n \in \mathbb{N}. \quad (20)$$

In view of the recursive relation in (12), it is easy to deduce the four values in (11). By induction and with the aid of the inequality (20), it is not difficult to prove that $a_n < 0$ for $n \in \mathbb{N}$.

It is clear that $H(0^+) = a_0 = \frac{3}{8}$. In light of the formula (17), we acquire

$$\begin{aligned} \lim_{r \rightarrow 1^-} H(r) &= \lim_{r \rightarrow 1^-} \frac{1 - {}_2F_1\left(-\frac{1}{2}, \frac{1}{2}; 1; r\right)}{1 - (1-r) {}_2F_1\left(\frac{1}{2}, 1; \frac{3}{2}; r\right)} \\ &= 1 - {}_2F_1\left(-\frac{1}{2}, \frac{1}{2}; 1; 1\right) \\ &= 1 - \frac{[\Gamma(1)]^2}{\Gamma\left(\frac{3}{2}\right)\Gamma\left(\frac{1}{2}\right)} \\ &= 1 - \frac{2}{\pi}. \end{aligned}$$

Since $a_n < 0$ for $n \in \mathbb{N}$, considering the series expansion $H(r) = \sum_{n=0}^{\infty} a_n r^n$, we obtain

$$-H'(r) = -\sum_{n=0}^{\infty} (n+1)a_{n+1}r^n = \sum_{n=0}^{\infty} |(n+1)a_{n+1}|r^n$$

and

$$H_1(r) = \frac{1}{r} \left[\frac{3}{8} - H(r) \right] = -\sum_{n=0}^{\infty} a_{n+1}r^n = \sum_{n=0}^{\infty} |a_{n+1}|r^n.$$

Consequently, the functions $-H'(r)$ and $H_1(r)$ are absolutely monotonic on $(0, 1)$.

It is ready that $H_1(0^+) = |a_1| = \frac{3}{640}$ and $H_1(1^-) = \frac{3}{8} - H(1^-) = \frac{2}{\pi} - \frac{5}{8}$.

In [11, p. 61], [34, p. 473, Eq. 83], and [49, p. 109], we find

$$\frac{\operatorname{arctanh} r}{r} = {}_2F_1\left(\frac{1}{2}, 1; \frac{3}{2}; r^2\right).$$

Hence, we have

$$1 - (1-r^2) \frac{\operatorname{arctanh} r}{r} = 1 - (1-r^2) {}_2F_1\left(\frac{1}{2}, 1; \frac{3}{2}; r^2\right). \quad (21)$$

Accordingly, we derive

$$H(r^2) = \frac{1 - \frac{2}{\pi} \mathcal{E}(r)}{1 - (1-r^2) \frac{\operatorname{arctanh} r}{r}} = \sum_{n=0}^{\infty} a_n r^{2n}.$$

Combining (12) and (21) yields (13). The proof of Theorem 1 is thus complete. \square

3.2. Proof of Theorem 2

Let

$$g_1(r) = \frac{\pi}{2} \left[{}_2F_1\left(-\frac{1}{2}, \frac{1}{2}; 1; r\right) - 1 \right] + r {}_2F_1\left(\frac{1}{2}, 1; \frac{3}{2}; r\right)$$

and

$$g_2(r) = r {}_2F_1(1, 1; 2; r).$$

Then $G(r) = \frac{g_1(r)}{g_2(r)} = \sum_{n=0}^{\infty} b_n r^n$. By the definition in (1), we have

$$g_1(r) = \sum_{n=0}^{\infty} \left[\frac{\pi}{2} \frac{\left(-\frac{1}{2}\right)_{n+1} \left(\frac{1}{2}\right)_{n+1}}{[(n+1)!]^2} + \frac{1}{2n+1} \right] r^{n+1}$$

and

$$g_2(r) = r {}_2F_1(1, 1; 2; r) = \sum_{n=0}^{\infty} \frac{1}{n+1} r^{n+1}.$$

Accordingly, we obtain

$$G(r) = \frac{\sum_{n=0}^{\infty} \left[\frac{\pi}{2} \frac{\left(-\frac{1}{2}\right)_{n+1} \left(\frac{1}{2}\right)_{n+1}}{[(n+1)!]^2} + \frac{1}{2n+1} \right] r^{n+1}}{\sum_{n=0}^{\infty} \frac{1}{n+1} r^{n+1}} = \sum_{n=0}^{\infty} b_n r^n,$$

that is,

$$\sum_{n=0}^{\infty} \left[\frac{\pi \left(-\frac{1}{2}\right)_{n+1} \left(\frac{1}{2}\right)_{n+1}}{2 [(n+1)!]^2} + \frac{1}{2n+1} \right] r^n = \sum_{n=0}^{\infty} \frac{1}{n+1} r^n \sum_{n=0}^{\infty} b_n r^n. \quad (22)$$

In other words,

$$\frac{\pi \left(-\frac{1}{2}\right)_{n+1} \left(\frac{1}{2}\right)_{n+1}}{2 [(n+1)!]^2} + \frac{1}{2n+1} = \sum_{k=0}^n \frac{b_k}{n-k+1} = b_n + \sum_{k=0}^{n-1} \frac{b_k}{n-k+1}. \quad (23)$$

As a result, we derive the recursive relation in (15).

Set

$$B_n^{(1)} = (n+1) \left[\frac{\pi \left(-\frac{1}{2}\right)_{n+1} \left(\frac{1}{2}\right)_{n+1}}{2 [(n+1)!]^2} + \frac{1}{2n+1} \right].$$

Making use of Lemma 1, we reveal

$$\begin{aligned} B_{n+1}^{(1)} - B_n^{(1)} &= \frac{\pi}{2} \left[\frac{(n+2) \left(-1/2, n+2\right) \left(\frac{1}{2}\right)_{n+2}}{[(n+2)!]^2} - \frac{(n+1) \left(-\frac{1}{2}\right)_{n+1} \left(\frac{1}{2}\right)_{n+1}}{[(n+1)!]^2} \right] \\ &\quad - \frac{1}{(2n+1)(2n+3)} \\ &= \frac{\pi}{2} \frac{\left(-\frac{1}{2}\right)_{n+1} \left(\frac{1}{2}\right)_{n+1}}{[(n+1)!]^2} \left[\frac{(2n+1)(2n+3)}{4(n+2)} - (n+1) \right] \\ &\quad - \frac{1}{(2n+1)(2n+3)} \\ &= \frac{\pi}{16} \frac{(4n+5) \left(\frac{1}{2}\right)_n \left(\frac{1}{2}\right)_{n+1}}{(n+2)[(n+1)!]^2} - \frac{1}{(2n+1)(2n+3)} \\ &< 0. \end{aligned}$$

This means that $B_n^{(1)}$ is decreasing in $n \in \mathbb{N}$.

On the other hand, let

$$B_n^{(2)} = (n+1) \sum_{k=0}^n \frac{b_k}{n-k+1}.$$

By the equations in (23), we see easily that $B_n^{(1)} = B_n^{(2)}$. Hence, the sequence $B_n^{(2)}$ is decreasing in $n \in \mathbb{N}$. As a result, we obtain

$$\begin{aligned} B_{n+1}^{(2)} - B_n^{(2)} &= (n+2) \sum_{k=0}^{n+1} \frac{b_k}{n-k+2} - (n+1) \sum_{k=0}^n \frac{b_k}{n-k+1} \\ &= (n+2)b_{n+1} + \sum_{k=1}^n \left[\frac{(n+2)b_k}{n-k+2} - \frac{(n+1)b_k}{n-k+1} \right] \\ &= (n+2)b_{n+1} - \sum_{k=1}^n \frac{kb_k}{(n-k+1)(n-k+2)} \\ &< 0. \end{aligned}$$

Consequently, we have

$$b_{n+1} < \frac{1}{n+2} \sum_{k=1}^n \frac{kb_k}{(n-k+1)(n-k+2)}. \quad (24)$$

By the equation (22), we have $b_0 = 1 - \frac{\pi}{8}$. With the help of the recursive relation in (15), we derive the rest values in (14). By induction and (24), it is not difficult to verify that $b_n < 0$ for $n \in \mathbb{N}$.

By the L'Hôpital rule and $\frac{d\mathcal{E}(r)}{dr} = \frac{\mathcal{E}(r) - \mathcal{K}(r)}{r}$, we acquire the limit

$$\begin{aligned} \lim_{r \rightarrow 1^-} G(r) &= \lim_{r \rightarrow 1^-} \frac{\frac{\pi}{2} [{}_2F_1(-\frac{1}{2}, \frac{1}{2}; 1; r) - 1] + r {}_2F_1(\frac{1}{2}, 1; \frac{3}{2}; r)}{r {}_2F_1(1, 1; 2; r)} \\ &= \lim_{r \rightarrow 1^-} \frac{(r-1)[\mathcal{K}(\sqrt{r}) - \mathcal{E}(\sqrt{r})] + (1-r)\sqrt{r} \operatorname{arctanh} \sqrt{r}}{2r} + \frac{1}{2} \\ &= \frac{1}{2}. \end{aligned}$$

Since $b_n < 0$ for $n \in \mathbb{N}$, by the expression $G(r) = \sum_{n=0}^{\infty} b_n r^n$, we arrive at

$$-G'(r) = -\sum_{n=0}^{\infty} (n+1)b_{n+1}r^n = \sum_{n=0}^{\infty} |(n+1)b_{n+1}|r^n$$

and

$$G_1(r) = \frac{1}{r} \left[1 - \frac{\pi}{8} - G(r) \right] = -\sum_{n=0}^{\infty} b_{n+1}r^n = \sum_{n=0}^{\infty} |b_{n+1}|r^n.$$

Accordingly, the functions $-G'(r)$ and $G_1(r)$ are absolutely monotonic on $(0, 1)$. Moreover, it is obvious that the limits

$$G_1(0^+) = |b_1| = \frac{64 - 15\pi}{384}$$

and

$$G_1(1^-) = 1 - \frac{\pi}{8} - G(1^-) = \frac{4 - \pi}{8}$$

are valid.

By the definition in (1), we acquire

$$-\ln(1-r) = \sum_{n=0}^{\infty} \frac{1}{n+1} r^{n+1} = r {}_2F_1(1, 1; 2; r). \quad (25)$$

Combining this with (3) gives

$$G(r^2) = \frac{\mathcal{E}(r) - \frac{\pi}{2} + r \operatorname{arctanh} r}{-\ln(1-r^2)} = \sum_{n=0}^{\infty} b_n r^{2n}.$$

Consequently, we deduce the expansion (16). The proof of Theorem 2 is thus complete. \square

4. Corolaries and remarks

In this section, we deduce several corollaries from Theorems 1 and 2, while we list several remarks on our main results.

COROLLARY 1. *Let the sequence a_n for $n \in \mathbb{N}_0$ be defined as in Theorem 1 and let*

$$\alpha_n = |a_{n+1}|, \quad \beta_n = \frac{2}{\pi} - 1 + \sum_{k=0}^n a_k, \quad P_n(r) = \sum_{k=0}^{n+1} a_k r^{2k}.$$

Then the function

$$H_{2,n}(r) = \frac{1}{r^{n+1}} \left[\sum_{k=0}^n a_k r^k - H(r) \right] \quad (26)$$

is absolutely monotonic from $(0, 1)$ onto $(\alpha_n, \alpha_n + \beta_n)$. Consequently, the double inequalities

$$\begin{aligned} \frac{\pi}{2} \left\{ 1 - \left[1 - (1-r^2) \frac{\operatorname{arctanh} r}{r} \right] [P_n(r) - \alpha_{n+1} r^{2n+4}] \right\} &< \mathcal{E}(r) \\ &< \frac{\pi}{2} \left\{ 1 - \left[1 - (1-r^2) \frac{\operatorname{arctanh} r}{r} \right] [P_n(r) - \beta_{n+1} r^{2n+4}] \right\} \end{aligned} \quad (27)$$

and

$$\begin{aligned} \frac{\pi}{2} \left\{ 1 - \left[1 - (1-r^2) \frac{\operatorname{arctanh} r}{r} \right] \left(\frac{3}{8} - \frac{3}{640} r^2 - \frac{171}{89600} r^4 - \frac{179200 - 56591\pi}{89600\pi} r^6 \right) \right\} \\ > \mathcal{E}(r) > \\ \frac{\pi}{2} \left\{ 1 - \left[1 - (1-r^2) \frac{\operatorname{arctanh} r}{r} \right] \left(\frac{3}{8} - \frac{3}{640} r^2 - \frac{171}{89600} r^4 - \frac{30161}{28672000} r^6 \right) \right\} \end{aligned} \quad (28)$$

are sound for $r \in (0, 1)$.

Proof. Since $H(r) = \sum_{n=0}^{\infty} a_n r^n$, by Theorem 1, we arrive at

$$H_{2,n}(r) = \frac{1}{r^{n+1}} \left(\sum_{k=0}^n a_k r^k - \sum_{k=0}^{\infty} a_k r^k \right) = - \sum_{k=n+1}^{\infty} a_k r^{k-n-1} = \sum_{k=0}^{\infty} |a_{k+n+1}| r^k.$$

Accordingly, the function $H_{2,n}(r)$ is absolutely monotonic on $(0, 1)$.

By Theorem 1, we acquire

$$H_{2,n}(0^+) = |a_{n+1}|, \quad H_{2,n}(1^-) = \sum_{k=0}^n a_k - H(1^-) = \frac{2}{\pi} - 1 + \sum_{k=0}^n a_k.$$

Since $H_{2,n}(r)$ is increasing and convex on $(0, 1)$, we discover

$$\begin{aligned} |a_{n+1}| + |a_{n+2}|r &< H_{2,n}(r) < |a_{n+1}| + \left[\sum_{k=0}^n a_k - H(1^-) - |a_{n+1}| \right] r \\ &= |a_{n+1}| + \beta_{n+1}r. \end{aligned} \quad (29)$$

In view of (4), (21), and (29), we derive the double inequality (27).

Taking $n = 1$ in (27) leads to the double inequality (28). The proof of Corollary 1 is complete. \square

REMARK 1. The double inequality (27) is better than (8) and (9).

REMARK 2. Suppose that a real infinitely differentiable function $f(x)$ has a formal Maclaurin power series expansion

$$\sum_{j=0}^{\infty} f^{(j)}(0) \frac{x^j}{j!}. \quad (30)$$

If $f^{(k+1)}(0) \neq 0$ for some $k \in \mathbb{N}_0$, then we call the function

$$T_k[f(x)] = \begin{cases} \frac{1}{f^{(k+1)}(0)} \frac{(k+1)!}{x^{k+1}} \left[f(x) - \sum_{j=0}^k f^{(j)}(0) \frac{x^j}{j!} \right], & x \neq 0 \\ 1, & x = 0 \end{cases} \quad (31)$$

the normalized remainder, or normalized tail, of the Maclaurin series expansion (30). This concept was invented by the second author and his coworkers starting from April 2023. The first several articles implicitly discussed about the normalized remainders are [23, 24, 25, 37]. The latest articles explicitly considered about the normalized remainders are [7, 22, 26, 31, 32, 35, 50, 57, 66, 67, 68, 69], especially the review and research article [36].

The Stirling numbers of the second kind $S(j, \ell)$ can be analytically generated by

$$\left(\frac{e^z - 1}{z} \right)^\ell = \sum_{j=0}^{\infty} \frac{S(j + \ell, \ell)}{\binom{j + \ell}{\ell}} \frac{z^j}{j!}, \quad \ell \geq 0; \quad (32)$$

see [28, Example 2.76] and [47, Eq. (9.59)]. The r -associate Stirling numbers of the second kind $S_r(j, \ell)$ for $j \geq \ell \geq 0$ and $r \geq 0$ are defined [17, p. 303, Eq. (1.2)] by

$$\left(e^z - \sum_{j=0}^r \frac{z^j}{j!} \right)^\ell = \left(\sum_{j=r+1}^{\infty} \frac{z^j}{j!} \right)^\ell = \ell! \sum_{j=(r+1)\ell}^{\infty} S_r(j, \ell) \frac{z^j}{j!}; \quad (33)$$

see also [42, Section 1.12]. The equation (33) can be reformulated as

$$\begin{aligned} (T_r[e^z])^\ell &= \left[\frac{(r+1)!}{z^{r+1}} \left(e^z - \sum_{j=0}^r \frac{z^j}{j!} \right) \right]^\ell \\ &= \frac{\ell! [(r+1)!]^\ell}{[(r+1)\ell]!} \sum_{j=0}^{\infty} \frac{S_r(j + (r+1)\ell, \ell)}{\binom{j + (r+1)\ell}{\ell}} \frac{z^j}{j!} \end{aligned} \quad (34)$$

for $\ell, r \in \mathbb{N}_0$. The equation (32) is a special case $r = 0$ of (34). Consequently, the integer powers of the normalized remainder $T_r[e^z]$ for $r \in \mathbb{N}_0$, which was investigated

in [7, 35] and [36, Section 1.7], can be regarded as the generating functions of the Stirling numbers of the second kind $S(j, \ell)$ and the r -associate Stirling numbers of the second kind $S_r(j, \ell)$ for $j \geq \ell \geq 0$ and $r \geq 0$.

The Bernoulli numbers and polynomials B_j and $B_j(t)$ for $j \in \mathbb{N}_0 = \{0\} \cup \mathbb{N}$ are generated [49, p. 3] by

$$\frac{z}{e^z - 1} = \sum_{j=0}^{\infty} B_j \frac{z^j}{j!} = 1 - \frac{z}{2} + \sum_{j=1}^{\infty} B_{2j} \frac{z^{2j}}{(2j)!}, \quad |z| < 2\pi \quad (35)$$

and

$$\frac{ze^z}{e^z - 1} = \sum_{j=0}^{\infty} B_j(t) \frac{z^j}{j!}, \quad |z| < 2\pi. \quad (36)$$

In the thesis [16] and the papers [9, 14, 15, 18, 19], among other things, Carlitz and Howard defined the sequences and polynomials A_j , $A_j(t)$, $A_{k,j}$, and $A_{k,j}(t)$ by

$$\frac{z^2}{2} \frac{1}{e^z - z - 1} = \sum_{j=0}^{\infty} A_j \frac{z^j}{j!}, \quad (37)$$

$$\frac{z^2}{2} \frac{e^z}{e^z - z - 1} = \sum_{j=0}^{\infty} A_j(t) \frac{z^j}{j!}, \quad (38)$$

$$\frac{z^k}{k!} \frac{1}{e^z - \sum_{j=0}^{k-1} \frac{z^j}{j!}} = \sum_{j=0}^{\infty} A_{k,j} \frac{z^j}{j!}, \quad (39)$$

and

$$\frac{z^k}{k!} \frac{e^z}{e^z - \sum_{j=0}^{k-1} \frac{z^j}{j!}} = \sum_{j=0}^{\infty} A_{k,j}(t) \frac{z^j}{j!} \quad (40)$$

for $k \in \mathbb{N}$, and they examined a lot of algebraic properties of these sequences and polynomials. It is easy to see that $A_j(0) = A_j$, $A_{1,j} = B_j$, $A_{2,j} = A_j$, $A_{k,j}(0) = A_{k,j}$, $A_{1,j}(0) = B_j$, $A_{2,j}(0) = A_j$, and $A_{1,j}(t) = B_j(t)$ for $j \in \mathbb{N}_0$. The sequence A_j for $j \in \mathbb{N}_0$ was recently investigated in [7, Remark 2] and [42, Section 2]. In terms of the notation $T_{k-1}[e^z]$, the equation (40) can be reformulated as

$$\frac{e^z}{T_{k-1}[e^z]} = \sum_{j=0}^{\infty} A_{k,j}(t) \frac{z^j}{j!}.$$

Meanwhile, the generating functions of the sequences and polynomials B_j , $B_j(t)$, A_j , $A_j(t)$, and $A_{k,j}$ in (35), (36), (37), (38), and (39) can be respectively rewritten as $\frac{1}{T_0[e^z]}$, $\frac{e^z}{T_0[e^z]}$, $\frac{1}{T_1[e^z]}$, $\frac{e^z}{T_1[e^z]}$, and $\frac{1}{T_{k-1}[e^z]}$ for $k \in \mathbb{N}$.

We say that the series $\sum_{j=0}^{\infty} a_j$ envelops the number A if the relations

$$\left| A - \sum_{j=0}^n a_j \right| < |a_{n+1}|, \quad n \in \mathbb{N}_0$$

are satisfied. For more detailed information, please refer to [33, Chapter 4]. The normalized remainder defined in (31) has close connection to the well-known “enveloping series”, a topic covered in [33, Chapter 4].

In terms of the notion of normalized remainders mentioned above, the function $H_{2,n}(r)$ defined in (26) can be reformulated as

$$T_n[H(r)] = -\frac{1}{a_{n+1}}H_{2,n}(r), \quad r \in (0, 1)$$

for $n \in \mathbb{N}_0$. Considering the negativity in (12) and employing Corollary 1, we conclude that the normalized remainder $T_n[H(r)]$ for $n \in \mathbb{N}_0$ is absolutely monotonic from $(0, 1)$ onto $(1, 1 + \frac{\beta_n}{\alpha_n})$.

COROLLARY 2. *Let the sequence b_n for $n \in \mathbb{N}_0$ be defined as in Theorem 2. Set*

$$\delta_n = |b_{n+1}|, \quad \lambda_n = -\frac{1}{2} + \sum_{k=0}^n b_k, \quad Q_n(r) = \sum_{k=0}^{n+1} b_k r^{2k}$$

for $r \in (0, 1)$. Then the function

$$G_{2,n}(r) = \frac{1}{r^{n+1}} \left[\sum_{k=0}^n b_k r^k - G(r) \right] \quad (41)$$

is absolutely monotonic from $(0, 1)$ onto $(\delta_n, \delta_n + \lambda_n)$. Consequently, for $r \in (0, 1)$, the double inequalities

$$\begin{aligned} \frac{\pi}{2} - r \operatorname{arctanh} r - [Q_n(r) - \lambda_{n+1} r^{2n+4}] \ln(1 - r^2) &< \mathcal{E}(r) \\ &< \frac{\pi}{2} - r \operatorname{arctanh} r - [Q_n(r) - \delta_{n+1} r^{2n+4}] \ln(1 - r^2) \end{aligned} \quad (42)$$

and

$$\begin{aligned} \left(\frac{8 - \pi}{4} - \frac{64 - 15\pi}{192} r^2 - \frac{384 - 95\pi}{3840} r^4 - \frac{2716 - 85\pi}{3840} r^6 \right) \frac{\ln(1 - r^2)}{2} \\ > \frac{\pi}{2} - r \operatorname{arctanh} r - \mathcal{E}(r) > \\ \left(\frac{8 - \pi}{4} - \frac{64 - 15\pi}{192} r^2 - \frac{384 - 95\pi}{3840} r^4 - \frac{274432 - 69195\pi}{516095} r^6 \right) \frac{\ln(1 - r^2)}{2} \end{aligned} \quad (43)$$

are valid.

Proof. From Theorem 2, it follows that

$$G_{2,n}(r) = \frac{1}{r^{n+1}} \left(\sum_{k=0}^n b_k r^k - \sum_{k=0}^{\infty} b_k r^k \right) = - \sum_{k=n+1}^{\infty} b_k r^{k-n-1} = \sum_{k=0}^{\infty} |b_{k+n+1}| r^k.$$

Consequently, the function $G_{2,n}(r)$ is absolutely monotonic on $(0, 1)$ and it is easy to see that the limits $G_{2,n}(0^+) = |b_{n+1}|$ and

$$G_{2,n}(1^-) = \sum_{k=0}^n b_k - G(1^-) = \sum_{k=0}^n b_k - \frac{1}{2}$$

are valid.

Since the function $G_{2,n}(r)$ is increasing and convex on $(0, 1)$, we deduce the double inequality

$$\begin{aligned} |b_{n+1}| + |b_{n+2}|r < G_{2,n}(r) < |b_{n+1}| + \left[\sum_{k=0}^n b_k - G(1^-) - |b_{n+1}| \right] r \\ &= |b_{n+1}| + \lambda_{n+1}r. \end{aligned} \quad (44)$$

Combining (4), (25), and (44) leads to (42).

Taking $n = 1$ in (42) results in (43). The proof of Corollary 2 is complete. \square

REMARK 3. The double inequality (42) is better than (10).

REMARK 4. As discussed in Remark 2, in terms of the notion of normalized remainders defined by (31), the function $G_{2,n}(r)$ defined in (41) can be reformulated as

$$T_n[G(r)] = -\frac{1}{b_{n+1}}G_{2,n}(r), \quad r \in (0, 1)$$

for $n \in \mathbb{N}_0$. Making use of the negativity in (15) and utilizing Corollary 2, we reveal that the normalized remainder $T_n[G(r)]$ for $n \in \mathbb{N}_0$ is absolutely monotonic from $(0, 1)$ onto $(1, 1 + \frac{\lambda_n}{\delta_n})$.

Making use of (7) and Corollaries 1 and 2, we derive the following corollary.

COROLLARY 3. For $r \in (0, 1)$ and $\kappa \in (1, \infty)$,

$$\varphi_\kappa(r) < r^{1/\kappa} \exp \left\{ \frac{4 \ln 2}{\pi - 2} \left(1 - \frac{1}{\kappa} \right) \min\{S(r), T(r)\} \right\}, \quad (45)$$

where a_n and b_n for $n \in \mathbb{N}_0$ are defined as in Theorems 1 and 2,

$$S(r) = \frac{\pi}{2} - 1 - \frac{\pi}{2} \left[1 - (1 - r^2) \frac{\operatorname{arctanh} r}{r} \right] \left[\sum_{k=0}^{n+1} a_k r^{2k} - \left(\sum_{k=0}^{n+1} a_k - 1 + \frac{2}{\pi} \right) r^{2n+4} \right],$$

and

$$T(r) = \frac{\pi}{2} - 1 - r \operatorname{arctanh} r - \left(\sum_{k=0}^{n+1} b_k r^{2k} + b_{n+2} r^{2n+4} \right) \ln(1 - r^2).$$

REMARK 5. The upper bound in (45) is better than the corresponding one in (7).

Acknowledgements. The authors appreciate anonymous referees for their helpful suggestions, careful corrections, and valuable comments on the original version of this paper.

REFERENCES

- [1] M. ABRAMOWITZ AND I. A. STEGUN (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series **55**, reprint of the 1972 edition, Dover Publications, Inc., New York, 1992.
- [2] R. P. AGARWAL, E. KARAPINAR, M. KOSTIĆ, J. CAO, AND W.-S. DU, *A brief overview and survey of the scientific work by Feng Qi*, *Axioms* **11** (2022), no. 8, Paper No. 385, 27 pages; available online at <https://doi.org/10.3390/axioms11080385>.
- [3] H. ALZER AND S.-L. QIU, *Monotonicity theorems and inequalities for the complete elliptic integrals*, *J. Comput. Appl. Math.* **172** (2004), no. 2, 289–312; available online at <https://doi.org/10.1016/j.cam.2004.02.009>.
- [4] H. ALZER AND K. RICHARDS, *A note on a function involving complete elliptic integrals: monotonicity, convexity, inequalities*, *Anal. Math.* **41** (2015), no. 3, 133–139; available online at <https://doi.org/10.1007/s10476-015-0201-7>.
- [5] G. D. ANDERSON, M. K. VAMANAMURTHY, AND M. VUORINEN, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.
- [6] G. E. ANDREWS, R. ASKEY, AND R. ROY, *Special Functions*, *Encyclopedia of Mathematics and its Applications* **71**, Cambridge University Press, Cambridge, 1999; available online at <https://doi.org/10.1017/CB09781107325937>.
- [7] Z.-H. BAO, R. P. AGARWAL, F. QI, AND W.-S. DU, *Some properties on normalized tails of Maclaurin power series expansion of exponential function*, *Symmetry* **16** (2024), no. 8, Art. 989, 15 pages; available online at <https://doi.org/10.3390/sym16080989>.
- [8] C. BERG, *Integral representation of some functions related to the gamma function*, *Mediterr. J. Math.* **1** (2004), no. 4, 433–439; available online at <https://doi.org/10.1007/s00009-004-0022-6>.
- [9] L. CARLITZ, *The Staudt–Clausen theorem*, *Math. Mag.* **34** (1960/61), 131–146; available online at <https://doi.org/10.2307/2688488>.
- [10] W.-S. DU, R. P. AGARWAL, E. KARAPINAR, M. KOSTIĆ, AND J. CAO, *Preface to the Special Issue “A Themed Issue on Mathematical Inequalities, Analytic Combinatorics and Related Topics in Honor of Professor Feng Qi”*, *Axioms* **12** (2023), no. 9, Art. 846, 5 pages; available online at <https://doi.org/10.3390/axioms12090846>.
- [11] I. S. GRADSHTEYN AND I. M. RYZHIK, *Table of Integrals, Series, and Products*, Translated from the Russian, Translation edited and with a preface by Daniel Zwillinger and Victor Moll, Eighth edition, Revised from the seventh edition, Elsevier/Academic Press, Amsterdam, 2015; available online at <https://doi.org/10.1016/B978-0-12-384933-5.00013-8>.
- [12] B.-N. GUO AND F. QI, *A property of logarithmically absolutely monotonic functions and the logarithmically complete monotonicity of a power-exponential function*, *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.* **72** (2010), no. 2, 21–30.
- [13] B.-N. GUO AND F. QI, *Some bounds for the complete elliptic integrals of the first and second kind*, *Math. Inequal. Appl.* **14** (2011), no. 2, 323–334; available online at <https://doi.org/10.7153/mia-14-26>.
- [14] F. T. HOWARD, *A special class of Bell polynomials*, *Math. Comp.* **35** (1980), no. 151, 977–989; available online at <https://doi.org/10.2307/2006208>.
- [15] F. T. HOWARD, *A sequence of numbers related to the exponential function*, *Duke Math. J.* **34** (1967), 599–615; available online at <http://projecteuclid.org/euclid.dmj/1077377163>.
- [16] F. T. HOWARD, *A sequence of numbers related to the exponential function*, Thesis (Ph.D.)–Duke University. ProQuest LLC, Ann Arbor, MI, 1966. 48 pp. available online at http://gateway.proquest.com/openurl?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:dissertation&res_dat=xri:pqdiss&rft_dat=xri:pqdiss:6706872.
- [17] F. T. HOWARD, *Associated Stirling numbers*, *Fibonacci Quart.* **18** (1980), no. 4, 303–315.
- [18] F. T. HOWARD, *Numbers generated by the reciprocal of $e^x - x - 1$* , *Math. Comp.* **31** (1977), no. 138, 581–598; available online at <https://doi.org/10.2307/2006440>.
- [19] F. T. HOWARD, *Some sequences of rational numbers related to the exponential function*, *Duke Math. J.* **34** (1967), 701–716; available online at <http://projecteuclid.org/euclid.dmj/1077377305>.
- [20] O. HÜBNER, *Remarks on a paper by Ławrynowicz on quasiconformal mappings*, *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.* **18** (1970), 183–186.

- [21] O. LEHTO AND K. I. VIRTANEN, *Quasiconformal Mappings in the Plane*, second edition, Translated from the German by K. W. Lucas. Die Grundlehren der mathematischen Wissenschaften, Band 126. Springer-Verlag, New York-Heidelberg, 1973.
- [22] Y.-W. LI AND F. QI, *A new closed-form formula of the Gauss hypergeometric function at specific arguments*, *Axioms* **13** (2024), no. 5, Art. 317, 24 pages; available online at <https://doi.org/10.3390/axioms13050317>.
- [23] Y.-F. LI AND F. QI, *A series expansion of a logarithmic expression and a decreasing property of the ratio of two logarithmic expressions containing cosine*, *Open Math.* **21** (2023), no. 1, Paper No. 20230159, 12 pages; available online at <https://doi.org/10.1515/math-2023-0159>.
- [24] Y.-W. LI, F. QI, AND W.-S. DU, *Two forms for Maclaurin power series expansion of logarithmic expression involving tangent function*, *Symmetry* **15** (2023), no. 9, Art. 1686, 18 pages; available online at <https://doi.org/10.3390/sym15091686>.
- [25] X.-L. LIU, H.-X. LONG, AND F. QI, *A series expansion of a logarithmic expression and a decreasing property of the ratio of two logarithmic expressions containing sine*, *Mathematics* **11** (2023), no. 14, Art. 3107, 12 pages; available online at <https://doi.org/10.3390/math11143107>.
- [26] X.-L. LIU AND F. QI, *Monotonicity results of ratio between two normalized remainders of Maclaurin series expansion for square of tangent function*, *Math. Slovaca* **75** (2025), no. 3, 699–705; available online at <https://doi.org/10.1515/ms-2025-0051>.
- [27] X. MA, Y. CHU, AND F. WANG, *Monotonicity and inequalities for the generalized distortion function*, *Acta Math. Sci. Ser. B (Engl. Ed.)* **33** (2013), no. 6, 1759–1766; available online at [https://doi.org/10.1016/S0252-9602\(13\)60121-6](https://doi.org/10.1016/S0252-9602(13)60121-6).
- [28] T. MANSOUR AND M. SCHORK, *Commutation Relations, Normal Ordering, and Stirling Numbers*, Discrete Mathematics and its Applications. CRC Press, Boca Raton, FL, 2016.
- [29] G. V. MILOVANOVIĆ AND F. QI, *Closed-form formulas of two Gauss hypergeometric functions of specific parameters*, *J. Math. Anal. Appl.* **543** (2025), no. 2, part 3, Paper No. 129024, 35 pages; available online at <https://doi.org/10.1016/j.jmaa.2024.129024>.
- [30] D. S. MITRINOVIĆ, J. E. PEČARIĆ, AND A. M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht-Boston-London, 1993; available online at <https://doi.org/10.1007/978-94-017-1043-5>.
- [31] D.-W. NIU AND F. QI, *Monotonicity results of ratios between normalized tails of Maclaurin power series expansions of sine and cosine*, *Mathematics* **12** (2024), no. 12, Art. 1781, 20 pages; available online at <https://doi.org/10.3390/math12121781>.
- [32] W.-J. PEI AND B.-N. GUO, *Monotonicity, convexity, and Maclaurin series expansion of Qi's normalized remainder of Maclaurin series expansion with relation to cosine*, *Open Math.* **22** (2024), no. 1, Paper No. 20240095, 11 pages; available online at <https://doi.org/10.1515/math-2024-0095>.
- [33] G. PÓLYA AND G. SZEGÖ, *Problems and Theorems in Analysis*, Volume I, *Series, integral calculus, theory of functions*, Translated from the German by D. Aeppli. Corrected printing of the revised translation of the fourth German edition. Grundlehren der Mathematischen Wissenschaften, 193. Springer-Verlag, Berlin-New York, 1978.
- [34] A. P. PRUDNIKOV, YU. A. BRYCHKOV, AND O. I. MARICHEV, *Integrals and Series*, Vol. 3. *More Special Functions*. Translated from the Russian by G. G. Gould. Gordon and Breach Science Publishers, New York, 1990.
- [35] F. QI, *Absolute monotonicity of normalized tail of power series expansion of exponential function*, *Mathematics* **12** (2024), no. 18, Art. 2859, 11 pages; available online at <https://doi.org/10.3390/math12182859>.
- [36] F. QI, *Series and connections among central factorial numbers, Stirling numbers, inverse of Vandermonde matrix, and normalized remainders of Maclaurin series expansions*, *Mathematics* **13** (2025), no. 2, Art. 223, 52 pages; available online at <https://doi.org/10.3390/math13020223>.
- [37] F. QI, R. P. AGARWAL, AND D. LIM, *Decreasing property of ratio of two logarithmic expressions involving tangent function*, *Math. Comput. Model. Dyn. Syst.* **31** (2025), no. 1, 1–16; available online at <https://doi.org/10.1080/13873954.2024.2449322>.
- [38] F. QI AND C.-P. CHEN, *A complete monotonicity property of the gamma function*, *J. Math. Anal. Appl.* **296** (2004), 603–607; available online at <https://doi.org/10.1016/j.jmaa.2004.04.026>.

- [39] F. QI, L.-H. CUI, AND S.-L. XU, *Some inequalities constructed by Tchebysheff's integral inequality*, Math. Inequal. Appl. **2** (1999), no. 4, 517–528; available online at <https://doi.org/10.7153/mia-02-42>.
- [40] F. QI AND C.-J. HUANG, *Computing sums in terms of beta, polygamma, and Gauss hypergeometric functions*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **114** (2020), Art. 191, 9 pages; available online at <https://doi.org/10.1007/s13398-020-00927-y>.
- [41] F. QI AND Z. HUANG, *Inequalities of the complete elliptic integrals*, Tamkang J. Math. **29** (1998), no. 3, 165–169; available online at <https://doi.org/10.5556/j.tkjm.29.1998.4242>.
- [42] F. QI, D.-W. NIU, D. LIM, AND Y.-H. YAO, *Special values of the Bell polynomials of the second kind for some sequences and functions*, J. Math. Anal. Appl. **491** (2020), no. 2, Art. 124382, 31 pages; available online at <https://doi.org/10.1016/j.jmaa.2020.124382>.
- [43] S.-L. QIU, X.-Y. MA, AND Y.-M. CHU, *Sharp Landen transformation inequalities for hypergeometric functions, with applications*, J. Math. Anal. Appl. **474** (2019), no. 2, 1306–1337; available online at <https://doi.org/10.1016/j.jmaa.2019.02.018>.
- [44] S.-L. QIU, X.-Y. MA, AND Y.-M. CHU, *Transformation properties of hypergeometric functions and their applications*, Comput. Methods Funct. Theory **22** (2022), no. 2, 323–366; available online at <https://doi.org/10.1007/s40315-021-00390-9>.
- [45] S.-L. QIU AND M. K. VAMANAMURTHY, *Elliptic integrals and the modulus of Grötzsch ring*, PanAmer. Math. J. **5** (1995), no. 2, 41–60.
- [46] S.-L. QIU AND M. VUORINEN, *Infinite products and normalized quotients of hypergeometric functions*, SIAM J. Math. Anal. **30** (1999), no. 5, 1057–1075; available online at <https://doi.org/10.1137/S0036141097326805>.
- [47] J. QUAINANCE AND H. W. GOULD, *Combinatorial Identities for Stirling Numbers*, The unpublished notes of H. W. Gould. With a foreword by George E. Andrews. World Scientific Publishing Co. Pte. Ltd., Singapore, 2016.
- [48] R. L. SCHILLING, R. SONG, AND Z. VONDRAČEK, *Bernstein Functions*, 2nd ed., de Gruyter Studies in Mathematics **37**, Walter de Gruyter, Berlin, Germany, 2012; available online at <https://doi.org/10.1515/9783110269338>.
- [49] N. M. TEMME, *Special Functions: An Introduction to Classical Functions of Mathematical Physics*, A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1996; available online at <https://doi.org/10.1002/9781118032572>.
- [50] A. WAN AND F. QI, *Power series expansion, decreasing property, and concavity related to logarithm of normalized tail of power series expansion of cosine*, Electron. Res. Arch. **32** (2024), no. 5, 3130–3144; available online at <https://doi.org/10.3934/era.2024143>.
- [51] M. K. WANG, Y. M. CHU, Y. P. JIANG, AND D. D. YAN, *A class of quadratic transformation inequalities for zero-balanced hypergeometric functions*, Acta Math. Sci. Ser. A (Chinese Ed.) **34** (2014), no. 4, 999–1007. (Chinese)
- [52] M.-K. WANG, Y.-M. CHU, AND S.-L. QIU, *Sharp bounds for generalized elliptic integrals of the first kind*, J. Math. Anal. Appl. **429** (2015), no. 2, 744–757; available online at <https://doi.org/10.1016/j.jmaa.2015.04.035>.
- [53] M.-K. WANG, Y.-M. CHU, AND Y.-Q. SONG, *Asymptotical formulas for Gaussian and generalized hypergeometric functions*, Appl. Math. Comput. **276** (2016), 44–60; available online at <https://doi.org/10.1016/j.amc.2015.11.088>.
- [54] M.-K. WANG, S.-L. QIU, Y.-M. CHU, AND Y.-P. JIANG, *Generalized Hersch-Pfluger distortion function and complete elliptic integrals*, J. Math. Anal. Appl. **385** (2012), no. 1, 221–229; available online at <https://doi.org/10.1016/j.jmaa.2011.06.039>.
- [55] F. WANG, B.-N. GUO, AND F. QI, *Monotonicity and inequalities related to complete elliptic integrals of the second kind*, AIMS Math. **5** (2020), no. 3, 2732–2742; available online at <https://doi.org/10.3934/math.2020176>.
- [56] F. WANG AND F. QI, *Monotonicity and sharp inequalities related to complete (p, q) -elliptic integrals of the first kind*, C. R. Math. Acad. Sci. Paris **358** (2020), no. 8, 961–970; available online at <https://doi.org/10.5802/crmath.119>.
- [57] F. WANG AND F. QI, *Power series expansion and decreasing property related to normalized remainders of power series expansion of sine*, Filomat **38** (2024), no. 29, 10447–10462; available online at <https://doi.org/10.2298/FIL2429447W>.

- [58] G. D. WANG, X. H. ZHANG, Y. M. CHU, AND S. L. QIU, *Complete elliptic integrals and the Hersch–Pfluger distortion function*, Acta Math. Sci. Ser. A (Chinese Ed.) **28** (2008), no. 4, 731–734. (Chinese)
- [59] D. V. WIDDER, *The Laplace Transform*, Princeton Mathematical Series **6**, Princeton University Press, Princeton, N. J., 1946.
- [60] Z.-H. YANG, Y.-M. CHU, AND W. ZHANG, *Accurate approximations for the complete elliptic integral of the second kind*, J. Math. Anal. Appl. **438** (2016), no. 2, 875–888; available online at <https://doi.org/10.1016/j.jmaa.2016.02.035>.
- [61] Z.-H. YANG AND J. TIAN, *Sharp inequalities for the generalized elliptic integrals of the first kind*, Ramanujan J. **48** (2019), no. 1, 91–116; available online at <https://doi.org/10.1007/s11139-018-0061-4>.
- [62] Z.-H. YANG; J.-F. TIAN, AND Y.-R. ZHU, *A sharp lower bound for the complete elliptic integrals of the first kind*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM **115** (2021), no. 1, Paper No. 8, 17 pp.; available online at <https://doi.org/10.1007/s13398-020-00949-6>.
- [63] Z.-H. YANG, M.-K. WANG, J.-F. TIAN, *Series expansion, higher-order monotonicity properties and inequalities for the modulus of the Grötzsch ring*, Proc. Edinburgh Math. Soc. Published online 2024:1-28; available online at <https://doi.org/10.1017/S0013091524000440>.
- [64] L. YIN, X.-L. LIN, AND F. QI, *Monotonicity, convexity and inequalities related to complete (p, q, r) -elliptic integrals and generalized trigonometric functions*, Publ. Math. Debrecen **97** (2020), no. 1-2, 181–199; available online at <https://doi.org/10.5486/PMD.2020.8793>.
- [65] L. YIN AND F. QI, *Some inequalities for complete elliptic integrals*, Appl. Math. E-Notes **14** (2014), 192–199.
- [66] H.-C. ZHANG, B.-N. GUO, AND W.-S. DU, *On Qi's normalized remainder of Maclaurin power series expansion of logarithm of secant function*, Axioms **13** (2024), no. 12, Art. 860, 11 pages; available online at <https://doi.org/10.3390/axioms13120860>.
- [67] G.-Z. ZHANG AND F. QI, *On convexity and power series expansion for logarithm of normalized tail of power series expansion for square of tangent*, J. Math. Inequal. **18** (2024), no. 3, 937–952; available online at <https://doi.org/10.7153/jmi-2024-18-51>.
- [68] G.-Z. ZHANG, Z.-H. YANG, AND F. QI, *On normalized tails of series expansion of generating function of Bernoulli numbers*, Proc. Amer. Math. Soc. **153** (2025), no. 1, 131–141; available online at <https://doi.org/10.1090/proc/16877>.
- [69] T. ZHANG, Z.-H. YANG, F. QI, AND W.-S. DU, *Some properties of normalized tails of Maclaurin power series expansions of sine and cosine*, Fractal Fract. **8** (2024), no. 5, Art. 257, 17 pages; available online at <https://doi.org/10.3390/fractalfract8050257>.
- [70] X. ZHANG, G. WANG, AND Y. CHU, *Remarks on generalized elliptic integrals*, Proc. Roy. Soc. Edinburgh Sect. A **139** (2009), no. 2, 417–426; available online at <https://doi.org/10.1017/S0308210507000327>.

(Received December 20, 2024)

Fei Wang

Department of Mathematics
Zhejiang Polytechnic University of Mechanical and Electrical Engineering
Hangzhou 310053, Zhejiang, China
e-mail: wf509529@163.com
<https://orcid.org/0000-0002-7175-2272>

Feng Qi

School of Mathematics and Informatics
Henan Polytechnic University
Jiaozuo, Henan, 454010, China
17709 Sabal Court, Dallas, TX 75252-8024, USA
and
School of Mathematics and Physics
Hulunbuir University
Hulunbuir, Inner Mongolia, 021008, China
e-mail: qifeng618@gmail.com
<https://orcid.org/0000-0001-6239-2968>