

STRICT SINGULARITY AND CLOSED RANGE OF VOLTERRA INTEGRATION OPERATOR ON DIRICHLET–MORREY TYPE SPACES

XIANGLING ZHU

(Communicated by L. Mihoković)

Abstract. The strict singularity and closed range property of the Volterra integral operator T_g with the symbol g on the Dirichlet–Morrey type space $\mathcal{D}_{\lambda,k}$ were investigate in this paper.

1. Introduction

Let \mathbb{D} represent the open unit disk within the complex plane \mathbb{C} , and let $H(\mathbb{D})$ denote the class of functions that are analytic in \mathbb{D} . Let $H^\infty = H^\infty(\mathbb{D})$ denote the space of all bounded analytic functions on \mathbb{D} . The norm in this space is given by $\|f\|_\infty = \sup_{w \in \mathbb{D}} |f(w)|$. The Bloch space, denoted by $\mathcal{B}(\mathbb{D}) = \mathcal{B}$, consists of all $f \in H(\mathbb{D})$ satisfying

$$\|f\|_{\mathcal{B}} = |f(0)| + \sup_{w \in \mathbb{D}} (1 - |w|^2) |f'(w)| < \infty.$$

The space \mathcal{B} is a Banach space with respect to the norm $\|\cdot\|_{\mathcal{B}}$, meaning it is complete under this norm. It can be shown that $H^\infty \subset \mathcal{B}$. The little Bloch space, denoted by \mathcal{B}_0 , contains all $f \in H(\mathbb{D})$ such that $\lim_{|w| \rightarrow 1} (1 - |w|^2) |f'(w)| = 0$.

Let $0 < p < \infty$ and $\alpha > -1$. The weighted Bergman space A_α^p consists of all analytic functions f in \mathbb{D} , i.e., $f \in H(\mathbb{D})$, for which the following norm is finite:

$$\|f\|_{A_\alpha^p}^p = (\alpha + 1) \int_{\mathbb{D}} |f(w)|^p (1 - |w|^2)^\alpha dA(w) < \infty.$$

In the above expression, dA represents the normalized area measure on \mathbb{D} . A function $f \in H(\mathbb{D})$ is said to be in the weighted Dirichlet space $\mathcal{D}_\alpha^p(\mathbb{D}) = \mathcal{D}_\alpha^p$ if

$$\|f\|_{\mathcal{D}_\alpha^p}^p = |f(0)|^p + \int_{\mathbb{D}} |f'(w)|^p (1 - |w|^2)^\alpha dA(w) < \infty.$$

It is worth to note that when $\alpha = 1$ and $p = 2$, the weighted Dirichlet space \mathcal{D}_α^p is identical to the classical Hardy space H^2 .

Mathematics subject classification (2020): 30H99, 47B38.

Keywords and phrases: Dirichlet–Morrey type space, Volterra integral operator, closed range, strict singularity.

Let $0 < p < \infty$, $-2 < q < \infty$, $0 \leq s < \infty$. A function $g \in H(\mathbb{D})$ is in $F(p, q, s)$ if

$$\|g\|_{F(p,q,s)}^p = |g(0)|^p + \sup_{b \in \mathbb{D}} \int_{\mathbb{D}} |g'(w)|^p (1 - |w|^2)^q (1 - |\sigma_b(w)|^2)^s dA(w) < \infty.$$

Here $\sigma_b(w) = \frac{b-w}{1-\bar{b}w}$ denotes the automorphism of \mathbb{D} exchanging 0 for b . A function g belongs to the space $F_0(p, q, s)$ if

$$\lim_{|b| \rightarrow 1} \int_{\mathbb{D}} |g'(w)|^p (1 - |w|^2)^q (1 - |\sigma_b(w)|^2)^s dA(w) = 0.$$

Zhao introduced $F(p, q, s)$ in [41]. From [41], when $q = p - 2$, $F(p, p - 2, s) = \mathcal{B}$ if $s > 1$, $F(p, p - 2, 0) = B_p$. When $p = 2$, $F(p, p - 2, s) = Q_s$. In particular, $F(2, 0, 1) = BMOA$, the set of all analytic functions of bounded mean oscillation. For a generalization of the space on the unit ball and some operators acting from or into it see, e.g., [14, 32, 33] and the references therein.

Let $K : [0, \infty) \rightarrow [0, \infty)$ be a non-decreasing, right-continuous function. Assume K is not identically zero, with $K(0) = 0$, $K(t) > 0$ for every $t > 0$, and $K(t) = K(1)$ for all $t \geq 1$. For $-1 < \lambda < 0$, as Hu and Liu [8] defined, a class of Dirichlet-Morrey type space $\mathcal{D}_{\lambda,K}$ consists of all functions $f \in H(\mathbb{D})$ such that

$$\|f\|_{\mathcal{D}_{\lambda,K}} = |f(0)| + \sup_{b \in \mathbb{D}} \frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} \|f \circ \sigma_b - f(b)\|_{\mathcal{D}_{\lambda}^1} < \infty.$$

We denote $\|f\|_{\mathcal{D}_{\lambda,K}} - |f(0)|$ by $\|f\|_{*,\mathcal{D}_{\lambda,K}}$. Analogously, an analytic function f is said to belong to the little Dirichlet-Morrey space $\mathcal{D}_{\lambda,K}^0$ if

$$\lim_{|b| \rightarrow 1} \frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} \|f \circ \sigma_b - f(b)\|_{\mathcal{D}_{\lambda}^1} = 0.$$

See [26, 35, 36, 38–40, 42, 44, 45] for various analytic Morrey type spaces, including Hardy-Morrey spaces, Bergman-Morrey spaces, Dirichlet-Morrey spaces and Besov-Morrey spaces.

Let $g \in H(\mathbb{D})$. The Volterra integral operator T_g is defined as follows:

$$T_g f(z) = \int_0^z f(w) g'(w) dw, \quad f \in H(\mathbb{D}), \quad z \in \mathbb{D}.$$

Pommerenke [22] was probably the first who studied the operator. He showed that T_g is bounded on the Hardy space H^2 if and only if $g \in BMOA$ —the space of bounded mean oscillation. Up to the present day, the operator T_g along with its generalizations, like generalized Cesàro or Volterra operators or their products with composition operators, have attracted significant attention within the mathematical community. Extensive research has been carried out on the boundedness, compactness, norm, and essential norm of T_g over various function spaces, yielding rich results. For a comprehensive overview of additional findings regarding this operator and its various generalizations, one may refer to [1, 2, 4–9, 11–16, 20, 21, 23–36, 38–40, 42, 44, 45].

Recently, the study of the strict singularity of the integral operator T_g in various function spaces has piqued interest. Let us recall that for a bounded linear operator $T : X \rightarrow Y$ between Banach spaces, as introduced by Kato in [10], it is said to be strictly singular if for any infinite-dimensional closed subspace $M \subset X$, the restriction $T|_M$ is not an isomorphism onto its image. Precisely, there does not exist a positive constant c such that $\|T(y)\| \geq c\|y\|$ for all $y \in M$. Evidently, every compact operator is strictly singular. A well-known example of strictly singular yet non-compact operators is the inclusion mapping $i_{p,q} : \ell^p \rightarrow \ell^q$, where $1 \leq p < q < \infty$. In 2017, Miihkinen, in the work [20], demonstrated that for a non-compact operator $T_g : H^p \rightarrow H^p$ with $1 \leq p < \infty$, it contains an isomorphic copy of l^p within H^p . As a consequence, for the operator T_g on H^p , the properties of compactness and strict singularity are equivalent. In the context of Bergman spaces A^p for $1 \leq p < \infty$, the strict singularity of the operator T_g is equivalent to its compactness. In [7], Chen established that for the operator $T_g : F(p, p-2, s) \rightarrow F(p, p-2, s)$ with $s, p > 0$ satisfying $s+p > 1$, the compactness and strict singularity are identical. See [17] for the study of the strict singularity of another integral operator S_g on H^p .

Meanwhile, studying the closed range of the operator has also attracted a lot of attention. Using the closed graph theorem, we see that T has closed range if and only if it is bounded below when T is a one-to-one bounded linear operator on X . Hence, to investigate the closed range of the operator T_g , we only need to consider the boundedness below of the operator T_g . Recall that a linear operator T on a quasi-Banach space $(X, \|\cdot\|)$ is said to be bounded below if there exists $C > 0$ such that

$$\|Tx\| \geq C\|x\|$$

for all $x \in X$. Anderson [3] showed that T_g can be bounded below on weighted Bergman spaces. In 2014, Anderson, Jovovic, and Smith showed that T_g is never bounded below on the Hardy space H^2 , the Bloch space \mathcal{B} , and the space $BMOA$ [4]. Chen [7] proved that T_g also has no lower bound on $F(p, pa-2, s)$.

Motivated by the above-mentioned works, in this paper, we study the strictly singular property and closed range property of the integration operator T_g on the Dirichlet-Morrey type space $\mathcal{D}_{\lambda,K}$. We show that when $g \in F(1, -1, \lambda+1) \setminus F_0(1, -1, \lambda+1)$, T_g is not strictly singular. Also, T_g has no closed range on $\mathcal{D}_{\lambda,K}$. Moreover, we also investigate some properties of $\mathcal{D}_{\lambda,K}^0$.

Throughout the paper, we write $A \lesssim B$ (or $B \gtrsim A$) to denote that there is some inessential constant C such that $A \leq CB$. If $A \lesssim B \lesssim A$, then we write $A \asymp B$.

2. Vanishing K -Carleson measure and embedding of $\mathcal{D}_{\lambda,K}^0$

In this section, we investigate some basic properties of $\mathcal{D}_{\lambda,K}^0$. We require certain properties of K . Throughout the remainder of this paper, we consistently assume that the subsequent condition regarding K is satisfied (refer to [37]):

$$\int_1^\infty \frac{\varphi_K(x)}{x^{1+\delta}} dx < \infty, \quad \delta > 0, \quad (2.1)$$

where

$$\varphi_K(x) = \sup_{0 < s \leq 1} \frac{K(sx)}{K(s)}, \quad 0 < x < \infty.$$

Obviously, $K(x) = x^p$ satisfies inequality (2.1) for $0 < p < \delta$.

LEMMA 2.1. ([37], Theorem 3.7) *If K satisfies condition (2.1) for some $\delta > 0$, then there exists a weight function K_1 such that*

$$\lim_{t \rightarrow 0^+} \frac{K_1(t)}{t^\delta} = \infty.$$

Furthermore, K_1 still satisfies all standing assumptions on weights, K_1 is comparable with K on $(0, 1)$.

For arc I on the unit circle $\partial\mathbb{D}$, $|I|$ is the normalized arc length ($|\partial\mathbb{D}| = 1$). Let

$$S(I) = \{w = re^{i\theta} : 1 - |I| < |w| < 1, e^{i\theta} \in I\}$$

denote the Carleson box based on I . A positive Borel measure μ on \mathbb{D} is called a K -Carleson measure if (see [35])

$$\|\mu\|_K = \sup_{I \subset \partial\mathbb{D}} \frac{\mu(S(I))}{K(|I|)} < \infty.$$

When $K(t) = t^s$, $0 < s < \infty$, the K -Carleson measure is an s -Carleson measure. When $K(t) = t$, μ is the classical Carleson measure. The following lemma gives a characterization for K -Carleson measure.

LEMMA 2.2. ([35]) *Suppose K satisfies (2.1) for some $\delta \in (0, 2)$. Let μ be a positive Borel measure on \mathbb{D} . The measure μ is a K -Carleson measure if and only if*

$$\sup_{b \in \mathbb{D}} \frac{1}{K(1 - |b|^2)} \int_{\mathbb{D}} \left(\frac{1 - |b|^2}{|1 - \bar{b}w|} \right)^t d\mu(w) < \infty, \quad \delta \leq t < \infty.$$

Similarly, we call μ a vanishing K -Carleson measure if

$$\lim_{|I| \rightarrow 0} \frac{\mu(S(I))}{K(|I|)} = 0.$$

Combining the proofs of Theorem 1 in [19] and Lemma 2.2, or Theorem 2.1 in [35], we can get the following lemma. Since the proof is routine, we omit the detail.

LEMMA 2.3. *Suppose K satisfies (2.1) for some $\delta \in (0, 2)$. Let μ be a positive Borel measure on \mathbb{D} . The measure μ is a vanishing K -Carleson measure if and only if*

$$\lim_{|b| \rightarrow 1} \frac{1}{K(1 - |b|^2)} \int_{\mathbb{D}} \left(\frac{1 - |b|^2}{|1 - \bar{b}w|} \right)^t d\mu(w) = 0, \quad \delta \leq t < \infty.$$

From the definition of $\mathcal{D}_{\lambda,K}^0$ and Lemma 2.3 we get the following characterization for the space $\mathcal{D}_{\lambda,K}^0$.

PROPOSITION 2.4. *Let $-1 < \lambda < 0$ and K satisfy (2.1) for some $\delta \in (0, \lambda + 1]$. Let $f \in H(\mathbb{D})$ and $d\mu_f(w) = |f'(w)|(1 - |w|^2)^\lambda dA(w)$. Then the following statements are equivalent:*

- (i) $f \in \mathcal{D}_{\lambda,K}^0$;
- (ii)

$$\lim_{|b| \rightarrow 1} \frac{(1 - |b|^2)^{1+\lambda}}{K(1 - |b|)} \int_{\mathbb{D}} |f'(w)|(1 - |w|^2)^{-1} (1 - |\sigma_b(w)|^2)^{\lambda+1} dA(w) = 0;$$

- (iii) μ is a vanishing K -Carleson measure.

The following integral estimate is of great importance in our proof.

LEMMA 2.5. [43, Lemma 3.10] *Suppose $z \in \mathbb{D}$, c is real, $t > -1$, and*

$$I_{c,t}(z) = \int_{\mathbb{D}} \frac{(1 - |w|^2)^t}{|1 - z\bar{w}|^{2+t+c}} dA(w).$$

(a) *If $c < 0$, then as a function of z , $I_{c,t}(z)$ is bounded from above and bounded from below on \mathbb{D} .*

(b) *If $c > 0$, then*

$$I_{c,t}(z) \asymp \frac{1}{(1 - |z|^2)^c}, \quad |z| \rightarrow 1^-.$$

The following two propositions illustrate the relationship between polynomials and $\mathcal{D}_{\lambda,K}^0$. We show that $\mathcal{D}_{\lambda,K}^0$ is the closure of all polynomials in $\mathcal{D}_{\lambda,K}$.

PROPOSITION 2.6. *Let $-1 < \lambda < 0$ and K satisfy (2.1) for some $\delta \in (0, \lambda + 1]$. Then $\mathcal{D}_{\lambda,K}^0$ (and so $\mathcal{D}_{\lambda,K}$) contains all polynomials.*

Proof. Since K satisfies (2.1) for some $\delta \in (0, \lambda + 1]$, by Lemma 2.1, there exists a weight function K_1 such that

$$t^\delta / K_1(t) \rightarrow 0,$$

as $t \rightarrow 0$. Because $\delta \leq \lambda + 1$, we have

$$\frac{(1 - |b|^2)^{\lambda+1}}{K_1(1 - |b|^2)} \leq \frac{(1 - |b|^2)^\delta}{K_1(1 - |b|^2)}.$$

Furthermore, K_1 still satisfies all standing assumptions on weights, K_1 is comparable with K on $(0, 1)$. Hence

$$\lim_{|b| \rightarrow 1} \frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} = 0. \quad (2.2)$$

If f is a polynomial, then $\sup_{w \in \mathbb{D}} |f'(w)| \leq M < \infty$. By Lemma 2.5,

$$\begin{aligned} & \frac{(1-|b|^2)^{1+\lambda}}{K(1-|b|^2)} \int_{\mathbb{D}} |f'(w)| (1-|w|^2)^{-1} (1-|\sigma_b(w)|^2)^{1+\lambda} dA(w) \\ & \leq M \frac{(1-|b|^2)^{1+\lambda}}{K(1-|b|^2)} \int_{\mathbb{D}} (1-|w|^2)^{-1} (1-|\sigma_b(w)|^2)^{1+\lambda} dA(w) \\ & \lesssim M \frac{(1-|b|^2)^{1+\lambda}}{K(1-|b|^2)} \int_{\mathbb{D}} \frac{(1-|w|^2)^\lambda}{|1-\bar{b}w|^{2+2\lambda}} dA(w) \\ & \lesssim M \frac{(1-|b|^2)^{1+\lambda}}{K(1-|b|^2)}. \end{aligned}$$

By (2.2), we obtain that

$$\lim_{|b| \rightarrow 1} \frac{(1-|b|^2)^{1+\lambda}}{K(1-|b|^2)} \int_{\mathbb{D}} |f'(w)| (1-|w|^2)^{-1} (1-|\sigma_b(w)|^2)^{1+\lambda} dA(w) = 0.$$

Thus $f \in \mathcal{D}_{\lambda,K}^0$. \square

Similarly to the proof of Proposition 2.15 in [41], we get the following proposition.

PROPOSITION 2.7. *Let $-1 < \lambda < 0$ and K satisfy (2.1) for some $\delta \in (0, \lambda + 1]$ and $g \in \mathcal{D}_{\lambda,K}^0$. Then*

$$\lim_{r \rightarrow 1} \|g_r - g\|_{\mathcal{D}_{\lambda,K}} = 0,$$

where $g_r(w) = g(rw)$. In particular, $\mathcal{D}_{\lambda,K}^0$ is the closure of all polynomials in $\mathcal{D}_{\lambda,K}$.

Let μ be a positive Borel measure on \mathbb{D} . The space $\mathcal{T}_K(\mu)$ consists of all measurable functions f that satisfy

$$\|f\|_{\mathcal{T}_K(\mu)} = \sup_{I \subset \partial\mathbb{D}} \frac{1}{K(|I|)} \int_{S(I)} |f(w)| d\mu(w) < \infty.$$

Let $\mathcal{T}_{K,0}(\mu)$ denote the space of all measure functions f such that

$$\lim_{|I| \rightarrow 0} \frac{1}{K(|I|)} \int_{S(I)} |f(w)| d\mu(w) = 0.$$

PROPOSITION 2.8. *Let $-1 < \lambda < 0$ and K satisfy (2.1) for some $\delta \in (0, \lambda + 1]$. Let μ be a positive Borel measure on \mathbb{D} . Then the identity operator $I_d : \mathcal{D}_{\lambda,K}^0 \rightarrow \mathcal{T}_{K,0}(\mu)$ is bounded if and only if μ is a $(\lambda + 1)$ -Carleson measure.*

Proof. Necessity. Suppose $I_d : \mathcal{D}_{\lambda,K}^0 \rightarrow \mathcal{T}_{K,0}(\mu)$ is bounded. Let $I \subset \partial\mathbb{D}$, $e^{i\theta}$ be the center of I and $a = (1 - |I|)e^{i\theta}$. It is easy to see that

$$|1 - \bar{a}w| \approx 1 - |a|^2 \approx |I|, \quad w \in S(I).$$

Let

$$f_a(w) = \frac{(1 - |a|^2)^{1+\lambda} K(1 - |a|^2)}{(1 - \bar{a}w)^{2\lambda+2}}, \quad w \in \mathbb{D}.$$

Clearly, we have that $f_a \in \mathcal{D}_{\lambda,K}^0$. Thus

$$\begin{aligned} \frac{\mu(S(I))}{|I|^{1+\lambda}} &= \frac{K(|I|)}{|I|^{1+\lambda}} \frac{1}{K(|I|)} \int_{S(I)} d\mu \\ &\approx \frac{1}{K(|I|)} \int_{S(I)} |f_a(w)| d\mu(w) \lesssim \|f_a\|_{\mathcal{T}_K(\mu)} \leq \|f_a\|_{\mathcal{D}_{\lambda,K}} < \infty, \end{aligned}$$

which implies that μ is a $(\lambda + 1)$ -Carleson measure.

Sufficiency. Suppose μ is a $(\lambda + 1)$ -Carleson measure. Carefully check the proof of Theorem 3.2 in [8], we see that the identity operator $I_d : \mathcal{D}_{\lambda,K} \rightarrow \mathcal{T}_K(\mu)$ is bounded. By Proposition 2.7, for any $\varepsilon > 0$, there exists a polynomial $P(z)$ such that

$$\|f - P\|_{\mathcal{D}_{\lambda,K}} < \varepsilon \quad (2.3)$$

for any $f \in \mathcal{D}_{\lambda,K}^0$. Thus,

$$\begin{aligned} \frac{1}{K(|I|)} \int_{S(I)} |f(w)| d\mu(w) &\lesssim \frac{1}{K(|I|)} \int_{S(I)} |f(w) - P(w)| d\mu(w) + \frac{1}{K(|I|)} \int_{S(I)} |P(w)| d\mu(w) \\ &\lesssim \|f - P\|_{\mathcal{T}_K(\mu)} + \|P\|_{H^\infty} \frac{\mu(S(I))}{|I|^{1+\lambda}} \frac{|I|^{1+\lambda}}{K(|I|)}. \end{aligned}$$

By Lemma 2.1, for $\delta \in (0, 1 + \lambda]$, we have that

$$\lim_{|I| \rightarrow 0} \frac{|I|^{1+\lambda}}{K(I)} = 0. \quad (2.4)$$

Combining (2.3) and (2.4), we get that

$$\lim_{|I| \rightarrow 0} \frac{1}{K(|I|)} \int_{S(I)} |f(w)| d\mu(w) = 0,$$

which implies $f \in \mathcal{T}_{K,0}(\mu)$. This completes the proof of this proposition. \square

3. Strict singularity

In this section, we mainly show that a non-compact operator T_g is not strictly singular, i.e., there is a subspace $M \subset \mathcal{D}_{\lambda,K}$ such that $T_g|_M$ is bounded below on M which yields the compactness and strict singularity are equivalent for T_g on $\mathcal{D}_{\lambda,K}$. In order to prove the main result in this section, we need the following lemma, see [8, Theorem 1 and Corollary 1].

LEMMA 3.1. *Let $g \in H(\mathbb{D})$, $-1 < \lambda < 0$ and K satisfy (2.1) for some $\delta \in (0, \lambda + 1]$. Then the following statements hold.*

1. T_g is bounded on $\mathcal{D}_{\lambda,k}$ if and only if $g \in F(1, -1, \lambda + 1)$;
2. T_g is compact on $\mathcal{D}_{\lambda,k}$ if and only if $g \in F_0(1, -1, \lambda + 1)$.

COROLLARY 3.2. *Let $g \in H(\mathbb{D})$, $-1 < \lambda < 0$ and K satisfy (2.1) for some $\delta \in (0, \lambda + 1]$. Then T_g is bounded on $\mathcal{D}_{\lambda,K}^0$ if and only if $g \in F(1, -1, \lambda + 1)$.*

Proof. Since $g \in F(1, -1, \lambda + 1)$ if and only if $d\mu_g = |g'(w)|(1 - |w|^2)^\lambda$ is a $(\lambda + 1)$ -Carleson measure, by Proposition 2.8, we can get this corollary. \square

Using Theorem 9 in [18], we immediately obtain the following result.

LEMMA 3.3. *Let $-1 < \lambda < 0$ and μ be a positive Borel measure on \mathbb{D} . Then μ is a $(\lambda + 1)$ -Carleson measure if and only if for all functions $f \in \mathcal{D}_\lambda^1$,*

$$\int_{\mathbb{D}} |f(w)| d\mu(w) \lesssim \|f\|_{\mathcal{D}_\lambda^1}.$$

COROLLARY 3.4. *Let $-1 < \lambda < 0$. Let $g \in F(1, -1, \lambda + 1)$. Then T_g is bounded on \mathcal{D}_λ^1 .*

Proof. By the assumption we see that $d\mu_g(w) := |g'(w)|(1 - |w|^2)^\lambda dA(w)$ is a $(\lambda + 1)$ -Carleson measure. By Lemma 3.3, $I_d : \mathcal{D}_\lambda^1 \rightarrow L(\mathbb{D}, d\mu_g)$ is bounded. Hence, for any $f \in \mathcal{D}_\lambda^1$,

$$\|T_g f\|_{\mathcal{D}_\lambda^1} = \int_{\mathbb{D}} |f(w)| |g'(w)| (1 - |w|^2)^\lambda dA(w) \lesssim \|f\|_{\mathcal{D}_\lambda^1},$$

as desired. \square

To prove the main result in this section, one of important steps is to construct the operator $Q : \mathcal{C}_0 \rightarrow \mathcal{D}_{\lambda,K}^0$ such that Q is bounded and is an isomorphism onto its range. Here \mathcal{C}_0 denote the Banach space of complex sequences converging to zero endowed with the supremum norm $\|\cdot\|_{l^\infty}$. We have the following result.

PROPOSITION 3.5. *Let $-1 < \lambda < 0$ and K satisfy (2.1) for some $\delta \in (0, 1 + \lambda]$. Let $\{f_n\}_{n=1}^\infty$ be a sequence in $\mathcal{D}_{\lambda,K}^0$ with $\|f_n\|_{*, \mathcal{D}_{\lambda,K}} \approx 1$ and $\lim_{n \rightarrow \infty} \|f_n\|_{\mathcal{D}_\lambda^1} = 0$. Then there exists a subsequence $\{f_{n_k}\} \subseteq \{f_n\}_{n=1}^\infty$ such that the operator $Q : \mathcal{C}_0 \rightarrow \mathcal{D}_{\lambda,K}^0$ defined as follows*

$$Q(\eta) = \sum_{k=1}^{\infty} \eta_k f_{n_k}, \quad \eta = \{\eta_k\} \subset \mathcal{C}_0,$$

is an isomorphism onto its range.

Proof. We denote

$$\Phi(f_n, b) = \frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} \int_{\mathbb{D}} |f'_n(w)| (1 - |w|^2)^{-1} (1 - |\sigma_b(w)|^2)^{\lambda+1} dA(w).$$

Since $f_n \in \mathcal{D}_{\lambda, K}^0$, for each $n \geq 1$, we have that

$$\lim_{|b| \rightarrow 1} \Phi(f_n, b) = 0. \quad (3.1)$$

If $|b| \leq t_0$ for some $t_0 \in (0, 1)$, according to the monotonicity of K and the fact that $\lim_{n \rightarrow \infty} \|f_n\|_{\mathcal{D}_{\lambda}^1} = 0$, we get

$$\Phi(f_n, b) \lesssim \frac{1}{K(1 - t_0^2)} \int_{\mathbb{D}} |f'_n(w)| (1 - |w|^2)^{\lambda} dA(w) \lesssim \frac{1}{K(1 - t_0^2)} \|f_n\|_{\mathcal{D}_{\lambda}^1} \rightarrow 0,$$

as $n \rightarrow \infty$. Therefore, for any $t \in (0, 1)$,

$$\lim_{n \rightarrow \infty} \sup_{|b| \leq t} \Phi(f_n, b) = 0. \quad (3.2)$$

By the assumed condition $\|f_n\|_{*, \mathcal{D}_{\lambda, K}} \asymp 1$, there are $S_1, S_2 > 0$ such that,

$$S_1 \leq \|f_n\|_{*, \mathcal{D}_{\lambda, K}} \leq S_2 \quad (3.3)$$

for any $n \geq 1$.

Combining (3.1), (3.2) with $\lim_{n \rightarrow \infty} \|f_n\|_{\mathcal{D}_{\lambda}^1} = 0$, there exists a subsequence $\{f_{n_k}\}$ and an increasing sequence $\{t_k\}$ with $t_k \in (0, 1)$ such that, for any $k \geq 1$,

$$\sup_{|b| \leq t_k} \Phi(f_{n_k}, b) < 3^{-k-1} S_1,$$

and

$$\sup_{|b| > t_{k+1}} \Phi(f_{n_k}, b) < 3^{-k-1} S_1$$

and

$$\|f_{n_k}\|_{\mathcal{D}_{\lambda}^1} < 3^{-k-1} S_1.$$

Since b belongs to at most one circular region $t_k < |b| \leq t_{k+1}$ within \mathbb{D} , for any $b \in \mathbb{D}$, the inequality

$$\Phi(f_{n_k}, b) < 3^{-k-1} S_1$$

holds for all but at most one index k . For the exceptional index k , $t_k < |b| \leq t_{k+1}$, the inequality $\Phi(f_{n_k}, b) \leq S_2$ is satisfied. In addition, from (3.3), for any $k \geq 1$, there is $b_k \in \mathbb{D}$ such that

$$\Phi(f_{n_k}, b_k) \geq \frac{2S_1}{3}.$$

By the triangle inequality, for any $\eta = \{\eta_k\} \in \mathcal{C}_0$, we obtain that

$$\begin{aligned} \|Q(\eta)\|_{\mathcal{D}_{\lambda,K}} &= \left| \sum_{k=1}^{\infty} \eta_k f_{n_k}(0) \right| + \sup_{b \in \mathbb{D}} \Phi(Q(\eta), b) \\ &\leq \|\eta\|_{l^\infty} \sum_{k=1}^{\infty} \|f_{n_k}\|_{\mathcal{D}_{\lambda}^1} + \sup_{b \in \mathbb{D}} \sum_{k=1}^{\infty} |\eta_k| \Phi(f_{n_k}, b) \\ &< \|\eta\|_{l^\infty} \sum_{k=1}^{\infty} 3^{-k-1} S_1 + \|\eta\|_{l^\infty} \left(\sum_{k=1}^{\infty} 3^{-k-1} S_1 + S_2 \right) \\ &< (S_1 + S_2) \|\eta\|_{l^\infty}, \end{aligned}$$

which implies that $Q(\eta) \in \mathcal{D}_{\lambda,K}$ and $Q: \mathcal{C}_0 \rightarrow \mathcal{D}_{\lambda,K}$ is bounded.

Next, we prove that $Q(\eta) \in \mathcal{D}_{\lambda,K}^0$. In fact, for any $b \in \mathbb{D}$ and $N > 1$,

$$\begin{aligned} \Phi(Q(\eta), b) &\leq \sum_{k=1}^{\infty} |\eta_k| \Phi(f_{n_k}, b) \leq \|\eta\|_{l^\infty} \sum_{k=1}^N \Phi(f_{n_k}, b) + \sup_{k > N} |\eta_k| \sum_{k=N+1}^{\infty} \Phi(f_{n_k}, b) \\ &\lesssim \|\eta\|_{l^\infty} \sum_{k=1}^N \Phi(f_{n_k}, b) + \sup_{k > N} |\eta_k|. \end{aligned}$$

Because $f_{n_k} \in \mathcal{D}_{\lambda,K}^0$ for each $k \geq 1$, we get

$$\overline{\lim}_{|b| \rightarrow 1} \Phi(Q(\eta), b) \lesssim \sup_{k > N} |\eta_k|.$$

Letting $N \rightarrow \infty$, we have that

$$\overline{\lim}_{|b| \rightarrow 1} \Phi(Q(\eta), b) = 0$$

and hence $Q(\eta) \in \mathcal{D}_{\lambda,K}^0$, which implies that $Q: \mathcal{C}_0 \rightarrow \mathcal{D}_{\lambda,K}^0$ is bounded.

Finally we will prove that $Q: \mathcal{C}_0 \rightarrow \mathcal{D}_{\lambda,K}^0$ is bounded below. For any $\eta = \{\eta_k\}$ and $j \geq 1$, by triangle inequality we obtain

$$\begin{aligned} \|Q(\eta)\|_{\mathcal{D}_{\lambda,K}} &\geq \Phi(Q(\eta), b_j) \geq |\eta_j| \Phi(f_{n_j}, b_j) - \sum_{k \neq j} |\eta_k| \Phi(f_{n_k}, b_j) \\ &\geq \frac{2S_1}{3} |\eta_j| - \sum_{k \neq j} |\eta_k| 3^{-k-1} S_1 \\ &\geq \frac{2S_1}{3} |\eta_j| - \frac{S_1}{6} \|\eta\|_{l^\infty}. \end{aligned}$$

The arbitrariness of j gives that

$$\|Q(\eta)\|_{\mathcal{D}_{\lambda,K}} \gtrsim \frac{S_1}{2} \|\eta\|_{l^\infty}.$$

Thus $Q: \mathcal{C}_0 \rightarrow \mathcal{D}_{\lambda,K}^0$ is an isomorphism onto its range. \square

THEOREM 3.6. *Let $-1 < \lambda < 0$ and K satisfy (2.1) for some $\delta \in (0, 1 + \lambda]$. Suppose $g \in F(1, -1, \lambda + 1) \setminus F_0(1, -1, \lambda + 1)$. Then there is a subspace $M \subset \mathcal{D}_{\lambda, K}$ isomorphic to \mathcal{C}_0 such that $T_g|_M : M \rightarrow T_g(M)$ is an isomorphism. In particular, T_g is not strictly singular.*

Proof. Let

$$\Upsilon := \limsup_{|I| \rightarrow 0} \frac{1}{|I|^{1+\lambda}} \int_{S(I)} |g'(w)|(1 - |w|^2)^\lambda dA(w).$$

Because $g \in F(1, -1, \lambda + 1) \setminus F_0(1, -1, \lambda + 1)$, we get that $\Upsilon > 0$ and there exists $\{I_n\} \subset \partial\mathbb{D}$ with $\lim_{n \rightarrow \infty} |I_n| = 0$ such that

$$\Upsilon = \lim_{n \rightarrow \infty} \frac{1}{|I_n|^{1+\lambda}} \int_{S(I_n)} |g'(w)|(1 - |w|^2)^\lambda dA(w) > 0. \quad (3.4)$$

For each $n \geq 1$, set $\gamma_n = (1 - |I_n|)\eta_n$, where η_n is the center of I_n . It is known that

$$|1 - \bar{\gamma}_n w| \approx 1 - |\gamma_n|^2 \approx |I_n|, \quad w \in S(I_n). \quad (3.5)$$

Take

$$f_n(w) = \frac{(1 - |\gamma_n|^2)^{\lambda+1} K(1 - |\gamma_n|^2)}{(1 - \bar{\gamma}_n w)^{2\lambda+2}}, \quad w \in \mathbb{D}.$$

It is easy to check that $f_n \in \mathcal{D}_{\lambda, K}^0$ and $\|f_n\|_{\mathcal{D}_{\lambda, K}} \lesssim 1$ for any $n \geq 1$. From (3.5), it is easy to see that

$$|f_n(w)| \gtrsim \frac{K(|I_n|)}{|I_n|^{1+\lambda}}, \quad w \in S(I_n). \quad (3.6)$$

Then by using (3.4) and (3.6), there exist some constant $\beta > 0$ such that

$$\frac{1}{K(|I_n|)} \int_{S(I_n)} |f_n(w)| |g'(w)|(1 - |w|^2)^\lambda dA(w) \geq \frac{\Upsilon\beta}{3}$$

for n larger than some fixed positive number $N \geq 1$.

Since $g \in F(1, -1, \lambda + 1)$, T_g is bounded on $\mathcal{D}_{\lambda, K}$, which implies that $T_g f_n \in \mathcal{D}_{\lambda, K}$ and

$$\|T_g f_n\|_{\mathcal{D}_{\lambda, K}} \lesssim 1.$$

Moreover, by Corollary 3.2, T_g is bounded also on $\mathcal{D}_{\lambda, K}^0$ and then $T_g f_n \in \mathcal{D}_{\lambda, K}^0$. Thus, passing to a further subsequence if necessary, we assume that

$$\frac{1}{K(|I_{n+1}|)} \int_{S(I_{n+1})} |f_n(w)| |g'(w)|(1 - |w|^2)^\lambda dA(w) \leq \frac{\Upsilon\beta}{4}$$

for any $n \geq N$.

Set

$$H_n = f_{n+1} - f_n, n \geq N.$$

Then $H_n \in \mathcal{D}_{\lambda,K}^0$ and $\|H_n\|_{\mathcal{D}_{\lambda,K}} \lesssim 1$. According to the above facts, we have that

$$\begin{aligned} & \frac{1}{K(|I_{n+1}|)} \int_{S(I_{n+1})} |H_n(z)| |g'(w)| (1 - |w|^2)^\lambda dA(w) \\ &= \frac{1}{K(|I_{n+1}|)} \int_{S(I_{n+1})} |f_{n+1}(w) - f_n(w)| |g'(w)| (1 - |w|^2)^\lambda dA(w) \\ &\geq \frac{1}{K(|I_{n+1}|)} \int_{S(I_{n+1})} |f_{n+1}(w)| |g'(w)| (1 - |w|^2)^\lambda dA(w) \\ &\quad - \frac{1}{K(|I_{n+1}|)} \int_{S(I_{n+1})} |f_n(w)| |g'(w)| (1 - |w|^2)^\lambda dA(w) \\ &\geq \frac{\Upsilon\beta}{12}. \end{aligned}$$

Therefore, for any $n \geq N$,

$$\begin{aligned} \frac{\Upsilon\beta}{12} &\leq \frac{1}{K(|I_{n+1}|)} \int_{S(I_{n+1})} |H_n(w)| |g'(w)| (1 - |w|^2)^\lambda dA(w) \\ &\lesssim \|T_g H_n\|_{*, \mathcal{D}_{\lambda,K}} \lesssim \|H_n\|_{*, \mathcal{D}_{\lambda,K}} \lesssim 1. \end{aligned}$$

Thus $T_g H_n \in \mathcal{D}_{\lambda,K}^0$ and

$$\|T_g H_n\|_{*, \mathcal{D}_{\lambda,K}} \asymp \|H_n\|_{*, \mathcal{D}_{\lambda,K}} \asymp 1$$

for any $n \geq N$.

Since $H_n \in \mathcal{D}_{\lambda,K}^0$, we have that $H_n \in \mathcal{D}_\lambda^1$. Set $P_n^\lambda = (1 - |\gamma_n|^2)^{\lambda+1} K(1 - |\gamma_n|^2)$.

$$\begin{aligned} & \int_{\mathbb{D}} |H'_n(w)| (1 - |w|^2)^\lambda dA(w) = \int_{\mathbb{D}} |f'_{n+1}(w) - f'_n(w)| (1 - |w|^2)^\lambda dA(w) \\ &= (2\lambda + 2) \int_{\mathbb{D}} \left| \frac{P_{n+1}^\lambda \gamma_{n+1}}{(1 - \overline{\gamma_{n+1}} w)^{2\lambda+3}} - \frac{P_n^\lambda \gamma_n}{(1 - \overline{\gamma_n} w)^{2\lambda+3}} \right| (1 - |w|^2)^\lambda dA(w) \\ &\lesssim \int_{\mathbb{D}} \frac{|P_{n+1}^\lambda \gamma_{n+1} (1 - \overline{\gamma_n} w)^{2\lambda+3} - P_n^\lambda \gamma_n (1 - \overline{\gamma_{n+1}} w)^{2\lambda+3}|}{|1 - \overline{\gamma_{n+1}} w|^{2\lambda+3} |1 - \overline{\gamma_n} w|^{2\lambda+3}} (1 - |w|^2)^\lambda dA(w) \\ &\lesssim \int_{\mathbb{D}} \frac{P_{n+1}^\lambda |(1 - \overline{\gamma_n} w)^{2\lambda+3} - (1 - \overline{\gamma_{n+1}} w)^{2\lambda+3}|}{|1 - \overline{\gamma_{n+1}} w|^{2\lambda+3} |1 - \overline{\gamma_n} w|^{2\lambda+3}} (1 - |w|^2)^\lambda dA(w) \\ &\quad + \int_{\mathbb{D}} \frac{|P_{n+1}^\lambda \gamma_{n+1} - P_n^\lambda \gamma_n| |1 - \overline{\gamma_{n+1}} w|^{2\lambda+3}}{|1 - \overline{\gamma_{n+1}} w|^{2\lambda+3} |1 - \overline{\gamma_n} w|^{2\lambda+3}} (1 - |w|^2)^\lambda dA(w) \\ &\lesssim \int_{\mathbb{D}} \frac{P_{n+1}^\lambda (1 - |w|^2)^\lambda dA(w)}{|1 - \overline{\gamma_{n+1}} w|^{2\lambda+3}} + \int_{\mathbb{D}} \frac{P_{n+1}^\lambda (1 - |w|^2)^\lambda dA(w)}{|1 - \overline{\gamma_n} w|^{2\lambda+3}} + \int_{\mathbb{D}} \frac{P_n^\lambda (1 - |w|^2)^\lambda dA(w)}{|1 - \overline{\gamma_n} w|^{2\lambda+3}}. \end{aligned} \tag{3.7}$$

By Lemma 2.5,

$$\begin{aligned} & \int_{\mathbb{D}} \frac{P_n^\lambda (1 - |w|^2)^\lambda dA(w)}{|1 - \overline{\gamma}_n w|^{2\lambda+3}} \\ &= (1 - |\gamma_n|^2)^{\lambda+1} K(1 - |\gamma_n|^2) \int_{\mathbb{D}} \frac{(1 - |w|^2)^\lambda dA(w)}{|1 - \overline{\gamma}_n w|^{2\lambda+3}} \lesssim K(1 - |\gamma_n|^2) \rightarrow 0 \end{aligned} \quad (3.8)$$

as $n \rightarrow \infty$. Similarly,

$$\int_{\mathbb{D}} \frac{P_{n+1}^\lambda (1 - |w|^2)^\lambda dA(w)}{|1 - \overline{\gamma}_{n+1} w|^{2\lambda+3}} \rightarrow 0 \quad \text{and} \quad \int_{\mathbb{D}} \frac{P_{n+1}^\lambda (1 - |w|^2)^\lambda dA(w)}{|1 - \overline{\gamma}_n w|^{2\lambda+3}} \rightarrow 0 \quad (3.9)$$

as $n \rightarrow \infty$. Combining (3.7), (3.8) and (3.9), we obtain that

$$\lim_{n \rightarrow \infty} \|H_n\|_{\mathcal{D}_\lambda^1} = 0.$$

Since T_g is bounded on \mathcal{D}_λ^1 by Corollary 3.4, we get

$$\lim_{n \rightarrow \infty} \|T_g H_n\|_{\mathcal{D}_\lambda^1} = 0.$$

Using Proposition 3.5, we can construct a subsequence $\{H_{n_k}\}$ of $\{H_n\}$ such that $Q : \mathcal{C}_0 \rightarrow \mathcal{D}_{\lambda,K}^0$ and $V : \mathcal{C}_0 \rightarrow \mathcal{D}_{\lambda,K}^0$ are both isomorphism onto their respective ranges, where

$$Q(\eta) = \sum_{k=1}^{\infty} \eta_k H_{n_k}, \quad \eta = \{\eta_k\} \subset \mathcal{C}_0,$$

and

$$V(\eta) = \sum_{k=1}^{\infty} \eta_k T_g H_{n_k}, \quad \eta = \{\eta_k\} \subset \mathcal{C}_0.$$

Let M be the closure of span $\{H_{n_k}\}$ in $\mathcal{D}_{\lambda,K}^0$. Then M is isomorphic to \mathcal{C}_0 and $T_g|_M : M \rightarrow T_g(M)$ is an isomorphism. This complete the proof of this theorem. \square

4. Closed range of T_g

In this section, we will study the closed range property of T_g on $\mathcal{D}_{\lambda,k}$ and show that $T_g : \mathcal{D}_{\lambda,k} \rightarrow \mathcal{D}_{\lambda,k}$ does not have closed range.

THEOREM 4.1. *Let $-1 < \lambda < 0$, $g \in F(1, -1, \lambda + 1)$ and K satisfy (2.1) for some $\delta \in (0, 1 + \lambda]$. Then $T_g : \mathcal{D}_{\lambda,k} \rightarrow \mathcal{D}_{\lambda,k}$ does not have closed range, or equivalently to say that T_g is not bounded below on $\mathcal{D}_{\lambda,k}$.*

Proof. If $g \in F_0(1, -1, 1)$, by Lemma 3.1, we have that T_g is compact on $\mathcal{D}_{\lambda,k}$ and then T_g is not bounded below on $\mathcal{D}_{\lambda,k}$. Thus we just need to prove that the conclusion is correct when $g \in F(1, -1, \lambda + 1) \setminus F_0(1, -1, \lambda + 1)$ happens.

From the proof of Theorem 2.9, for any $\varepsilon > 0$, there is a positive constant $\rho_0 \in (0, 1)$ such that

$$\frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} < \varepsilon \quad (4.1)$$

for any $\rho_0 < |b| < 1$.

Let $f_n(w) = w^n$. Clearly, $f_n \in \mathcal{D}_{\lambda,k}$ for any $n \geq 1$. Then, for any $n \geq 1$, by (4.1) we get that

$$\begin{aligned} & \sup_{\rho_0 < |b| < 1} \frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} \int_{\mathbb{D}} |w^n| |g'(w)| (1 - |w|^2)^{-1} (1 - |\sigma_b(w)|^2)^{2\lambda+2} dA(w) \\ & \leq \|g\|_{F(1,-1,\lambda+1)} \sup_{\rho_0 < |b| < 1} \frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} \lesssim \varepsilon. \end{aligned} \quad (4.2)$$

Now let us prove the case of $|b| \leq \rho_0$. Since $g \in F(1, -1, \lambda + 1)$, we see that $g \in \mathcal{D}_\lambda^1$. Thus there exists $r \in (0, 1)$ such that

$$\int_{\mathbb{D} \setminus r\mathbb{D}} |g'(w)| (1 - |w|^2)^\lambda dA(w) < \varepsilon.$$

Thus, by the monotonicity of K we have that

$$\begin{aligned} & \sup_{|b| \leq \rho_0} \frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} \int_{\mathbb{D} \setminus r\mathbb{D}} |w^n| |g'(w)| (1 - |w|^2)^{-1} (1 - |\sigma_b(w)|^2)^{\lambda+1} dA(w) \\ & = \sup_{|b| \leq \rho_0} \frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} \int_{\mathbb{D} \setminus r\mathbb{D}} |w^n| |g'(w)| (1 - |w|^2)^\lambda \frac{1}{|1 - \bar{b}w|^{2\lambda+2}} dA(w) \\ & \lesssim \sup_{|b| \leq \rho_0} \frac{1}{K(1 - |b|^2)} \int_{\mathbb{D} \setminus r\mathbb{D}} |g'(w)| (1 - |w|^2)^\lambda dA(w) \\ & \lesssim \frac{\varepsilon}{K(1 - |\rho_0|^2)}. \end{aligned} \quad (4.3)$$

In addition, we can choose a sufficiently large N such that $r^n < \varepsilon$ whenever $n > N$. Then

$$\begin{aligned} & \sup_{|b| \leq \rho_0} \frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} \int_{r\mathbb{D}} |w^n| |g'(w)| (1 - |w|^2)^{-1} (1 - |\sigma_b(w)|^2)^{\lambda+1} dA(w) \\ & \lesssim \frac{\varepsilon \|g\|_{F(1,-1,\lambda+1)}}{K(1 - |\rho_0|^2)}. \end{aligned} \quad (4.4)$$

Combing (4.3) with (4.4) we get

$$\begin{aligned} & \sup_{|b| \leq \rho_0} \frac{(1 - |b|^2)^{\lambda+1}}{K(1 - |b|^2)} \int_{\mathbb{D}} |w^n| |g'(w)| (1 - |w|^2)^{-1} (1 - |\sigma_b(w)|^2)^{\lambda+1} dA(w) \\ & \lesssim \frac{\varepsilon \|g\|_{F(1,-1,\lambda+1)}}{K(1 - |\rho_0|^2)} \lesssim \varepsilon. \end{aligned} \quad (4.5)$$

By (4.2), (4.5) and the fact that $\varepsilon > 0$ is arbitrary, we conclude that T_g is not bounded below on $\mathcal{D}_{\lambda,k}$. \square

Acknowledgements. The author was supported by GuangDong Basic and Applied Basic Research Foundation (No. 2023A1515010614).

Data availability. Data sharing is not applicable for this article as no datasets were generated or analyzed during the current study.

Conflict of interest. The author declare no competing interests.

REFERENCES

- [1] A. ALEMAN AND A. SISKAKIS, *An integral operator on H^p* , Complex Var. Theory Appl. **28** (1995), 149–158.
- [2] A. ALEMAN AND A. SISKAKIS, *Integration operators on Bergman spaces*, Indiana Univ. Math. J. **46** (1997), 337–356.
- [3] A. ANDERSON, *Some closed range integral operators on spaces of analytic functions*, Integral Equations Operator Theory **69** (2011), 87–99.
- [4] A. ANDERSON, M. JOVOCIC AND W. SMITH, *Some integral operators acting on H^∞* , Integral Equations Operator Theory **80** (2014), 275–291.
- [5] K. AVETISYAN AND S. STEVIĆ, *Extended Cesàro operators between different Hardy spaces*, Appl. Math. Comput. **207** (2009), 346–350.
- [6] D. C. CHANG, S. LI AND S. STEVIĆ, *On some integral operators on the unit polydisk and the unit ball*, Taiwanese J. Math. **11** (5) (2007), 1251–1285.
- [7] J. CHEN, *Some properties of the integration operators on the spaces $F(p, q, s)$* , Acta Math. Sci. **44 B** (2024), 173–188.
- [8] L. HU AND X. LIU, *Volterra integral operators on a family of Dirichlet-Morrey spaces*, Opuscula Math. **43** (2023), no. 5, 633–649.
- [9] L. HU, R. YANG AND S. LI, *Dirichlet-Morrey type spaces and Volterra integral operators*, J. Non-linear Var. Anal. **5** (2021), 477–491.
- [10] T. KATO, *Perturbation theory for nullity, deficiency and other quantities of linear operators*, J. Anal. Math. (1958), 261–322.
- [11] S. LI, J. LIU AND C. YUAN, *Embedding theorems for Dirichlet type spaces*, Canad. Math. Bull. **63** (2020), 106–117.
- [12] S. LI AND S. STEVIĆ, *Integral type operators from mixed-norm spaces to α -Bloch spaces*, Integral Transforms Spec. Funct. **18** (7) (2007), 485–493.
- [13] S. LI AND S. STEVIĆ, *Riemann-Stieltjes operators on Hardy spaces in the unit ball of \mathbb{C}^n* , Bull. Belg. Math. Soc. Simon Stevin **14** (4) (2007), 621–628.
- [14] S. LI AND S. STEVIĆ, *Compactness of Riemann-Stieltjes operators between $F(p, q, s)$ spaces and α -Bloch spaces*, Publ. Math. Debrecen **72** (2008), 111–128.
- [15] S. LI AND S. STEVIĆ, *Riemann-Stieltjes operators between different weighted Bergman spaces*, Bull. Belg. Math. Soc. Simon Stevin **15** (4) (2008), 677–686.
- [16] Q. LIN, J. LIU AND Y. WU, *Volterra type operators on $S^p(\mathbb{D})$ spaces*, J. Math. Anal. Appl. **461** (2018), 1100–1114.
- [17] Q. LIN, J. LIU AND Y. WU, *Strict singularity of Volterra type operators on Hardy spaces*, J. Math. Anal. Appl. **492** (2020), 124438.
- [18] J. LIU, Z. LOU AND K. ZHU, *Embedding of Möbius invariant function spaces into tent spaces*, J. Geom. Anal. **27** (2017), 1013–1028.
- [19] B. MACCLUER AND R. ZHAO, *Vanishing logarithmic Carleson measures*, Illinois J. Math. **46** (2002), 507–518.
- [20] S. MIIHKINEN, *Strict singularity of a Volterra-type integral operator on H^p* , Proc. Amer. Math. Soc. **145** (2017), 165–175.

- [21] J. PAU AND R. ZHAO, *Carleson measures, Riemann-Stieltjes and multiplication operators on a general family of function spaces*, Integral Equations Operator Theory **78** (2014), 483–514.
- [22] C. POMMERENKE, *Schlichte funktionen und analytische funktionen von beschränkter mittlerer Oszillation*, Comment. Math. Helv. **52** (1977), 591–602.
- [23] D. QU AND X. ZHU, *Embedding theorem for Besov-Morrey type spaces and Volterra integral operators*, J. Math. Inequal. **16** (2022), 1661–1674.
- [24] B. SEHBA AND S. STEVIĆ, *On some product-type operators from Hardy-Orlicz and Bergman-Orlicz spaces to weighted-type spaces*, Appl. Math. Comput. **233** (2014), 565–581.
- [25] C. SHEN, Z. LOU AND S. LI, *Embedding of $BMOA_{\log}$ into tent spaces and Volterra integral operators*, Comput. Meth. Funct. Theory **20** (2020), 217–234.
- [26] Y. SHI AND S. LI, *Essential norm of integral operators on Morrey type spaces*, Math. Inequal. Appl. **19** (2016), 385–393.
- [27] S. STEVIĆ, *Boundedness and compactness of an integral operator on a weighted space on the polydisc*, Indian J. Pure Appl. Math. **37** (6) (2006), 343–355.
- [28] S. STEVIĆ, *Norms of some operators from Bergman spaces to weighted and Bloch-type space*, Util. Math. **76** (2008), 59–64.
- [29] S. STEVIĆ, *Integral-type operators from a mixed norm space to a Bloch-type space on the unit ball*, Sib. Math. J. **50** (6) (2009), 1098–1105.
- [30] S. STEVIĆ, *On an integral-type operator from Zygmund-type spaces to mixed-norm spaces on the unit ball*, Abstr. Appl. Anal. **2010**, Article ID 198608, (2010), 7 pp.
- [31] S. STEVIĆ, *On operator P_ϕ^g from the logarithmic Bloch-type space to the mixed-norm space on unit ball*, Appl. Math. Comput. **215** (2010), 4248–4255.
- [32] S. STEVIĆ, *On some integral-type operators between a general space and Bloch-type spaces*, Appl. Math. Comput. **218** (2011), 2600–2618.
- [33] S. STEVIĆ, *Boundedness and compactness of an integral-type operator from Bloch-type spaces with normal weights to $F(p, q, s)$ space*, Appl. Math. Comput. **218** (2012), 5414–5421.
- [34] S. STEVIĆ AND S. I. UEKI, *Integral-type operators acting between weighted-type spaces on the unit ball*, Appl. Math. Comput. **215** (2009), 2464–2471.
- [35] F. SUN AND H. WULAN, *Characterizations of Morrey type spaces*, Canad. Math. Bull. **65** (2022), 328–344.
- [36] H. WULAN AND J. ZHOU, *Q_K and Morrey type spaces*, Ann. Acad. Sci. Fenn. Math. **38** (2013), 193–207.
- [37] H. WULAN AND K. ZHU, *Möbius Invariant Q_K Spaces*, Springer, Cham, (2017).
- [38] Y. YANG AND J. LIU, *Integral operators on Bergman-Morrey spaces*, J. Geom. Anal. **32** (2022), 181.
- [39] R. YANG AND X. ZHU, *Besov-Morrey spaces and Volterra integral operator*, Math. Inequal. Appl. **24** (2021), 857–871.
- [40] Y. ZHANG AND X. ZHU, *Volterra integral operator on Dirichlet-Morrey type spaces*, Complex Var. Ellipt. Equ., doi:10.1080/17476933.2024.2426154.
- [41] R. ZHAO, *On a general family of function spaces*, Ann. Acad. Sci. Fenn. Math. Diss. **105** (1996), 56.
- [42] J. ZHOU AND X. ZHU, *Essential norm of a Volterra-type integral operator from Hardy spaces to some analytic function spaces*, J. Integral Equ. Appl. **28** (2016), 581–593.
- [43] K. ZHU, *Operator Theory in Function Spaces*, 2nd edn, American Mathematical Society, Providence, 2007.
- [44] X. ZHU, E. ABBASI AND D. QU, *Bergman-Morrey type spaces and Volterra integral operators*, J. Nonlinear Var. Anal. **6** (2022), 359–370.
- [45] X. ZHU, L. HU AND D. QU, *Dirichlet-Morrey type spaces and Volterra integral operators*, Complex Var. Ellipt. Equ. **69** (2024), 301–316.

(Received May 4, 2025)

Xiangling Zhu
University of Electronic Science and Technology of China
Zhongshan Institute
528402, Zhongshan, Guangdong, P. R. China
e-mail: jyuzxl@163.com