Mathematical
Inequalities
& Applications
Volume 1, Number 1 (1998), 113–121

SOME HILBERT SPACE CHARACTERIZATIONS AND BANACH SPACE INEQUALITIES

TAE-HWA KIM * AND HONG-KUN XU **

(*communicated by J. Pečarić*)

Abstract. It is well known that if X is a normed linear space with dimension not less than three such that the radial projection from X onto the closed unit ball is nonexpansive, then X must be an inner product space. Using this fact, we are able to give a characterization of Hilbert spaces. Two other Hilbert space characterizations and some Banach space inequalities are established via duality maps.

Let X be a real normed linear space. We denote by B the closed unit ball of X and by P the radial projection from X onto B, i.e.,

$$Px = \begin{cases} x & \text{if } x \in B, \\ \dfrac{x}{\|x\|} & \text{if } x \notin B. \end{cases} \tag{1}$$

The following characterization of inner product spaces will be used in order to prove our first result.

LEMMA [4]. *A normed linear space $(X, \|\cdot\|)$ with dimension not less than three is an inner product space if and only if the radial projection $P : X \to B$ given by (1) is nonexpansive, i.e.,*

$$\|Px - Py\| \leqslant \|x - y\|, \text{ for all } x, y \in X.$$

In [7], Kim and Shin proved the following result.

THEOREM. *Let X be a uniformly convex Banach space satisfying Condition A (see definition below), C a nonempty closed convex subset of X and $f : C \to X$ a nonexpansive mapping (i.e., $\|f(x) - f(x)\| \leqslant \|x - y\|$, $x, y \in C$) such that $f(C)$ is bounded and $f(\partial C) \subseteq C$, where ∂C is the boundary of C. For each $k \in [0, 1)$, let $x_k \in C$ be the unique fixed point of the contraction $T_k : C \to X$ defined by*

$$T_k x = kTx + (1 - k)y, \quad x \in C,$$

where y is a fixed point in C. Then the strong $\lim_{k\uparrow 1} x_k$ exists and is a fixed point of T.

Mathematics subject classification (1991): Primary 46B20; Secondary 47H09.

Key words and phrases: Radial projection, nonexpansive mapping, condition A, duality map, uniformly smooth Banach space, inequality.

* Supported in part by BSRI-97-1440 and the Research Fund of Pukyong National University, 1996.

** Supported in part by FRD.

© ELEMENT, Zagreb
Paper MIA-01-09

DEFINITION [7]. A normed linear space $(X, \|\cdot\|)$ is said to satisfy Condition A if given $r > 0$, $y \in S_r := \{x \in X : \|x\| = r\}$ and $w \notin B_r := \{x \in X : \|x\| \leqslant r\}$ we have

$$\|y - w\| < \|y - \lambda w\| \quad \text{for all } \lambda > 1. \tag{2}$$

Clearly Condition (A) is equivalent to the validity of the inequality (2) for all $y, w \in X$ such that $\|y\| = 1$ and $\|w\| > 1$. In this note we shall prove that Condition A in fact characterizes inner product spaces with dimension not less than three; hence the above Theorem is reduced precisely to the Theorem of Singh and Watson [10] (see [1], [6], [12] for more and latest results about the strong convergence of $\{x_k\}$). Two more characterizations of Hibert spaces and certain inequalities in Banach spaces will also be obtained via the duality map (see below for definition).

PROPOSITION 1. *A normed linear space $(X, \|\cdot\|)$ satisfies Condition A if and only if $(X, \|\cdot\|)$ is an inner product space.*

Proof. It is easily seen that an inner product space satisfies Condition A. To show the converse assertion, by the Lemma above, it suffices to show that the radial projection P given by (1) is nonexpansive. Towards this end, we distinguish two cases.

Case 1: $\|x\| < 1$, $\|y\| > 1$. It follows from Condition A (with $r := \|x\|$) that

$$\begin{aligned}\|Px - Py\| &= \left\|x - \frac{y}{\|y\|}\right\| < \left\|x - \frac{\lambda}{\|y\|}y\right\| \qquad (\forall \lambda > 1)\\ &= \|x - y\| \qquad (\text{let } \lambda = \|y\|).\end{aligned}$$

Case 2: $\|x\| \geqslant 1$, $\|y\| \geqslant 1$. Without loss of generality we assume $\|y\| \geqslant \|x\|$. For any $\varepsilon > 0$ and $\lambda > 1$, Condition A yields that

$$\begin{aligned}\|Px - (1+\varepsilon)Py\| &= \left\|\frac{x}{\|x\|} - (1+\varepsilon)\frac{y}{\|y\|}\right\|\\ &< \left\|\frac{x}{\|x\|} - \lambda(1+\varepsilon)\frac{y}{\|y\|}\right\|\\ &= \frac{1}{\|x\|}\left\|x - \lambda(1+\varepsilon)\frac{\|x\|}{\|y\|}y\right\|\\ &\leqslant \left\|x - \lambda(1+\varepsilon)\frac{\|x\|}{\|y\|}y\right\| \quad \text{as } \|x\| \geqslant 1.\end{aligned}$$

Taking $\lambda = (1+\varepsilon)\|y\|/\|x\|$ and then letting $\varepsilon \to 0$ we get $\|Px - Py\| \leqslant \|x - y\|$. Therefore P is nonexpansive. $\square$

REMARK. The above proof shows that the conclusion of Proposition 1 remains valid if the strict "$<$" in (2) is replaced by the nonstrict "$\leqslant$", i.e., $\|y - x\| \leqslant \|y - \lambda w\|$ for all $y \in B$, $w \notin B$, and $\lambda > 1$.

Recall that the (normalized) *duality* map $J : X \to X^*$ is defined by

$$J(x) = \{x^* \in X^* : \langle x, x^*\rangle = \|x\|^2 = \|x^*\|^2\}, \quad x \in X.$$

It is known (cf. [5]) that X is *smooth* if and only if J is single-valued on X.

In [8] and [9], Park studied the convergence of the Mann iteration for accretive operators by imposing the following assumption on the underlying smooth Banach space X:

$$\langle x-y, J(x)-J(y)\rangle \leqslant \|x-y\|^2 \quad \text{for all } x,y \in X. \tag{3}$$

Our next result shows that the assumption (3) also characterizes Hilbert spaces; thus reducing the main results of [8] and [9] to a Hilbert space setting which have been proved already (cf. Bruck [2]). We shall work in a more general setting; i.e., we do not assume smoothness, instead we use the following assumption:

$$\langle x-y, j_x-j_y\rangle \leqslant \|x-y\|^2 \quad \forall j_x \in J(x),\ \forall j_y \in J(y),\ x,y \in X. \tag{4}$$

It is obvious that if X is smooth, then (4) is reduced to (3).

PROPOSITION 2. *Assume that a Banach space $(X, \|\cdot\|)$ satisfies the condition (4). Then X is a Hilbert space.*

Proof. Let $x,y \in X$ and $t \geqslant 0$ and let $j_{x+ty} \in J(x+ty)$ satisfy

$$\sup\{\langle y,j\rangle : j \in J(x+ty)\} = \langle y, j_{x+ty}\rangle .$$

Then we have for any $j_x \in J(x)$,

$$\begin{aligned}
\frac{d^+}{dt}\frac{1}{2}\|x+ty\|^2 &:= \lim_{s\downarrow 0} \frac{\frac{1}{2}\|x+ty+sy\|^2 - \frac{1}{2}\|x+ty\|^2}{s} \\
&= \sup\{\langle y,j\rangle : j \in J(x+ty)\} \\
&= \langle y, j_{x+ty}\rangle \\
&= \frac{1}{t}\langle (x+ty)-x, j_{x+ty}-j_x\rangle + \langle y, j_x\rangle \\
&\leqslant t\|y\|^2 + \langle y, j_x\rangle \quad \text{by (4).}
\end{aligned}$$

It follows that

$$\begin{aligned}
\frac{1}{2}\|x+y\|^2 &\leqslant \frac{1}{2}\|x\|^2 + \int_0^1 t\|y\|^2 dt + \langle y, j_x\rangle \\
&= \frac{1}{2}\|x\|^2 + \frac{1}{2}\|y\|^2 + \langle y, j_x\rangle ,
\end{aligned}$$

or

$$\|x+y\|^2 \leqslant \|x\|^2 + \|y\|^2 + 2\langle y, j_x\rangle , \quad x,y \in X. \tag{5}$$

Substitute $-y$ for y into (5) to get

$$\|x-y\|^2 \leqslant \|x\|^2 + \|y\|^2 - 2\langle y, j_x\rangle , \quad x,y \in X. \tag{6}$$

Adding (5) and (6) gets

$$\|x+y\|^2 + \|x-y\|^2 \leqslant 2(\|x\|^2 + \|y\|^2), \quad x,y \in X. \tag{7}$$

Replacing x by $\frac{x+y}{2}$ and y by $\frac{x-y}{2}$ in (7) respectively, we obtain

$$\|x+y\|^2 + \|x-y\|^2 \geqslant 2(\|x\|^2 + \|y\|^2), \quad x,y \in X. \tag{8}$$

By (7) and (8) we conclude that the Parallelogram identity

$$\|x+y\|^2 + \|x-y\|^2 = 2(\|x\|^2 + \|y\|^2) \tag{9}$$

holds for all $x, y \in X$; hence X is an inner product space. $\square$

COROLLARY 1. *A Banach space X is a Hilbert space if and only if the duality map J is Lipschitz with constant one in the sense that*

$$\|j_x - j_y\| \leqslant \|x-y\|, \quad \forall j_x \in J(x),\ \forall j_y \in J(y),\ x, y \in X.$$

Next we consider the dual version to Proposition 2; namely, we consider the case where there exists a selection j of J (i.e., a single-valued map $j : X \to X^*$ such that $j(x) \in J(x)$ for all $x \in X$) satisfying

$$\langle x-y, j(x)-j(y)\rangle \geqslant \|x-y\|^2, \quad x, y \in X. \tag{10}$$

We shall show the inequality (10) also characterizes inner product spaces.

PROPOSITION 3. *If a Banach space X satisfies the inequality* (10)*, then X is a Hilbert space.*

Proof. Since for every $x \in X$, $J(x) = \partial \frac{1}{2}\|x\|^2$, the subdifferential of the function $\frac{1}{2}\|\cdot\|^2$ at x, by the subdifferential inequality, we have for $x, y \in X$ and $t \geqslant 0$,

$$\begin{aligned}\frac{d^+}{dt}\frac{1}{2}\|x+ty\|^2 := \lim_{s\to t^+} \frac{\frac{1}{2}\|x+sy\|^2 - \frac{1}{2}\|x+ty\|^2}{s-t} \\ \geqslant \langle y, j(x+ty)\rangle .\end{aligned}$$

This combined with the inequality (10) yields

$$\begin{aligned}\frac{d^+}{dt}\frac{1}{2}\|x+ty\|^2 &\geqslant \frac{1}{t}\langle (x+ty)-x, j(x+ty)-j(x)\rangle + \langle y, j(x)\rangle \\ &\geqslant t\|y\|^2 + \langle y, j(x)\rangle .\end{aligned}$$

Hence

$$\begin{aligned}\frac{1}{2}\|x+y\|^2 &\geqslant \frac{1}{2}\|x\|^2 + \int_0^1 (t\|y\|^2 + \langle y, j(x)\rangle)dt \\ &= \frac{1}{2}\|x\|^2 + \frac{1}{2}\|y\|^2 + \langle y, j(x)\rangle .\end{aligned}$$

Replacing y by $-y$ leads to

$$\frac{1}{2}\|x-y\|^2 \geqslant \frac{1}{2}\|x\|^2 + \frac{1}{2}\|y\|^2 - \langle y, j(x)\rangle .$$

Adding these last two inequalities gets

$$\|x+y\|^2 + \|x-y\|^2 \geqslant 2(\|x\|^2 + \|y\|^2) \quad \forall x, y \in X.$$

This is inequality (8). Replacing in (8) x by $\frac{x+y}{2}$ and y by $\frac{x-y}{2}$, respectively yields inequality (7). Hence the Parallelogram indentity (9) holds for all $x, y \in X$ and X is an inner product space. $\square$

REMARK. In Proposition 3, we need that inequality (10) is valid only for some selection j for J; while in Proposition 2 we need the validity of inequality (4) for all $j_x \in J(x)$ and $j_y \in J(y)$. We do not know whether Proposition 2 is still valid if inequality (4) holds only for some selection j of J.

COROLLARY 2 [3]. *A normed linear space $(X, \|\cdot\|)$ is an inner product space if and only if the duality map $J : X \to X^*$ is linear, i.e.,*

$$J(ax+by) = aJ(x) + bJ(y) \quad x, y \in X \;\; a, b \in \mathbb{R}.$$

This means that given any $j_x \in J(x)$ and $j_y \in J(y)$, there exists some $j_{ax+by} \in J(ax+by)$ such that $aj_x + bj_y = j_{ax+by}$.

Proof. Given $x, y \in X$ and any selection j of J, we can find some $j_{x-y} \in J(x-y)$ satisfying $j(x) - j(y) = j_{x-y}$ so that

$$\langle x-y, j(x)-j(y)\rangle = \langle x-y, j_{x-y}\rangle = \|x-y\|^2.$$

So Corollary 2 follows from Proposition 3. $\square$

In the remaining part of this paper, we shall improve some Banach space inequalities established in [11]. Recall that the moduli of convexity and smoothness of a normed linear space X are defined respectively by

$$\delta_X(\varepsilon) = \inf\Big\{1 - \|\frac{1}{2}(x+y)\| : \|x\| = \|y\| = 1, \, , \|x-y\| = \varepsilon\Big\}, \quad 0 \leqslant \varepsilon \leqslant 2,$$

and

$$\rho_X(\tau) = \sup\Big\{\frac{1}{2}(\|x+y\| + \|x-y\|) - 1 : \|x\| = 1, \, , \|y\| = \tau\Big\}, \quad \tau > 0.$$

X is *uniformly convex* if $\delta_X(\varepsilon) > 0$ for $0 < \varepsilon \leqslant 2$ and *uniformly smooth* if $\lim\limits_{\tau \to 0} \frac{\rho_X(\tau)}{\tau} = 0$. Let $p, q > 1$ be real numbers. Then X is said to be *p-uniformly convex* (resp., *q-uniformly smooth*) if there is a constant $c > 0$ such that $\delta_X(\varepsilon) \geqslant c\varepsilon^p$, $0 \leqslant \varepsilon \leqslant 2$ (resp., $\rho_X(\tau) \leqslant c\tau^q$, $\tau > 0$). It is proved in [11] that X is p-uniformly convex if and only if there is a constant $c > 0$ such that for all $x, y \in X$ and $\lambda \in [0,1]$

$$\|\lambda x + (1-\lambda)y\|^p \leqslant \lambda\|x\|^p + (1-\lambda)\|y\|^p - W_p(\lambda)c\|x-y\|^p, \tag{11}$$

where

$$W_p(\lambda) = \lambda^p(1-\lambda) + \lambda(1-\lambda)^p.$$

It is also proved in [11] that if X is a smooth Banach space, then the dual version to (11), i.e., the inequality

$$\|\lambda x + (1-\lambda)y\|^q \geqslant \lambda\|x\|^q + (1-\lambda)\|y\|^q - W_q(\lambda)c\|x-y\|^q, \quad x, y \in X \text{ and } \lambda \in [0,1] \tag{12}$$

characterizes uniform smoothness. Here we are going to show that the smoothness assumption on X can be removed.

PROPOSITION 4. *Let $q > 1$ be a real number. Then a Banach space X is q-uniformly smooth if and only if there is a constant $c > 0$ such that inequality* (12) *holds.*

Proof. By [11, Theorem 1′], it suffices to show that (12) implies the smoothness of X. By replacing in (12) x by $\frac{x+y}{2}$ and y by $\frac{x-y}{2}$ and setting $\lambda = \frac{1}{2}$ we get

$$\frac{1}{2}(\|x+y\|^q + \|x-y\|^q) \leqslant \|x\|^q + c\|y\|^q, \quad x, y \in X. \tag{13}$$

Assume X is not smooth. Then there exists an $x_0 \in X$, $\|x_0\| = 1$, and $f, g \in X^*$, $f \neq g$, $\|f\| = \|g\| = 1$ such that $f(x_0) = g(x_0) = \|x_0\| = 1$. Take $y_0 \in X$, $\|y_0\| = 1$ such that $(f+g)(y_0) = 0$ and $f(y_0) > 0$. It follows that for any $\tau > 0$,

$$\begin{aligned}
\|x_0 + \tau y_0\|^q + \|x_0 - \tau y_0\|^q &\geqslant [f(x_0 + \tau y_0)]^q + [g(x_0 - \tau y_0)]^q \\
&= [1 + \tau f(y_0)]^q + [1 - \tau g(y_0)]^q \\
&= 2[1 + \tau f(y_0)]^q \qquad \text{as} \quad -g(y_0) = f(y_0) \\
&\geqslant 2[1 + \tau f(y_0)] \qquad \text{as} \quad f(y_0) > 0.
\end{aligned}$$

But inequality (13) then implies for all $\tau > 0$,

$$1 + c\tau^q \geqslant 1 + \tau f(y_0) \quad \text{or} \quad c\tau^{q-1} \geqslant f(y_0),$$

which is impossible since $\tau^{q-1} \to 0$ as $\tau \to 0^+$ and $f(y_0) > 0$. □

We now use the (generalized) *duality* map $J_q : X \to X^*$ defined by

$$J_q(x) = \{x^* \in X^* : \langle x, x^* \rangle = \|x\|^q \text{ and } \|x^*\| = \|x\|^{q-1}\}, \quad x \in X$$

to characterize uniform smoothness of Banach spaces. Again we remove the smoothness assumption in [11].

PROPOSITION 5. *Let $q > 1$ be a real number. Then the following are equivalent for a Banach space X.*

(*i*) *X is q-uniformly smooth.*

(*ii*) *There is a constant $c > 0$ such that*

$$\|x+y\|^q \leqslant \|x\|^q + q\langle y, j_q(x)\rangle + c\|y\|^q, \quad \forall x, y \in X,\ \forall j_q(x) \in J_q(x). \tag{14}$$

(*iii*) *There is a constant $\tilde{c} > 0$ such that*

$$\langle x-y, j_q(x) - j_q(y)\rangle \leqslant \tilde{c}\|x-y\|^q, \quad \forall x, y \in X,\ \forall j_q(x) \in J_q(x),\ \forall j_q(y) \in J_q(y). \tag{15}$$

Proof. (*i*) $\Rightarrow$ (*ii*). Since X is smooth, we have that J_q is single-valued and

$$q\langle y, J_q(x)\rangle = \lim_{\lambda \downarrow 0} \frac{\|x + \lambda y\|^q - \|x\|^q}{\lambda}, \quad x, y \in X.$$

By inequality (12), we have

$$\begin{aligned}\frac{\|x+\lambda y\|^q-\|x\|^q}{\lambda} &= \frac{\|(1-\lambda)x+\lambda(x+y)\|^q-\|x\|^q}{\lambda}\\ &\geqslant \frac{(1-\lambda)\|x\|^q+\lambda\|x+y\|^q-W_q(\lambda)c\|y\|^q-\|x\|^2}{\lambda}\\ &= \|x+y\|^q-\|x\|^q-[(1-\lambda)^q+\lambda^{q-1}(1-\lambda)]c\|y\|^q.\end{aligned}$$

Taking the limit as $\lambda\to 0^+$ we obtain

$$q\langle y,J_q(x)\rangle \geqslant \|x+y\|^q-\|x\|^q-c\|y\|^q,$$

which is (14).

$(ii)\Rightarrow(iii)$. By (14) we have for any $j_q(x)\in J_q(x)$ and $j_q(y)\in J_q(y)$,

$$\|y\|^q\leqslant\|x\|^q+q\langle y-x,j_q(x)\rangle+c\|y-x\|^q \tag{16}$$

and

$$\|x\|^q\leqslant\|y\|^q+q\langle x-y,j_q(y)\rangle+c\|x-y\|^q. \tag{17}$$

Upon adding (17) to (16) we get

$$\langle x-y,j_q(x)-j_q(y)\rangle\leqslant\frac{2c}{q}\|x-y\|^q.$$

Hence (15) is valid with $\tilde{c}=2c/q$.

$(iii)\Rightarrow(i)$. Let $j_q(x+ty)\in J_q(x+ty)$ satisfy

$$\langle y,j_q(x+ty)\rangle=\sup\{\langle y,j_q\rangle : j_q\in J_q(x+ty)\}.$$

It then follows that

$$\begin{aligned}\frac{d^+}{dt}\frac{1}{q}\|x+ty\|^q &:= \lim_{s\downarrow 0}\frac{\|x+ty+sy\|^q-\|x+ty\|^q}{qs}\\ &= \sup\{\langle y,j_q\rangle : j_q\in J_q(x+ty)\}\\ &= \langle y,j_q(x+ty)\rangle\\ &= \frac{1}{t}\langle (x+ty)-x,j_q(x+ty)-j_q(x)\rangle+\langle y,j_q(x)\rangle\\ &\leqslant \tilde{c}t^{q-1}\|y\|^q+\langle y,j_q(x)\rangle \qquad \text{by (15)}.\end{aligned}$$

Therefore,

$$\|x+y\|^q\leqslant\|x\|^q+q\int_0^1(\tilde{c}t^{q-1}\|y\|^q+\langle y,j_q(x)\rangle)dt,$$

or

$$\|x+y\|^q\leqslant\|x\|^q+\tilde{c}\|y\|^q+q\langle y,j_q(x)\rangle \quad \forall x,y\in X,\ \forall j_q(x)\in J_q(x). \tag{18}$$

Now for any $\lambda\in[0,1]$, write $z=\lambda x+(1-\lambda)y$. Then by (18) we have

$$\|x\|^q\leqslant\|z\|^q+\tilde{c}\|x-z\|^q+q\langle x-z,j_q(z)\rangle \tag{19}$$

and

$$\|y\|^q \leqslant \|z\|^q + \tilde{c}\|y-z\|^q + q\langle y-z, j_q(z)\rangle . \tag{20}$$

Since $x-z=(1-\lambda)(x-y)$ and $y-z=\lambda(y-x)$, it follows from (19) and (20) that

$$\lambda\|x\|^q + (1-\lambda)\|y\|^q \leqslant \|z\|^q + \tilde{c}[\lambda(1-\lambda)^q + \lambda^q(1-\lambda)]\|x-y\|^q.$$

Hence (12) holds with $c=\tilde{c}$ and X is q-uniformly smooth by Propositon 4. □

Finally we state the local versions of Propositoins 4 and 5 which again delete the smoothness assumption imposed on the space X in [11]. The proofs are omitted here as they are similar to those of Propositions 4 and 5. Recall that $B_r = \{x \in X : \|x\| \leqslant r\}$ is the closed ball centered at the origin with radius $r>0$. Let Γ be the family of continuous, strictly increasing and convex functions $g : [0,\infty) \to [0,\infty)$ such that $\lim_{\tau\to 0^+} g(\tau)/\tau = 0$.

PROPOSITION 6. *Let $q>1$ and $r>0$ be given. Then the following are equivalent for a Banach space X.*

(*i*) *X is uniformly smooth.*

(*ii*) *There exists a $g \in \Gamma$ (depending on r) such that*

$$\|\lambda x + (1-\lambda)y\|^q \geqslant \lambda\|x\|^q + (1-\lambda)\|y\|^q - W_q(\lambda)g(\|x-y\|) \quad \forall x,y \in B_r.$$

(*iii*) *There exists a $g \in \Gamma$ (depending on r) such that*

$$\|x+y\|^q \leqslant \|x\|^q + q\langle y, j_q(x)\rangle + g(\|y\|) \quad \forall x,y \in B_r,\ \forall j_q(x) \in J_q(x).$$

(*iv*) *There exists a $g \in \Gamma$ (depending on r) such that*

$$\langle x-y, j_q(x)-j_q(y)\rangle \leqslant g(\|x-y\|) \quad \forall x,y \in B_r,\ \forall j_q(x) \in J_q(x),\ \forall j_q(y) \in J_q(y).$$

Acknowledgement. This work was done while H. K. Xu was visiting Pukyong National University.

REFERENCES

[1] F. E. BROWDER, *Convergence of approximants to fixed points of nonexpansive nonlinear maps in Banach spaces*, Arch. Rational Mech. Acad. **24** (1967), 82–90.

[2] R. E. BRUCK, *The iterative solution of the equation $y \in x+Tx$ for a monotone operator T in Hilbert space*, Bull. Amer. Math. Soc. **79** (1973), 1258–1261.

[3] J. CIORANESCU, Geometry of Banach Spaces–Duality Mappings and Nonlinear Problems, Mathematics and Its Applications **62**, Kluwer Academic Publisher, Dordrecht, The Netherlands, 1990.

[4] D. G. DEFIGUEIREDO AND L. A. KARLOVITZ, *On the radial projection in normed spaces*, Bull. Amer. Math. Soc. **73** (1967), 364–368.

[5] J. DIESTEL, Geometry of Banach Spaces–Selected Topics, Lect. Notes in Mathematics **485**, Springer-Verlag, New-York, 1970.

[6] B. HARPERN, *Fixed points of nonexpanding maps*, Bull. Amer. Math. Soc. **73** (1967), 957–961.

[7] K. KIM AND S. S. SHIN, *On approximating fixed points in Banach spaces*, Preprint.

[8] J. A. PARK, *Mann-iteration process for the fixed point of strictly pseudocontractive mapping in some Banach spaces*, J. Korean Math. Soc. **31** (1994), 333–337.

[9] J. A. PARK, *Mann-iteration process to the solution of $y = x+Tx$ for an accretive operator T in some Banach spaces*, Comm. Korean Math. Soc. **9** (1994), 819–823.

[10] S. P. SINGH AND B. WATSON, *On approximating fixed points*, Proc. Symp. Pure Math. **45** (1986), Part 2, 393–395.
[11] H. K. XU, *Inequalities in Banach spaces with applications*, Nonlinear Analysis **16** (1991), 1127–1136.
[12] H. K. XU, *Approximating curves of nonexpansive nonself mappings in Banach spaces*, C. R. Acad. Sci., Paris, Ser. I, to appear.

(Received May 9, 1997)

Tae-Hwa Kim
Department of Applied Mathematics
Pukyong National University
Pusan 608-737
Korea
e-mail: taehwa@dolphin.pknu.ac.kr

Hong-Kun Xu
Department of Mathematics
University of Durban-Westville
Private Bag X54001
Durban 4000
South Africa
e-mail: hkxu@pixie.udw.ac.za