

UPPER AND LOWER SOLUTIONS METHOD FOR HIGHER ORDER DISCRETE BOUNDARY VALUE PROBLEMS

RAVI P. AGARWAL AND FU-HSIANG WONG

(communicated by J. Pečarić)

Abstract. We shall offer sufficient conditions on the function $f(i, u_1, \dots, u_{n-1})$, so that the higher order discrete boundary value problem

$$(BVP) \begin{cases} (E) \Delta^n u(i-1) + f(i, u(i), \Delta u(i), \dots, \Delta^{n-2} u(i)) = 0 \text{ for } i \in [1, T+1] \text{ and } n \geq 2, \\ (BC) \begin{cases} \Delta^m u(0) = 0, \quad 0 \leq m \leq n-3, \\ \alpha \Delta^{n-2} u(0) - \beta \Delta^{n-1} u(0) = 0, \\ \gamma \Delta^{n-2} u(T+1) + \delta \Delta^{n-1} u(T+1) = 0, \end{cases} \end{cases}$$

has at least one solution.

1. Introduction

In this paper we shall employ upper and lower solutions method to offer some new existence criteria for the higher order discrete boundary value problem

$$(BVP) \begin{cases} (E) \Delta^n u(i-1) + f(i, u(i), \Delta u(i), \dots, \Delta^{n-2} u(i)) = 0 \text{ for } i \in [1, T+1] \text{ and } n \geq 2, \\ (BC) \begin{cases} \Delta^m u(0) = 0, \quad 0 \leq m \leq n-3, \\ \alpha \Delta^{n-2} u(0) - \beta \Delta^{n-1} u(0) = 0, \\ \gamma \Delta^{n-2} u(T+1) + \delta \Delta^{n-1} u(T+1) = 0, \end{cases} \end{cases}$$

where $T \geq 1$ and $n \geq 2$ are fixed positive integers, Δ^m denotes the m -th forward difference operator with stepsize 1, and $[a, b] = \{a, a+1, \dots, b\} \subseteq \mathbf{Z}$ is the set of all integers.

The motivation for the present work stems from many recent investigations in [1–20, and the several references therein] for the above (BVP), and several of its particular cases.

Mathematics subject classification (1991): 34B15, 39A10.

Key words and phrases: Higher order discrete boundary value problem, Green's function, upper and lower solutions.

2. Main Results

Let $\alpha, \gamma, \beta, \delta \geq 0$, $\rho = \gamma\beta + \alpha\gamma(T+1) + \alpha\delta > 0$, and \mathbf{B} be the Banach space

$$\mathbf{B} = \{u : [0, T+n] \rightarrow \mathbb{R} \mid \Delta^m u(0) = 0, \ 0 \leq m \leq n-3\}$$

with the norm $\|u\| = \max_{i \in [0, T+2]} |\Delta^{n-2} u(i)|$. To abbreviate our discussion, we shall assume that the following conditions are satisfied:

(C₁) $K(i, j)$ defined on $[0, T+n] \times [1, T+1]$ is the Green's function of the difference equation

$$-\Delta^n u(i-1) = 0 \text{ for } i \in [1, T+1]$$

subject to the boundary conditions (BC).

(C₂) $k(i, j)$ defined on $[0, T+2] \times [1, T+1]$ is the Green's function of the difference equation

$$-\Delta^2 u(i-1) = 0 \text{ for } i \in [1, T+1]$$

subject to the boundary conditions

$$(BC^*) \quad \begin{cases} \alpha u(0) - \beta \Delta u(0) = 0, \\ \gamma u(T+1) + \delta \Delta u(T+1) = 0. \end{cases}$$

(C₃) $f \in C([1, T+1] \times \mathbb{R}^{n-1}; \mathbb{R})$.

(C₄) $v, w : [0, T+n] \rightarrow \mathbb{R}$ are lower and upper solutions of (BVP), i.e.,

$$\left\{ \begin{array}{l} (1) \quad \Delta^n v(i-1) + f(i, v(i), \Delta v(i), \dots, \Delta^{n-2} v(i)) \geq 0 \text{ for } i \in [1, T+1], \\ (2) \quad \Delta^n w(i-1) + f(i, w(i), \Delta w(i), \dots, \Delta^{n-2} w(i)) \leq 0 \text{ for } i \in [1, T+1], \\ (3) \quad \begin{cases} \Delta^m v(0) \leq 0, \ 0 \leq m \leq n-3, \\ \alpha \Delta^{n-2} v(0) - \beta \Delta^{n-1} v(0) \leq 0, \\ \gamma \Delta^{n-2} v(T+1) + \delta \Delta^{n-1} v(T+1) \leq 0, \end{cases} \\ (4) \quad \begin{cases} \Delta^m w(0) \geq 0, \ 0 \leq m \leq n-3, \\ \alpha \Delta^{n-2} w(0) - \beta \Delta^{n-1} w(0) \geq 0, \\ \gamma \Delta^{n-2} w(T+1) + \delta \Delta^{n-1} w(T+1) \geq 0. \end{cases} \end{array} \right.$$

(C₅) The lower and upper solutions $v(i)$ and $w(i)$, and the function $f(i, u_1, \dots, u_{n-2}, u_{n-1})$ satisfy the following:

(I) $\Delta^{n-2} v(i) \leq \Delta^{n-2} w(i)$ on $[0, T+1]$,

(II) $f(i, v(i), \dots, \Delta^{n-3} v(i), u_{n-1}) \leq f(i, u_1, \dots, u_{n-2}, u_{n-1})$
 $\leq f(i, w(i), \dots, \Delta^{n-3} w(i), u_{n-1})$, for $i \in [1, T+1]$, and $(v(i), \dots, \Delta^{n-3} v(i))$
 $\leq (u_1, \dots, u_{n-2}) \leq (w(i), \dots, \Delta^{n-3} w(i))$, where $(x_1, \dots, x_{n-2}) \leq (y_1, \dots, y_{n-2})$,
 if and only if, $x_m \leq y_m$ for $m = 1, \dots, n-2$.

REMARK 2.1. The following hold:

(a) If $f(i, u_1, \dots, u_{n-2}, u_{n-1})$ is increasing with respect to (u_1, \dots, u_{n-2}) on \mathbb{R}^{n-2} for each fixed $(i, u_{n-1}) \in [1, T+1] \times \mathbb{R}$, then (C₅) (II) holds.

(b) A simple calculation shows that

$$\Delta^{n-2}K(i,j) = k(i,j) = \begin{cases} \frac{1}{\rho}\{\beta + \alpha j\}\{\delta + \gamma(T+1-i)\}, & 1 \leq j \leq i-1, \\ \frac{1}{\rho}\{\beta + \alpha i\}\{\delta + \gamma(T+1-j)\}, & i \leq j \leq T+1. \end{cases}$$

(c) There are positive constants M and L such that

$$\begin{cases} (R_1) & \frac{k(i,j)}{k(j,j)} \leq L, \quad \text{for all } (i,j) \in [0, T+2] \times [1, T+1], \\ (R_2) & \frac{k(i,j)}{k(j,j)} \geq M, \quad \text{for all } (i,j) \in [1, T] \times [1, T+1]. \end{cases}$$

Our main result is embodied in the following:

THEOREM 1. *With respect to the (BVP) assume that conditions $(C_1) - (C_5)$ are satisfied. Then, (BVP) has at least one solution $u \in \mathbf{B}$ such that*

$$\Delta^m v(i) \leq \Delta^m u(i) \leq \Delta^m w(i) \quad \text{on } [0, T+n-m-1] \quad \text{for } m = 0, 1, \dots, n-2.$$

Proof. We separate the proof into the following steps:

Step (1) Consider the modified problem

$$(BVP^*) \quad \begin{cases} (E^*) & \Delta^n u(i-1) + f^*(i, u(i), \Delta u(i), \dots, \Delta^{n-2}u(i)) = 0 \quad \text{for } i \in [1, T+1], \\ (BC) & \begin{cases} \Delta^m u(0) = 0, & 0 \leq m \leq n-3, \\ \alpha \Delta^{n-2}u(0) - \beta \Delta^{n-1}u(0) = 0, \\ \gamma \Delta^{n-2}u(T+1) + \delta \Delta^{n-1}u(T+1) = 0, \end{cases} \end{cases}$$

where

$$f^*(i, u_1, \dots, u_{n-1}) = f(i, \eta_1, \dots, \eta_{n-1}) + \rho(\eta_{n-1} - u_{n-1}),$$

and

$$\eta_l = \begin{cases} \Delta^{l-1}w(i) & \text{if } u_l > \Delta^{l-1}w(i), \\ u_l & \text{if } \Delta^{l-1}v(i) \leq u_l \leq \Delta^{l-1}w(i), \\ \Delta^{l-1}v(i) & \text{if } u_l < \Delta^{l-1}v(i), \end{cases}$$

for all $l = 1, 2, \dots, n-1$, $i \in [1, T+1]$, and $\rho: \mathbf{R} \rightarrow [-1, 1]$ is the radial retraction defined by

$$\rho(r) = \begin{cases} r & \text{for } |r| \leq 1, \\ \frac{r}{|r|} & \text{for } |r| > 1. \end{cases}$$

It is clear that (BVP^*) has a solution $u = u(i)$, if and only if, u is a solution of the operator equation

$$u(i) = \sum_{j=1}^{T+1} K(i,j) f^*(j, u(j), \Delta u(j), \dots, \Delta^{n-2}u(j)) = (T^*u)(i), \quad u \in \mathbf{B}$$

for $i \in [0, T + n]$, or equivalently,

$$\Delta^{n-2}u(i) = \sum_{j=1}^{T+1} k(i, j)f^*(j, u(j), \Delta u(j), \dots, \Delta^{n-2}u(j)) = \Delta^{n-2}(T^*u)(i), \quad u \in \mathbf{B}$$

for $i \in [0, T + 2]$.

Since f^* is continuous and bounded on $[1, T + 1] \times \mathbf{R}^{n-1}$, $T^* : \mathbf{B} \rightarrow \mathbf{R}$ is continuous and compact. Therefore, it follows from Schauder's fixed point theorem that T^* has a fixed point $u \in \mathbf{B}$, i.e., (BVP^*) has a solution $u \in \mathbf{B}$.

Step (2) Let

$$H(i) = \Delta^{n-2}u(i) - \Delta^{n-2}w(i) \text{ on } [0, T + 2],$$

then for $\theta \in [1, T + 1]$ such that $H(\theta) > 0$, it follows that

$$\begin{aligned} \Delta^2 H(\theta - 1) &\geq -f^*(\theta, u(\theta), \dots, \Delta^{n-3}u(\theta), \Delta^{n-2}u(\theta)) \\ &\quad + f(\theta, w(\theta), \dots, \Delta^{n-3}w(\theta), \Delta^{n-2}w(\theta)) \\ &= -f(\theta, \eta_1, \dots, \eta_{n-2}, \Delta^{n-2}w(\theta)) - \rho(\Delta^{n-2}w(\theta) - \Delta^{n-2}u(\theta)) \\ &\quad + f(\theta, w(\theta), \dots, \Delta^{n-3}w(\theta), \Delta^{n-2}w(\theta)) \quad (\eta_l \text{ is defined in Step(1)}) \\ &\geq -\rho(\Delta^{n-2}w(\theta) - \Delta^{n-2}u(\theta)) > 0, \quad (\text{by using } (C_5)). \end{aligned}$$

Therefore, there is no $\theta \in [1, T + 1]$ such that $H(\theta) > 0$ and $\Delta^2 H(\theta - 1) \leq 0$.

Step (3) We claim that $H(i) \leq 0$ on $[0, T + 1]$. Suppose to the contrary that there exists $i_0 \in [0, T + 1]$ such that $H(i_0) > 0$. Then, there is a $\theta \in [0, T + 1]$ such that

$$H(\theta) = \max_{i \in [0, T+1]} H(i) > 0.$$

If $\theta \in [1, T]$, then we have

$$\Delta^2 H(\theta - 1) = [H(\theta + 1) - H(\theta)] - [H(\theta) - H(\theta - 1)] \leq 0,$$

which contradicts the conclusion of Step(2). Hence, $\theta \in \{0, T + 1\}$.

Case (1) Suppose that $\theta = T + 1$, i.e., $H(T + 1) > 0$. We shall first show that $\Delta H(T + 1) \leq 0$ is impossible. For this, assume that $\Delta H(\theta - 1) = \Delta H(T) < 0$, then $H(T + 1) < H(T)$. This implies that $H(\theta) = H(T + 1)$ cannot be the maximum of $H(i)$. Hence $\Delta H(T) \geq 0$. But, then we have

$$\Delta^2 H(\theta - 1) = \Delta^2 H(T) = \Delta H(T + 1) - \Delta H(T) \leq 0,$$

which contradicts the conclusion of Step(2). Thus $\Delta H(T + 1) > 0$. Now from (BC) and (4), if $\gamma = 0$ we find $\Delta H(T + 1) \leq 0$, which we have seen is impossible. Further, if $\gamma > 0$ then in view of $\Delta^{n-1}u(T + 1) > \Delta^{n-1}w(T + 1)$, we have

$$\Delta^{n-2}w(T + 1) \geq \frac{-\Delta^{n-1}w(T + 1)\delta}{\gamma} \geq \frac{-\Delta^{n-1}u(T + 1)\delta}{\gamma} = \Delta^{n-2}u(T + 1),$$

i.e., $H(T+1) \leq 0$. But, this contradicts $H(T+1) > 0$. Hence, $\theta \neq T+1$.

Case (2) Suppose that $\theta = 0$, i.e., $H(0) > 0$. We shall first show that $\Delta H(0) \geq 0$ is impossible. For this, assume that $\Delta H(\theta) = \Delta H(0) > 0$, then $H(0) < H(1)$. This implies that $H(\theta) = H(0)$ cannot be the maximum of $H(i)$, thus $\Delta H(0) = 0$. Now, we claim that $\Delta^2 H(\theta) = \Delta^2 H(0) \leq 0$. Assume to the contrary that $\Delta^2 H(0) > 0$, then $H(2) - H(1) > H(1) - H(0) = \Delta H(0) = 0$. This leads to $H(2) > H(1) = H(0) = H(\theta)$. Therefore, $H(\theta) = H(0)$ is not the maximum of $H(i)$, which gives a contradiction. But, then from $\Delta^2 H(\theta) = \Delta^2 H(0) = \Delta^2 H(1-1) \leq 0$, and the conclusion of Step(2), we obtain a contradiction. Thus, $\Delta H(0) < 0$. Now from (BC) and (4), if $\alpha = 0$ we find $\Delta H(0) \geq 0$, which we have seen is impossible. Further, if $\alpha > 0$ then in view of $\Delta^{n-1}u(0) < \Delta^{n-1}w(0)$, we have

$$\Delta^{n-2}w(0) \geq \frac{\Delta^{n-1}w(0)\beta}{\alpha} \geq \frac{\Delta^{n-1}u(0)\beta}{\alpha} = \Delta^{n-2}u(0),$$

i.e., $H(0) \leq 0$. But, this contradicts $H(0) > 0$. Hence, $\theta \neq 0$.

Step (4) From Step(3), we have

$$\Delta^{n-2}u(i) \leq \Delta^{n-2}w(i) \text{ on } [0, T+1].$$

Similarly, we can show that

$$\Delta^{n-2}v(i) \leq \Delta^{n-2}u(i) \text{ on } [0, T+1].$$

Thus, it follows that

$$\Delta^{n-2}v(i) \leq \Delta^{n-2}u(i) \leq \Delta^{n-2}w(i) \text{ on } [0, T+1].$$

Now the conditions $\Delta^m v(0) \leq 0$, $\Delta^m u(0) = 0$, and $\Delta^m w(0) \geq 0$, $0 \leq m \leq n-3$, immediately imply that

$$\Delta^m v(i) \leq \Delta^m u(i) \leq \Delta^m w(i) \text{ on } [0, T+n-m-1] \text{ for } m = 0, 1, \dots, n-2.$$

Therefore, from the definition of f^* , we have

$$f^*(i, u(i), \Delta u(i), \dots, \Delta^{n-2}u(i)) = f(i, u(i), \Delta u(i), \dots, \Delta^{n-2}u(i)) \text{ on } [1, T+1].$$

This implies that $u(i)$ is infact a solution of (BVP).

In our next result for a given $g(i, u_1, u_2, \dots, u_{n-1}) \in C([1, T+1] \times [0, \infty)^{n-1}; [0, \infty))$ we shall need the following:

$$\begin{aligned} \max g_0 &= \lim_{u_1, u_2, \dots, u_{n-1} \rightarrow 0^+} \max_{i \in [1, T+1]} \frac{g(i, u_1, u_2, \dots, u_{n-1})}{u_{n-1}}, \\ \min g_0 &= \lim_{u_1, u_2, \dots, u_{n-1} \rightarrow 0^+} \min_{i \in [n-1, T+1]} \frac{g(i, u_1, u_2, \dots, u_{n-1})}{u_{n-1}}, \\ \max g_\infty &= \lim_{u_1, u_2, \dots, u_{n-1} \rightarrow \infty} \max_{i \in [1, T+1]} \frac{g(i, u_1, u_2, \dots, u_{n-1})}{u_{n-1}}, \end{aligned}$$

$$\min g_{\infty} = \lim_{u_1, u_2, \dots, u_{n-1} \rightarrow \infty} \min_{i \in [n-1, T+1]} \frac{g(i, u_1, u_2, \dots, u_{n-1})}{u_{n-1}}.$$

We shall also need the numbers

$$D_1 = \left(\sum_{j=1}^{T+1} k(j, j) \right)^{-1}, \quad D_2 = \left(\sum_{j=n-1}^{T+1} k(n-1, j) \right)^{-1}, \quad \text{and} \quad M^* = \frac{M}{L}.$$

THEOREM 2. *Suppose that*

(H₁) *$f \in C([1, T+1] \times \mathbb{R}^{n-1}; \mathbb{R})$ is such that*

$$f(i, 0, 0, \dots, 0) \geq 0 \quad \text{on } [1, T+1] \quad (f \text{ may be negative for } u_i \neq 0).$$

(H₂) *There exists a function $g(i, u_1, u_2, \dots, u_{n-1}) \in C([1, T+1] \times [0, \infty)^{n-1}; [0, \infty))$ which satisfies*

$$g(i, |u_1|, |u_2|, \dots, |u_{n-1}|) \geq f(i, u_1, u_2, \dots, u_{n-1}) \quad \text{on } [1, T+1] \times [0, \infty)^{n-1}.$$

(H₃) *One of the following holds:*

(P₁) *$\max g_0 = A_1 \in [0, D_1/L]$ and $\min g_{\infty} = A_2 \in (D_2/M^*, \infty]$.*

(P₂) *$\min g_0 = A_3 \in (D_2/M^*, \infty]$ and $\max g_{\infty} = A_4 \in [0, D_1/L]$.*

(P₃) *There exist two functions $q : [1, T+1] \rightarrow [0, \infty)$, and $h(u_1, u_2, \dots, u_{n-1}) \in C([0, \infty)^{n-1}; [0, \infty))$, such that h is increasing with respect to u_{n-1} , and*

$$\begin{cases} g(i, u_1, u_2, \dots, u_{n-1}) = q(i)h(u_1, u_2, \dots, u_{n-1}), \\ \sup_{u_{n-1} \in (0, \infty)} \min_{(u_1, u_2, \dots, u_{n-2}) \in [0, \infty)} \frac{u_{n-1}}{Qh(u_1, u_2, \dots, u_{n-1})} > 1, \end{cases}$$

where $Q = \max_{i \in [1, T+1]} \sum_{j=1}^{T+1} k(i, j)q(j)$.

Then (BVP) has at least one non-negative solution.

Proof. Using the same techniques as in Agarwal and Wong [8–10], we can show that

$$(BVP^{**}) \begin{cases} (E^{**}) \quad \Delta^n w(i-1) + g(i, w(i), \Delta w(i), \dots, \Delta^{n-2} w(i)) = 0 \quad \text{for } i \in [1, T+1], \\ (BC^{**}) \quad \begin{cases} \Delta^m w(0) = 0, & 0 \leq m \leq n-3, \\ \alpha \Delta^{n-2} w(0) - \beta \Delta^{n-1} w(0) = 0, \\ \gamma \Delta^{n-2} w(T+1) + \delta \Delta^{n-1} w(T+1) = 0, \end{cases} \end{cases}$$

has at least one solution $w(i)$, $i \in [0, T+n]$ such that $\Delta^m w(i) \geq 0$, $i \in [0, T+n-m-1]$ for $m = 0, 1, \dots, n-2$. It is clear that this $w(i)$ and $v(i) = 0$ are upper and lower solutions of (BVP), respectively. The result now follows from Theorem 1.

REFERENCES

- [1] R. P. AGARWAL, *On boundary value problems for second order discrete systems*, *Applicable Analysis* **20** (1985), 1–17.
- [2] R. P. AGARWAL, *Difference Equations and Inequalities*, Marcel Dekker, New York, 1992.
- [3] R. P. AGARWAL AND D. O'REGAN, *Boundary value problems for discrete equations*, *Applied Mathematics Letters* **10**(4) (1997), 83–89.
- [4] R. P. AGARWAL AND D. O'REGAN, *A fixed point approach for nonlinear discrete boundary value problems*, *Computers Math. Applic.*, to appear.
- [5] R. P. AGARWAL AND D. O'REGAN, *Singular discrete boundary value problems*, *Applied Mathematics Letters*, to appear.
- [6] R. P. AGARWAL AND D. O'REGAN, *Nonpositone discrete boundary value problems*, *Nonlinear Analysis*, to appear.
- [7] R. P. AGARWAL AND D. O'REGAN, *Difference equations in abstract spaces*, *J. Austr. Math. Soc., Ser.(A)*, to appear.
- [8] R. P. AGARWAL AND F. H. WONG, *Existence of positive solutions for higher order difference equations*, *Applied Mathematics Letters* **10**(5) (1997), 67–74.
- [9] R. P. AGARWAL AND F. H. WONG, *Existence of positive solutions for non-positive difference equations*, *Mathl. Computer Modelling*, to appear.
- [10] R. P. AGARWAL AND F. H. WONG, *An application of topological transversality to non-positive higher order difference equations*, *Appl. Math. Comp.*, to appear.
- [11] R. P. AGARWAL AND P. J. Y. WONG, *Advanced Topics in Difference Equations*, Kluwer, Dordrecht, 1997.
- [12] R. GAINES, *Difference equations associated with boundary value problems for second order nonlinear ordinary differential equations*, *SIAM J. Numer. Anal.* **11** (1974), 411–434.
- [13] J. HENDERSON, *Singular boundary value problems for difference equations*, *Dynamic Systems and Applications* **1** (1992), 271–282.
- [14] J. HENDERSON, *Singular boundary value problems for higher order difference equations*, in *Proceedings of the First World Congress on Nonlinear Analysts 1992*, ed. V. Lakshmikantham, Walter de Gruyter and Co., 1996, pp. 1139–1150.
- [15] A. LASOTA, *A discrete boundary value problem*, *Ann. Polon. Math.* **20** (1968), 183–190.
- [16] A. C. PETERSON, *Existence and uniqueness theorems for nonlinear difference equations*, *J. Math. Anal. Appl.* **125** (1987), 185–191.
- [17] P. J. Y. WONG AND R. P. AGARWAL, *On the existence of solutions of singular boundary value problems for higher order difference equations*, *Nonlinear Analysis* **28** (1997), 277–287.
- [18] P. J. Y. WONG AND R. P. AGARWAL, *Topological Methods in Nonlinear Analysis*, to appear.
- [19] P. J. Y. WONG AND R. P. AGARWAL, *On the eigenvalues of boundary value problems for higher order difference equations*, *Rocky Mountain J. Math.*, to appear.
- [20] W. ZHUANG, Y. CHENG AND S. S. CHENG, *Monotone methods for a discrete boundary value problem*, *Computers Math. Applic.* **32** (1996), 41–49.

(Received February 16, 1998)

Ravi P. Agarwal
Department of Mathematics
National University of Singapore
10 Kent Ridge Crescent
Singapore 119260
e-mail: matravip@leonis.nus.sg

Fu-Hsiang Wong
Department of Mathematics and Science
National Taipei Teacher's College
134, Ho-Ping E. Rd. Sec. 2
Taipei 10659
Taiwan
Republic of China
e-mail: wongjyp@nievax.nie.ac.sg