

A NOTE ON SOME CLASSES OF FOURIER COEFFICIENTS

ŽIVORAD TOMOVSKI

(communicated by H. M. Srivastava)

Abstract. It is shown that the class S_p ($p > 1$) is a subclass of $C \cap BV$, where C is the Garret–Stanojević class and BV the class of sequences of bounded variation. A new direct proof of this theorem is given.

1. Introduction

A sequence $\{a_k\}$ belongs to the class S , if $a_k \rightarrow 0$ as $k \rightarrow \infty$ and there exists a monotonically decreasing sequence $\{A_k\}$ such that $\sum_{k=1}^{\infty} A_k < \infty$ and $|\Delta a_k| \leq A_k$ for all k .

Now, we say that a sequence $\{a_k\}$ belongs to the class S_p or $a_k \in S_p$ if $a_k \rightarrow 0$ as $k \rightarrow \infty$ and there exists a monotonically decreasing sequence $\{A_k\}$ such that $\sum_{k=1}^{\infty} A_k < \infty$ and $\frac{1}{n} \sum_{k=1}^n \frac{|\Delta a_k|^p}{A_k^p} = O(1)$.

Thus in view of the above definitions, it is obvious that $S \subset S_p$.

The class C was defined in [3] as follows: $\{a_n\} \in C$ if for every $\varepsilon > 0$ there is a $\delta > 0$ such that

$$\int_0^\delta \left| \sum_{k=n}^{\infty} \Delta a_k D_k(x) \right| dx < \varepsilon,$$

for all n , where

$$D_n(x) = \frac{1}{2} + \sum_{k=1}^n \cos kx$$

is the Dirichlet kernel.

2. Main result

THEOREM. $S_p \subseteq C \cap BV$.

Mathematics subject classification (1991): 26D15, 42A20.

Key words and phrases: Dirichlet kernel, Abel's transformation, Hölder inequality, Hausdorff–Young inequality, Cauchy condensation test.

Proof. It suffices to show that

$$C_n = \int_0^\pi \left| \sum_{k=n}^{\infty} \Delta a_k D_k(x) \right| dx = o(1), \quad n \rightarrow \infty.$$

For each n , let k_n be the least natural number such that $n \leq 2^{k_n} - 1$.

Then C_n can be majorized by

$$C_n \leq \int_0^\pi \left| \sum_{j=n}^{2^{k_n}-1} \Delta a_j D_j(x) \right| dx + \sum_{l=k_n}^{\infty} \int_0^\pi \left| \sum_{j=2^l}^{2^{l+1}-1} \Delta a_j D_j(x) \right| dx = I_1 + I_2.$$

The second term is written as follows:

$$I_2 = \sum_{l=k_n}^{\infty} \left\{ \int_0^{1/2^{l+1}} + \int_{1/2^{l+1}}^{\pi} \right\} \left| \sum_{j=2^l}^{2^{l+1}-1} \Delta a_j D_j(x) \right| dx = \Sigma_1 + \Sigma_2.$$

For the first term, the uniform estimate $|D_n(x)| \leq n + \frac{1}{2}$, is applied, i.e.

$$\begin{aligned} \Sigma_1 &\leq \sum_{l=k_n}^{\infty} \frac{1}{2^{l+1}} \sum_{j=2^l}^{2^{l+1}-1} |\Delta a_j| \left(j + \frac{1}{2} \right) \\ &= \sum_{i=2^{k_n}}^{\infty} \frac{1}{2i} \sum_{j=i}^{2i-1} |\Delta a_j| \left(j + \frac{1}{2} \right) \leq \sum_{i=2^{k_n}}^{\infty} \frac{1}{2i} \sum_{j=i}^{2i-1} |\Delta a_j| 2i \\ &= \sum_{i=2^{k_n}}^{\infty} \sum_{j=i}^{2i-1} |\Delta a_j|. \end{aligned}$$

By summation by parts, we have:

$$\begin{aligned} \sum_{i=2^{k_n}}^{\infty} |\Delta a_i| &= \sum_{i=2^{k_n}}^{\infty} \frac{|\Delta a_i|}{A_i} A_i = \sum_{i=2^{k_n}-1}^{\infty} \Delta A_i \sum_{j=1}^i \frac{|\Delta a_j|}{A_j} + A_{2^{k_n}} \sum_{j=1}^{2^{k_n}-1} \frac{|\Delta a_j|}{A_j} \\ &\leq \sum_{i=2^{k_n}-1}^{\infty} i(\Delta A_i) \left(\frac{1}{i} \sum_{j=1}^i \frac{|\Delta a_j|^p}{A_j^p} \right)^{1/p} + 2^{k_n} A_{2^{k_n}} \left(\frac{1}{2^{k_n}} \sum_{j=1}^{2^{k_n}} \frac{|\Delta a_j|^p}{A_j^p} \right)^{1/p} \\ &= O(1) \left[\sum_{i=2^{k_n}-1}^{\infty} i(\Delta A_i) + 2^{k_n} A_{2^{k_n}} \right]. \end{aligned}$$

Since $\sum_{n=1}^{\infty} A_n < \infty$, both terms on the right-hand side of the above inequality are $o(1)$, as $n \rightarrow \infty$.

Thus $\Sigma_1 = o(1)$, as $n \rightarrow \infty$

Let

$$\Sigma_2 = \sum_{l=k_n}^{\infty} \int_{1/2^{l+1}}^{\pi} \left| \sum_{j=2^l}^{2^{l+1}-1} \frac{|\Delta a_j|}{A_j} A_j D_j(x) \right| dx.$$

Applying Abel's transformation, we get:

$$\begin{aligned} \int_{1/2^{l+1}}^{\pi} \left| \sum_{j=2^l}^{2^{l+1}-1} \frac{\Delta a_j}{A_j} A_j D_j(x) \right| dx &\leq \sum_{j=2^l}^{2^{l+1}-2} \Delta A_j \int_{1/2^{l+1}}^{\pi} \left| \sum_{r=1}^j \frac{\Delta a_r}{A_r} D_r(x) \right| dx \\ &\quad + A_{2^l} \int_{1/2^{l+1}}^{\pi} \left| \sum_{r=1}^{2^l} \frac{\Delta a_r}{A_r} D_r(x) \right| dx, \quad \text{i.e.} \end{aligned}$$

$$\Sigma_2 \leq \sum_{l=k_n}^{\infty} \sum_{j=2^l}^{2^{l+1}-2} \Delta A_j \int_{1/2^{l+1}}^{\pi} \left| \sum_{r=1}^j \frac{\Delta a_r}{A_r} D_r(x) \right| dx + \sum_{l=k_n}^{\infty} A_{2^l} \int_{1/2^{l+1}}^{\pi} \left| \sum_{r=1}^{2^l} \frac{\Delta a_r}{A_r} D_r(x) \right| dx.$$

Applying the Hölder inequality, we get:

$$\begin{aligned} I_l &= \int_{1/2^{l+1}}^{\pi} \left| \sum_{r=1}^{2^l} \frac{\Delta a_r}{A_r} D_r(x) \right| dx = \int_{1/2^{l+1}}^{\pi} \frac{1}{\sin \frac{x}{2}} \left| \sum_{r=1}^{2^l} \frac{\Delta a_r}{A_r} \sin \left(r + \frac{1}{2} \right) x \right| dx \\ &\leq \left[\int_{1/2^{l+1}}^{\pi} \frac{dx}{\left(\sin \frac{x}{2} \right)^p} \right]^{1/p} \left[\int_{1/2^{l+1}}^{\pi} \left| \sum_{r=1}^{2^l} \frac{\Delta a_r}{A_r} \sin \left(r + \frac{1}{2} \right) x \right|^q dx \right]^{1/q} \end{aligned}$$

where $1/p + 1/q = 1$.

Since

$$\int_{1/2^{l+1}}^{\pi} \frac{dx}{\left(\sin \frac{x}{2} \right)^p} \leq \pi^p \int_{1/2^{l+1}}^{\pi} \frac{dx}{x^p} \leq M_p (2^{l+1})^{p-1},$$

(M_p is absolute constant depending on p), it follows that

$$I_l \leq (2^{l+1})^{1/q} (M_p)^{1/p} \left[\int_{1/2^{l+1}}^{\pi} \left| \sum_{r=1}^{2^l} \frac{\Delta a_r}{A_r} \sin \left(r + \frac{1}{2} \right) x \right|^q dx \right]^{1/q}.$$

Applying the Hausdorff–Young inequality to the last integral we get:

$$\left(\int_{1/2^{l+1}}^{\pi} \left| \sum_{r=1}^{2^l} \frac{\Delta a_r}{A_r} \sin \left(r + \frac{1}{2} \right) x \right|^q dx \right)^{1/q} \leq B_p \left(\sum_{r=1}^{2^l} \frac{|\Delta a_r|^p}{A_r^p} \right)^{1/p}.$$

Thus

$$I_l \leq 2^{l+1} C_p \left(\frac{1}{2^{l+1}} \sum_{r=1}^{2^{l+1}} \frac{|\Delta a_r|^p}{A_r^p} \right)^{1/p} \quad (C_p > 0).$$

Then

$$\Sigma_2 = O(1) \left[\sum_{l=k_n}^{\infty} \sum_{j=2^l}^{2^{l+1}-2} j \Delta A_j + 2 \sum_{l=k_n}^{\infty} 2^l A_{2^l} \right].$$

Applying the Cauchy condensation test,

$$\sum_{l=k_n}^{\infty} 2^l A_{2^l} = o(1), \quad n \rightarrow \infty.$$

But

$$\sum_{j=2^l}^{2^{l+1}-2} j \Delta A_j = \sum_{j=2^l}^{2^{l+1}-1} A_j - (2^{l+1} - 1) A_{2^{l+1}-1} \leq 2^l A_{2^l} - 2^{l+1} A_{2^{l+1}}.$$

Thus

$$\sum_{l=k_n}^{\infty} \sum_{j=2^l}^{2^{l+1}-2} j \Delta A_j \leq \sum_{l=k_n}^{\infty} [2^l A_{2^l} - 2^{l+1} A_{2^{l+1}}] = 2^{k_n} A_{2^{k_n}} = o(1), \quad n \rightarrow \infty,$$

i.e. $\Sigma_2 = o(1)$, $n \rightarrow \infty$. Finally $I_2 = o(1)$, $n \rightarrow \infty$.

The same method applied to I_1 yields the estimate

$$I_1 \leq O(1) \sum_{l=n}^{2^{k_n}-1} |\Delta a_l| + O(1) \left(\sum_{j=n}^{2^{k_n}-1} j \Delta A_j + 2n A_n \right).$$

Letting $n \rightarrow \infty$, the proof of the theorem follows.

REFERENCES

- [1] G. A. FOMINE, *A class of trigonometric series*, Mat. Zametki **23** (1978), 213–222.
- [2] J. W. GARRET AND Č. V. STANOJEVIĆ, *On L^1 convergence of certain cosine sums*, Proc. Amer. Math. Soc. **54** (1976), 102–105.
- [3] Č. V. STANOJEVIĆ, *Classes of L^1 convergence of Fourier and Fourier Stiltjes series*, Proc. Amer. Math. Soc. **82** (1981), 209–215.

(Received April 15, 1998)

*Faculty of Mathematical
and Natural Sciences
P.O. Box 162
91000 Skopje
Macedonia
e-mail: tomovski@iunona.pmf.ukim.edu.mk*