

SEVERAL PROPERTIES ON CLASS A INCLUDING p -HYPONORMAL AND LOG-HYPONORMAL OPERATORS

MASATOSHI ITO

Abstract. In this paper, we shall show that Yamazaki's result in [23] remains valid for class A introduced in [13] as a class of operators including p -hyponormal and log-hyponormal operators. Moreover we shall show several properties on class A which correspond to the properties on paranormal in [8][12][18] and [19].

Mathematics subject classification (1991): 47B20, 47A63.

Key words and phrases: class A operator, paranormal operator, order preserving operator inequality.

REFERENCES

- [1] A. ALUTHGE, *On p -hyponormal operators for $0 < p < 1$* , Integral Equations Operator Theory, **13** (1990), 307–315.
- [2] A. ALUTHGE AND D. WANG, *Powers of p -hyponormal operators*, preprint.
- [3] T. ANDO, *Operators with a norm condition*, Acta Sci. Math. (Szeged), **33** (1972), 169–178.
- [4] M. CHO AND M. ITOH, *Putnam's inequality for p -hyponormal operators*, Proc. Amer. Math. Soc., **123** (1995), 2435–2440.
- [5] M. FUJII, *Furuta's inequality and its mean theoretic approach*, J. Operator Theory, **23** (1990), 67–72.
- [6] M. FUJII, S. IZUMINO AND R. NAKAMOTO, *Classes of operators determined by the Heinz-Kato-Furuta inequality and the Hölder-McCarthy inequality*, Nihonkai Math. J., **5** (1994), 61–67.
- [7] M. FUJII, R. NAKAMOTO AND H. WATANABE, *The Heinz-Kato-Furuta inequality and hyponormal operators*, Math. Japon., **40** (1994), 469–472.
- [8] T. FURUTA, *On the class of paranormal operators*, Proc. Japan Acad., **43** (1967), 594–598.
- [9] T. FURUTA, *$A \geq B \geq 0$ assures $(B^r A P B^r)^{1/q} \geq B^{(p+2r)/q}$ for $r \geq 0$, $p \geq 0$, $q \geq 1$ with $(1+2r)q \geq p+2r$* , Proc. Amer. Math. Soc., **101** (1987), 85–88.
- [10] T. FURUTA, *An elementary proof of an order preserving inequality*, Proc. Japan Acad. Ser. A Math. Sci., **65** (1989), 126.
- [11] T. FURUTA, *Extension of the Furuta inequality and Ando-Hiai log-majorization*, Linear Algebra Appl., **219** (1995), 139–155.
- [12] T. FURUTA, M. HORIE AND R. NAKAMOTO, *A remark on a class of operators*, Proc. Japan Acad., **43** (1967), 607–609.
- [13] T. FURUTA, M. ITO AND T. YAMAZAKI, *A subclass of paranormal operators including class of log-hyponormal and several related classes*, Scientiae Mathematicae, **1** (1998), 389–403.
- [14] T. FURUTA, T. YAMAZAKI AND M. YANAGIDA, *Operator functions implying generalized Furuta inequality*, Math. Inequal. Appl., **1** (1998), 123–130.
- [15] T. FURUTA AND M. YANAGIDA, *Further extensions of Aluthge transformation on p -hyponormal operators*, Integral Equations Operator Theory, **29** (1997), 122–125.
- [16] T. FURUTA AND M. YANAGIDA, *On powers of p -hyponormal and log-hyponormal operators*, preprint.
- [17] T. HURUYA, *A note on p -hyponormal operators*, Proc. Amer. Math. Soc., **125** (1997), 3617–3624.
- [18] I. ISTRĂTESCU AND V. ISTRĂTESCU, *On some classes of operators. I*, Proc. Japan Acad., **43** (1967), 605–606.
- [19] V. ISTRĂTESCU, T. SAITO AND T. YOSHINO, *On a class of operators*, Tôhoku Math. J., **18** (1966), 410–413.

- [20] E. KAMEI, *A satellite to Furuta's inequality*, Math. Japon., **33** (1988), 883–886.
- [21] C. A. McCARTHY, c_p , Israel J. Math., **5** (1967), 249–271.
- [22] K. TANAHASHI, *On log-hyponormal operators*, preprint.
- [23] T. YAMAZAKI, *Extensions of the results on p -hyponormal and log-hyponormal operators by Aluthge and Wang*, to appear in SUT J. Math.
- [24] T. YOSHINO, *The p -hyponormality of the Aluthge transform*, Interdiscip. Inform. Sci., **3** (1997), 91–93.