

## NONCONVEX FUNCTIONS AND SEPARATION BY POWER MEANS

ZSOLT PÁLES

*Abstract.* In this note we show that, for a nonconvex function defined on a real interval, there exists a point where this function behaves like a strictly concave function. Due to this result, global convexity can be characterized as pointwise convexity everywhere. As an application, a necessary and sufficient condition for the separability of quasarithmetic means with power means is obtained.

*Mathematics subject classification (1991):* Primary 26A51, 26B25.

*Key words and phrases:* Convexity, quasarithmetic means, power means, separation.

### REFERENCES

- [1] G. H. HARDY, J. E. LITTLEWOOD, AND G. PÓLYA, *Inequalities*, Cambridge Univ. Press, Cambridge – New York – London, 1934 (first edition), 1952 (second edition).
- [2] M. KUCZMA, *An Introduction to the Theory of Functional Equations and Inequalities*, Państwowe Wydawnictwo Naukowe, Warszawa – Kraków – Katowice, 1985.
- [3] ZS. PÁLES, *Hölder-type inequalities for quasarithmetic means*, Acta Math. Hung., **47** (1986), 395-399.
- [4] A. W. ROBERTS AND D. E. VARBERG, *Convex Functions*, Academic Press, New York – London, 1973.