

THE EULER FORMULÆ AND CONVEX FUNCTIONS

LJ. DEDIĆ, C. E. M. PEARCE AND J. PEČARIĆ

Abstract. The Euler midpoint and Euler–Simpson formulæ are used with functions possessing various convexity and concavity properties to derive inequalities pertinent to numerical integration.

Mathematics subject classification (1991): 26D15, 26D20, 26D99.

Key words and phrases: Hadamard inequality, r -convexity, integral inequalities, Euler midpoint formula, Euler–Simpson formula.

REFERENCES

- [1] I. S. Berezin and N. P. Zhidkov, *Computing Methods*, Vol. I, Pergamon Press, Oxford-London-Edinburgh-New York-Paris-Frankfurt, 1965.
- [2] Lj. Dedić, C. E. M. Pearce and J. Pečarić, Hadamard and Dragomir–Agarwal inequalities, higher-order convexity and the Euler formula. Submitted.
- [3] S. S. Dragomir and R. P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, *Appl. Mat. Lett.* **11** (5), 91–95 (1998).
- [4] V. I. Krylov, *Approximate calculation of integrals*, The MacMillan Comp., New York, 1962.
- [5] C. E. M. Pearce and J. Pečarić, Inequalities for differentiable mappings with applications to special means and quadrature formulæ, to appear in *Appl. Math. Lett.*