

SMOOTHNESS CONDITIONS AND FOURIER SERIES

SERGEY TIKHONOV

Abstract. We study the relationships between smoothness properties of a 2π -periodic function and the asymptotic behavior of its Fourier coefficients. The smoothness conditions of a function are expressed in terms of it being in the generalized Lipschitz classes.

Mathematics subject classification (2000): 26A16, 42A16.

Key words and phrases: Lipschitz class, trigonometric series, moduli of smoothness of fractional order.

REFERENCES

- [1] N. K. BARI, *On best approximation of two conjugate functions by trigonometric polynomials*, Izv. Akad. Nauk SSSR, Ser. Mat., **19**, (5) (1955), 285–302.
- [2] N. K. BARI, S. B. STEČKIN, *Best approximations and differential properties of two conjugate functions*, Trudy Moskov. Mat. Obšč., **5**, (1956), 483–522.
- [3] N. H. BINGHAM, C. M. GOLDIE AND J. L. TEUGELS, *Regular variation*, Cambridge Univ. Press, 1987.
- [4] R. P. BOAS JR., *Fourier series with positive coefficients*, J. Math. Anal. Appl., **17**, (1967), 463–483.
- [5] R. P. BOAS JR., *Integrability theorems for trigonometric transforms*, Springer-Verlag, New York, 1967.
- [6] P. L. BUTZER, H. DYCKHOFF, E. GÖRLICH AND R. L. STENS, *Best trigonometric approximation, fractional order derivatives and Lipschitz classes*, Can. J. Math., **29** (1977), 781–793.
- [7] P. L. BUTZER, U. WESTPHAL, *An access to fractional differentiation via fractional difference quotients*, Fractional calculus and its applications. Lecture Notes in Math., **457**, Springer, 1975.
- [8] LING-YAU CHAN, *On Fourier series with nonnegative coefficients and two problems of R. P. Boas*, J. Math. Anal. Appl., **110**, (1) (1985), 116–129.
- [9] LING-YAU CHAN, *Generalized Lipschitz classes and asymptotic behavior of Fourier series*, J. Math. Anal. Appl., **155**, (2) (1991), 371–377.
- [10] M. IZUMI, S. IZUMI, *Lipschitz classes and Fourier coefficients*, J. Math. Mech., **18**, (1969), 857–870.
- [11] L. LEINDLER, *Power-monotone sequences and Fourier series with positive coefficients*, J. Inequal. Pure Appl. Math., **1** (1), Article No.1, (2000), Available online at URL: <http://jipam.vu.edu.au/>.
- [12] L. LEINDLER, J. NEMETH, *On the connection between quasi power-monotone and quasi geometrical sequences with application to integrability theorems for power series*, Acta Math. Hung., **68**, (1–2) (1995) 7–19.
- [13] J. NEMETH, *Fourier series with positive coefficients and generalized Lipschitz classes*, Acta Sci. Math., (Szeged), **54**, (3–4) (1990), 291–304.
- [14] J. NEMETH, *Power-monotone sequences and Fourier series with positive coefficients*, J. Inequal. Pure Appl. Math., **2** (2), Article No. 14 2001. Available online at URL: <http://jipam.vu.edu.au/>.
- [15] S. TIKHONOV, *On generalized Lipschitz classes and Fourier series*, Z. Anal. Anwendungen, **23**, (4) (2004), 745–764.
- [16] S. TIKHONOV, *On moduli of smoothness offractional order* Real Analysis Exchange, **30**, (2) (2004/2005), 507–518.
- [17] S. TIKHONOV, *On two theorem of Lorentz*, Izv. Ross. Akad. Nauk, Ser. Mat., **69**, (1) (2005), 165–178, translation in Russian Acad. Sci. Izv. Math, **69**, (1) (2005), 163–175.
- [18] V. TOTIK, *On the strong approximation of Fourier series*, Acta Math. Acad. Sci. Hung., **35**, (1980), 151–172.
- [19] A. ZYGMUND, *Trigonometric series*, Cambridge University Press, 1959.