

SOME NEW INEQUALITIES SIMILAR TO HARDY–HILBERT’S INEQUALITY

S. K. SUNANDA, C. NAHAK AND S. NANDA

(Communicated by R. Mohapatra)

Abstract. In this paper we have studied some new inequalities similar to Hardy-Hilbert’s inequality. As applications, we have considered the associated integral inequalities.

1. Introduction

If $a_n, b_n \geq 0$, $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, and $0 < \sum_{n=1}^{\infty} a_n^p < \infty$, $0 < \sum_{n=1}^{\infty} b_n^q < \infty$, then the famous Hardy-Hilbert inequality (see Hardy et al. [6]) is given by

$$\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{a_m b_n}{m+n} < \frac{\pi}{\sin(\pi/p)} \left\{ \sum_{n=1}^{\infty} a_n^p \right\}^{1/p} \left\{ \sum_{n=1}^{\infty} b_n^q \right\}^{1/q}, \quad (1.1)$$

where the constant factor $\frac{\pi}{\sin(\pi/p)}$ is the best possible. Inequality (1.1) led to many papers dealing with alternative proofs, generalizations, numerous variants, and applications in analysis (Hardy et al. [6]). Recently, Yang, Gao and Debnath [1, 2, 8] gave some strengthened version of (1.1). By introducing a parameter Yang [3] gave an extension of (1.1) as

$$\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{a_m b_n}{m^{\lambda} + n^{\lambda}} < \frac{\pi}{\lambda \sin(\pi/p)} \left\{ \sum_{n=1}^{\infty} n^{(p-1)(1-\lambda)} a_n^p \right\}^{1/p} \left\{ \sum_{n=1}^{\infty} n^{(q-1)(1-\lambda)} b_n^q \right\}^{1/q},$$

where the constant factor $\frac{\pi}{\lambda \sin(\pi/p)}$, ($0 < \lambda \leq \min p, q$) is the best possible.

In 2003 yang [4] gave an extension of (1.1) as

$$\sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \frac{a_m b_n}{\ln mn} < \frac{\pi}{\sin(\pi/p)} \left\{ \sum_{n=2}^{\infty} n^{p-1} a_n^p \right\}^{1/p} \left\{ \sum_{n=2}^{\infty} n^{q-1} b_n^q \right\}^{1/q}, \quad (1.2)$$

where the constant factor $\frac{\pi}{\sin(\pi/p)}$ is the best possible.

Mathematics subject classification (2010): 26D15.

Keywords and phrases: Holder’s inequality, Hardy-Hilbert’s inequality, generalized l_p space, generalized L_p space.

In this paper, we have generalized the results of [4], which is related to the double series of the form

$$\sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \frac{a_m b_n}{\ln mn}. \quad (1.3)$$

For this series $\sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \frac{a_m b_n}{\ln(mn)}$, we have estimated the weight co-efficient of the form

$$w_{r_k}(n) = \sum_{m=2}^{\infty} \frac{1}{m \ln(mn)} \left(\frac{\ln n}{\ln m} \right)^{1/r_k} \quad (r_k = (p_k, q_k) > 1, n \in \mathbf{N} \setminus \{1\}). \quad (1.4)$$

2. Some preliminary results

The sequence space l_p has been generalized to $l(p)$, in the following manner (see Maddox [7] and Simmons [9]).

DEFINITION 2.1. Let $p = (p_k)$ be a bounded sequence of strictly positive numbers, with $0 < p_k \leq \sup p_k = H < \infty$. Then

$$l(p) = \{x = (x_k) : \sum |x_k|^{p_k} < \infty\}. \quad (2.1)$$

A natural metric on $l(p)$ is

$$d(x, y) = \left(\sum_{k=0}^{\infty} |x_k - y_k|^{p_k} \right)^{1/M},$$

where $M = \max(1, H)$.

DEFINITION 2.2. Let $1 \leq p < \infty$, then the function space L_p is given by

$$L_p = \{f : \int_0^{\infty} |f(x)|^p dx < \infty\}. \quad (2.2)$$

The function space L_p has been generalized to $L(p)$ in the following manner.

DEFINITION 2.3. [10] Let p be a bounded measurable function, with $0 < p(x) \leq \sup p(x) = H < \infty$. Then

$$L(p) = \{f : \int_0^{\infty} |f(x)|^{p(x)} dx < \infty\}. \quad (2.3)$$

Note that $L(p)$ is a linear topological space paranormed by $d(f)$

$$d(f) = \left(\int_0^{\infty} |f(x)|^{p(x)} dx \right)^{1/M},$$

where $M = \max(1, H)$.

Das and Nanda [5] have generalized Holder's inequality in $l(p)$ space, which is given below.

LEMMA 2.1. (Das and Nanda, [5]) *Let $(p_n)_{n=1}^{\infty}$ be a real sequence where $p_n > 1$ for all n . $(q_n)_{n=1}^{\infty}$ is defined by $\frac{1}{p_n} + \frac{1}{q_n} = 1$, for all n . Let $a_n, b_n \geq 0$. We write*

$$\begin{aligned} A_m &= \sum_{n=1}^m a_n^{p_n}, & B_m &= \sum_{n=1}^m b_n^{q_n}, \\ A &= \sum_{n=1}^{\infty} a_n^{p_n}, & B &= \sum_{n=1}^{\infty} b_n^{q_n} \end{aligned}$$

and whenever the series on the right converge. Then

(i)

$$\sum_{k=1}^m a_k b_k \leq \alpha_m \beta_m, \quad (2.4)$$

where $\alpha_m = \sup_{1 \leq n \leq m} \frac{1}{p_n} + \sup_{1 \leq n \leq m} \frac{1}{q_n}$, $\beta_m = \sup_{1 \leq n \leq m} A_m^{1/p_n} B_m^{1/q_n}$.

(ii) If $a \in l(p)$, $b \in l(q)$, then $ab \in l$ and

$$\sum_{k=1}^{\infty} a_k b_k \leq \alpha \beta, \quad (2.5)$$

where $\alpha = \sup_{n \geq 1} \frac{1}{p_n} + \sup_{n \geq 1} \frac{1}{q_n}$, $\beta = \sup_{n \geq 1} (A^{1/p_n} B^{1/q_n})$.

If p_k and q_k are constants then (2.4) and (2.5) reduce to standard Holder's inequality in l_p space.

Sunanda et. al. [10] generalized the Holder's inequality in $L(p)$ space which is given below.

LEMMA 2.2. (a) Let p and q be bounded measurable functions such that $p(x)^{-1} + q(x)^{-1} = 1$ for all x . Let $f \in L(p)$, $g \in L(q)$. Let

$$A = \int_0^{\infty} |f(x)|^{p(x)} dx, \quad B = \int_0^{\infty} |g(x)|^{q(x)} dx$$

Then for $p(x) > 1$, $f g \in L_1$ and

$$\int_0^{\infty} |f(x)g(x)| dx \leq \alpha \beta, \quad (2.6)$$

where $\alpha = \sup_x 1/p(x) + \sup_x 1/q(x)$, $\beta = \sup_x A^{1/p(x)} B^{1/q(x)}$.

(b) Let $0 < p(x) < 1$ and $p(x)^{-1} + q(x)^{-1} = 1$. If $f \in L(p)$ and $g \in L(q)$, then

$$\int_0^{\infty} |f(x)|^{p(x)} dx \leq \alpha [\sup_x p(x) + \sup_x (1 - p(x))], \quad (2.7)$$

where $\alpha = \sup_x \left[\left(\int |g(x)|^{q(x)} dx \right)^{1-p(x)} \left(\int |f(x)g(x)| dx \right)^{p(x)} \right]$.

Proof. To prove (a) For $a, b > 0$, we have

$$ab \leq \frac{a^{p(x)}}{p(x)} + \frac{b^{q(x)}}{q(x)} \quad \text{for all } x. \quad (2.8)$$

Using the above inequality we have

$$\frac{|f(x)|}{A^{1/p(x)}} \frac{|g(x)|}{B^{1/q(x)}} \leq \frac{1}{p(x)} \frac{|f(x)|^{p(x)}}{A} + \frac{1}{q(x)} \frac{|g(x)|^{q(x)}}{B}.$$

Therefore,

$$\begin{aligned} \int_0^\infty \frac{|f(x)| |g(x)|}{A^{1/p(x)} B^{1/q(x)}} dx &\leq \frac{1}{A} \int_0^\infty \frac{|f(x)|^{p(x)}}{p(x)} dx + \frac{1}{B} \int_0^\infty \frac{|g(x)|^{q(x)}}{q(x)} dx \\ &\leq \sup_x \frac{1}{p(x)} + \sup_x \frac{1}{q(x)}. \end{aligned} \quad (2.9)$$

Also we have

$$\frac{1}{\sup_x A^{1/p(x)} B^{1/q(x)}} \int_0^\infty |f(x) g(x)| dx \leq \int_0^\infty \frac{|f(x) g(x)|}{A^{1/p(x)} B^{1/q(x)}} dx. \quad (2.10)$$

From (2.9) and (2.10), we get (2.6), that is

$$\int_0^\infty |f(x) g(x)| dx \leq \alpha \beta.$$

We now prove (b).

Let $p_1(x) = 1/p(x)$, so that $p_1(x) > 1$ for all x . Let

$$A(x) = |g(x)|^{-1/p_1(x)}, \quad B(x) = |f(x)g(x)|^{1/p_1(x)}.$$

So by (2.6)

$$\begin{aligned} \int_0^\infty |f(x)|^{p(x)} dx &= \int_0^\infty |A(x)B(x)| dx \\ &\leq \sup_x \left[\left(\int_0^\infty |A(x)|^{q_1(x)} dx \right)^{1/q_1(x)} \left(\int_0^\infty |B(x)|^{p_1(x)} dx \right)^{1/p_1(x)} \right] \\ &\quad \times \left(\sup_x \frac{1}{p_1(x)} + \sup_x \frac{1}{q_1(x)} \right), \end{aligned} \quad (2.11)$$

since $\frac{1}{p_1(x)} + \frac{1}{q_1(x)} = 1$, $\frac{1}{q_1(x)} = 1 - p(x)$ and $q_1(x) = \frac{1}{1-p(x)}$.

Substituting the values in (2.11), we get (2.7)

$$\begin{aligned} \int_0^\infty |f(x)|^{p(x)} dx &= \sup_x \left[\left(\int_0^\infty |g(x)|^{q(x)} dx \right)^{1-p(x)} \left(\int_0^\infty |f(x)g(x)| dx \right)^{p(x)} \right] \\ &\quad \times \left[\sup_x p(x) + \sup_x (1 - p(x)) \right] \\ &= \alpha [\sup_x p(x) + \sup_x (1 - p(x))]. \end{aligned}$$

This completes the proof of the lemma. \square

Note. Equality occurs in (2.6) when

$$(1) \alpha = 1.$$

$$(2) \beta |f(x)|^{p(x)} = \gamma |g(x)|^{q(x)}, \text{ where } \beta \text{ and } \gamma \text{ are constants not all zero.}$$

But $\alpha = 1$ holds only when $p(x)$ and $q(x)$ are constants.

If $p(x)$ and $q(x)$ are not constants and $\frac{1}{p(x)} + \frac{1}{q(x)} = 1$, then $\alpha = \sup \frac{1}{p(x)} + \sup \frac{1}{q(x)} > 1$ e.g.

For $p(x) = 1 + \frac{1}{x} > 1$, $x \in [2, 3]$ and $q(x) = x + 1 > 1$, $x \in [2, 3]$ not constants then $\alpha = \sup \frac{1}{p(x)} + \sup \frac{1}{q(x)} > 1$.

3. Main results

LEMMA 3.1. Let $(p_k)_{k=1}^{\infty}$ and $(q_k)_{k=1}^{\infty}$ be real bounded sequences defined by $\frac{1}{p_k} + \frac{1}{q_k} = 1$ where $p_k > 1$, $0 < \epsilon < q_k - 1$ for all $k \in \mathbb{N}$, then we have for any $k \geq 1$

$$\begin{aligned} & \sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \frac{1}{\ln(mn)} \frac{1}{m(\ln m)^{(1+\epsilon)/p_k}} \frac{1}{n(\ln n)^{(1+\epsilon)/q_k}} \\ & > \frac{1}{\epsilon} \left[\frac{\pi}{\sin(\pi/p_k)} + o(1) \right] \quad (\text{as } \epsilon \rightarrow 0^+). \end{aligned} \quad (3.1)$$

Proof. Obviously, we have

$$\begin{aligned} & \sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \frac{1}{\ln(mn)} \frac{1}{m(\ln m)^{(1+\epsilon)/p_k}} \frac{1}{n(\ln n)^{(1+\epsilon)/q_k}} \\ & > \int_e^{\infty} \int_e^{\infty} \frac{1}{\ln(xy)} \frac{1}{x(\ln x)^{(1+\epsilon)/p_k}} \frac{1}{y(\ln y)^{(1+\epsilon)/q_k}} dy dx. \end{aligned} \quad (3.2)$$

Setting $u = \frac{\ln y}{\ln x}$ in the following expression, for $x \geq e$ and $0 < \epsilon < q_k - 1$ we obtain for $k = 1, 2, \dots$

$$\begin{aligned} & \int_e^{\infty} \frac{1}{\ln(xy)} \frac{1}{y(\ln y)^{(1+\epsilon)/q_k}} dy \\ & = \frac{1}{(\ln x)^{(1+\epsilon)/q_k}} \int_{1/\ln x}^{\infty} \frac{1}{1+u} \frac{1}{u^{(1+\epsilon)/q_k}} du \\ & = \frac{1}{(\ln x)^{(1+\epsilon)/q_k}} \left[\int_0^{\infty} \frac{1}{1+u} \frac{1}{u^{(1+\epsilon)/q_k}} du - \int_0^{1/\ln x} \frac{1}{1+u} \frac{1}{u^{(1+\epsilon)/q_k}} du \right] \\ & > \frac{1}{(\ln x)^{(1+\epsilon)/q_k}} \left[\int_0^{\infty} \frac{1}{1+u} \frac{1}{u^{(1+\epsilon)/q_k}} du - \int_0^{1/\ln x} \frac{1}{u^{(1+\epsilon)/q_k}} du \right] \\ & = \frac{1}{(\ln x)^{(1+\epsilon)/q_k}} \left[\frac{\pi}{\sin(\pi/p_k)} + o(1) \right] - \frac{q_k}{(q_k - 1 - \epsilon)} \frac{1}{\ln x}. \end{aligned}$$

Hence we find for $k \geq 1$

$$\begin{aligned} & \int_e^\infty \int_e^\infty \frac{1}{\ln(xy)} \frac{1}{x(\ln x)^{(1+\epsilon)/p_k}} \frac{1}{y(\ln y)^{(1+\epsilon)/q_k}} dy dx \\ & > \int_e^\infty \frac{1}{x(\ln x)^{(1+\epsilon)}} \left[\frac{\pi}{\sin(\pi/p_k)} + o(1) \right] dx - \frac{q_k}{(q_k - 1 - \epsilon)} \int_e^\infty \frac{1}{x(\ln x)^{1+(1+\epsilon)/p_k}} dx \\ & = \left[\frac{\pi}{\sin(\pi/p_k)} + o(1) \right] \frac{1}{\epsilon} - \frac{q_k}{(q_k - 1 - \epsilon)(1 + \epsilon)} \quad (\epsilon \rightarrow 0), \end{aligned} \quad (3.3)$$

by (3.2) and (3.3), it follows that (3.1) is valid. Hence the lemma. \square

THEOREM 3.1. *If $a_n, b_n \geq 0$, (p_k) and (q_k) are real bounded sequences defined by $\frac{1}{p_k} + \frac{1}{q_k} = 1$, where $p_k > 1$ for all $k \in \mathbf{N}$, and $0 < \sum_{n=2}^\infty n^{p_k-1} a_n^{p_k} < \infty$, $0 < \sum_{n=2}^\infty n^{q_k-1} b_n^{q_k} < \infty$. Then*

$$\sum_{n=2}^\infty \sum_{m=2}^\infty \frac{a_m b_n}{\ln(mn)} < \alpha \sup_k \left\{ \frac{\pi}{\sin(\pi/p_k)} \left\{ \sum_{n=2}^\infty n^{p_k-1} a_n^{p_k} \right\}^{1/p_k} \left\{ \sum_{n=2}^\infty n^{q_k-1} b_n^{q_k} \right\}^{1/q_k} \right\}, \quad (3.4)$$

where $\alpha = \sup_{k \geq 1} \frac{1}{p_k} + \sup_{k \geq 1} \frac{1}{q_k}$.

Proof. By Holder's inequality and (1.4), we have

$$\begin{aligned} & \sum_{n=2}^\infty \sum_{m=2}^\infty \frac{a_m b_n}{\ln mn} \\ & = \sum_{n=2}^\infty \sum_{m=2}^\infty \left[\frac{a_m}{(\ln mn)^{\frac{1}{p_k}}} \left(\frac{\ln m}{\ln n} \right)^{\frac{1}{p_k q_k}} \left(\frac{m^{1/q_k}}{n^{1/p_k}} \right) \right] \left[\frac{b_n}{(\ln mn)^{1/q_k}} \left(\frac{\ln n}{\ln m} \right)^{\frac{1}{p_k q_k}} \left(\frac{n^{1/p_k}}{m^{1/q_k}} \right) \right] \\ & \leq \alpha \sup_{k \geq 1} \left[\left(\sum_{m=2}^\infty \sum_{n=2}^\infty \frac{a_m^{p_k}}{\ln mn} \left(\frac{\ln m}{\ln n} \right)^{\frac{1}{q_k}} \right. \right. \\ & \quad \times \left(\frac{m^{p_k}}{n} \right)^{\frac{1}{p_k}} \left(\sum_{n=2}^\infty \sum_{m=2}^\infty \frac{b_n^{q_k}}{\ln mn} \left(\frac{\ln n}{\ln m} \right)^{\frac{1}{p_k}} \left(\frac{n^{q_k}}{m} \right)^{\frac{1}{q_k}} \right)^{\frac{1}{q_k}} \left. \right] \\ & = \alpha \sup_{k \geq 1} \left[\left(\sum_{m=2}^\infty \sum_{n=2}^\infty \frac{1}{\ln mn} \left(\frac{\ln m}{\ln n} \right)^{\frac{1}{q_k}} \left(\frac{m^{p_k-1}}{n} \right) a_m^{p_k} \right)^{\frac{1}{p_k}} \right. \\ & \quad \times \left. \left(\sum_{n=2}^\infty \sum_{m=2}^\infty \frac{1}{\ln mn} \left(\frac{\ln n}{\ln m} \right)^{\frac{1}{p_k}} \left(\frac{n^{q_k-1}}{m} \right) b_n^{q_k} \right)^{\frac{1}{q_k}} \right] \\ & = \alpha \sup_{k \geq 1} \left[\left\{ \sum_{m=2}^\infty w_{q_k}(m) m^{p_k-1} a_m^{p_k} \right\}^{\frac{1}{p_k}} \left\{ \sum_{n=2}^\infty w_{p_k}(n) n^{q_k-1} b_n^{q_k} \right\}^{\frac{1}{q_k}} \right]. \end{aligned} \quad (3.5)$$

For $r_k = p_k, q_k$ for all k , and $n \geq 2$ in (1.4), setting $u = \frac{\ln x}{\ln n}$ in the following integral, we find

$$w_{r_k}(n) < \int_1^\infty \frac{1}{x \ln nx} \left(\frac{\ln n}{\ln x} \right)^{\frac{1}{r_k}} dx = \int_0^\infty \frac{1}{(1+u)u^{\frac{1}{r_k}}} du = \frac{\pi}{\sin \pi(1 - \frac{1}{r_k})}. \quad (3.6)$$

Since $\sin(\pi/p_k) = \sin(\pi/q_k)$, for all k , by (3.5) and (3.6), we have (3.4). \square

THEOREM 3.2. Let $a_n \geq 0, (p_k)$ and (q_k) be real bounded sequences defined by $\frac{1}{p_k} + \frac{1}{q_k} = 1$, where $p_k > 1$ for all $k \in \mathbf{N}$, and $0 < \sum_{n=2}^\infty n^{p_k-1} a_n^{p_k} < \infty$. Then

$$\sum_{n=2}^\infty \frac{1}{n} \left(\sum_{m=2}^\infty \frac{a_m}{\ln(mn)} \right)^{p_k} < \alpha^{\sup p_k} \sup_{k \geq 1} \left\{ \left[\frac{\pi}{\sin(\pi/p_k)} \right]^{p_k} \sum_{n=2}^\infty n^{p_k-1} a_n^{p_k} \right\}, \quad (3.7)$$

where $\alpha = \sup_{n \geq 1} \frac{1}{p_n} + \sup_{n \geq 1} \frac{1}{q_n}$.

Proof. Since $\sum_{n=2}^\infty n^{p_k-1} a_n^{p_k} > 0$, there exists $t_0 \geq 2$, such that for any $t > t_0$, $\sum_{n=2}^t n^{p_k-1} a_n^{p_k} > 0$, and $b_n(t) = \frac{1}{n} \left(\sum_{m=2}^t \frac{a_m}{\ln(mn)} \right)^{p_k-1} > 0$. Then we have

$$0 < \sum_{n=2}^t n^{q_k-1} b_n^{q_k}(t) = \sum_{n=2}^t \frac{1}{n} \left(\sum_{m=2}^t \frac{a_m}{\ln(mn)} \right)^{p_k} = \sum_{n=2}^t \sum_{m=2}^t \frac{a_m b_n(t)}{\ln(mn)}. \quad (3.8)$$

If we set $\tilde{a}_n = a_n$ and $\tilde{b}_n = b_n(t)$, for $n = 2, 3, \dots, t$; and $\tilde{a}_n = \tilde{b}_n = 0$, for $n > t$. By using (3.4), we may have

$$\begin{aligned} \sum_{n=2}^t \sum_{m=2}^t \frac{a_m b_n(t)}{\ln(mn)} &= \sum_{n=2}^\infty \sum_{m=2}^\infty \frac{\tilde{a}_m \tilde{b}_n}{\ln(mn)} \\ &< \alpha \sup_{k \geq 1} \left[\frac{\pi}{\sin(\pi/p_k)} \left\{ \sum_{n=2}^\infty n^{p_k-1} \tilde{a}_n^{p_k} \right\}^{\frac{1}{p_k}} \left\{ \sum_{n=2}^\infty n^{q_k-1} \tilde{b}_n^{q_k} \right\}^{\frac{1}{q_k}} \right] \\ &= \alpha \sup_{k \geq 1} \left[\frac{\pi}{\sin(\pi/p_k)} \left\{ \sum_{n=2}^t n^{p_k-1} a_n^{p_k} \right\}^{\frac{1}{p_k}} \left\{ \sum_{n=2}^t n^{q_k-1} b_n^{q_k}(t) \right\}^{\frac{1}{q_k}} \right]. \end{aligned}$$

Hence by (3.8), we have

$$\begin{aligned} 0 < \sum_{n=2}^\infty n^{q_k-1} b_n^{q_k}(t) &= \sum_{n=2}^t \frac{1}{n} \left(\sum_{m=2}^t \frac{a_m}{\ln mn} \right)^{p_k} \\ &< \sup_{k \geq 1} \left[\frac{\alpha \pi}{\sin(\pi/p_k)} \right]^{p_k} \sum_{n=2}^t n^{p_k-1} a_n^{p_k} \\ &= \alpha^{\sup p_k} \sup_{k \geq 1} \left\{ \left[\frac{\pi}{\sin(\pi/p_k)} \right]^{p_k} \sum_{n=2}^t n^{p_k-1} a_n^{p_k} \right\}. \end{aligned}$$

It follows that $0 < \sum_{n=2}^t n^{q_k-1} b_n^{q_k}(\infty) < \infty$, since $0 < \sum_{n=2}^t n^{p_k-1} a_n^{p_k} < \infty$. Hence by (3.4) we have,

$$\begin{aligned} & \sum_{n=2}^t \frac{1}{n} \left(\sum_{m=2}^{\infty} \frac{a_m}{\ln(mn)} \right)^{p_k} \\ &= \sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \frac{a_m b_n(\infty)}{\ln mn} \\ &< \sup_{k \geq 1} \left[\frac{\alpha \pi}{\sin(\pi/p_k)} \left\{ \sum_{n=2}^{\infty} n^{p_k-1} a_n^{p_k} \right\}^{\frac{1}{p_k}} \left\{ \sum_{n=2}^{\infty} n^{q_k-1} b_n^{q_k}(\infty) \right\}^{\frac{1}{q_k}} \right] \\ &< \sup_{k \geq 1} \left[\frac{\alpha \pi}{\sin(\pi/p_k)} \left\{ \sum_{n=2}^{\infty} n^{p_k-1} a_n^{p_k} \right\}^{\frac{1}{p_k}} \left\{ \sum_{n=2}^{\infty} \frac{1}{n} \left(\sum_{m=2}^{\infty} \frac{a_m}{\ln mn} \right)^{p_k} \right\}^{\frac{1}{q_k}} \right] \end{aligned}$$

by simplification, we have (3.7). \square

4. Associated Integral Inequalities

Define the weight function as in [4] $w_r(x)$ as

$$w_r(x) = \int_1^{\infty} \frac{1}{y \ln(xy)} \left(\frac{\ln x}{\ln y} \right)^{\frac{1}{r(x)}} dy \quad (r(x) = p(x), q(x) \geq 2, x \geq 1). \quad (4.1)$$

Setting $u = \frac{\ln y}{\ln x}$ in (4.1), we have

$$\begin{aligned} w_r(x) &= \int_0^{\infty} \frac{1}{1+u} \left(\frac{1}{u} \right)^{\frac{1}{r(x)}} du = \frac{\pi}{\sin \pi(1 - \frac{1}{r(x)})} \\ &\quad (r(x) = p(x), q(x) \geq 2, x \geq 1). \end{aligned} \quad (4.2)$$

THEOREM 4.1. Let $f, g \geq 0, p(x)$ and $q(x)$ be real bounded measurable functions defined by $\frac{1}{p(x)} + \frac{1}{q(x)} = 1$ where $p(x) > 1$ for all $x \geq 1$, and $0 < \int_1^{\infty} x^{p(x)-1} f^{p(x)}(x) dx < \infty$, $0 < \int_1^{\infty} x^{q(x)-1} g^{q(x)}(x) dx < \infty$, then

$$\begin{aligned} \int_1^{\infty} \int_1^{\infty} \frac{f(x)g(y)}{\ln(xy)} dx dy &< \alpha \sup_{x \geq 1} \left[\frac{\pi}{\sin(\pi/p(x))} \left\{ \int_1^{\infty} x^{p(x)-1} f^{p(x)}(x) dx \right\}^{\frac{1}{p(x)}} \right. \\ &\quad \times \left. \left\{ \int_1^{\infty} x^{q(x)-1} g^{q(x)}(x) dx \right\}^{\frac{1}{q(x)}} \right], \end{aligned} \quad (4.3)$$

and

$$\begin{aligned} & \int_1^\infty \frac{1}{y} \left(\int_1^\infty \frac{f(x)}{\ln(xy)} dx \right)^{p(y)} dy \\ & < \alpha^{\sup p(x)} \sup_{x \geq 1} \left[\frac{\pi}{\sin(\pi/p(x))} \right]^{p(x)} \int_1^\infty x^{p(x)-1} f^{p(x)}(x) dx \end{aligned} \quad (4.4)$$

where $\alpha = \sup p(t)^{-1} + \sup q(t)^{-1}$.

Proof. By generalized Holder's inequality (2.6), we have

$$\begin{aligned} \int_1^\infty \int_1^\infty \frac{f(x)g(y)}{\ln(xy)} dx dy &= \int_1^\infty \int_1^\infty \left[\frac{f(x)}{(\ln xy)^{1/p(x)}} \left(\frac{\ln x}{\ln y} \right)^{\frac{1}{p(x)q(x)}} \frac{x^{1/q(x)}}{y^{1/p(x)}} \right] \\ &\quad \times \left[\frac{g(y)}{(\ln xy)^{1/q(x)}} \left(\frac{\ln y}{\ln x} \right)^{\frac{1}{p(x)q(x)}} \frac{y^{1/p(x)}}{x^{1/q(x)}} \right] dx dy \\ &\leq \alpha \sup_{x \geq 1} \left[\left\{ \int_1^\infty \int_1^\infty \frac{f^{p(x)}(x)}{\ln(xy)} \left(\frac{\ln x}{\ln y} \right)^{\frac{1}{q(x)}} \frac{x^{p(x)-1}}{y} dy dx \right\}^{\frac{1}{p(x)}} \right. \\ &\quad \times \left. \left\{ \int_1^\infty \int_1^\infty \frac{g^{q(y)}(y)}{\ln(xy)} \left(\frac{\ln y}{\ln x} \right)^{\frac{1}{p(y)}} \frac{y^{q(y)-1}}{x} dy dx \right\}^{\frac{1}{q(x)}} \right]. \end{aligned} \quad (4.5)$$

From (2.6) equality holds in (4.5), when $p(x)$ and $q(x)$ are constants p and q , and for p, q there exists numbers a and b , such that

$$\frac{af^p(x)}{\ln xy} \left(\frac{\ln x}{\ln y} \right)^{\frac{1}{q}} \frac{x^{p-1}}{y} = \frac{bg^q(y)}{\ln xy} \left(\frac{\ln y}{\ln x} \right)^{\frac{1}{p}} \frac{y^{q-1}}{x} \quad \text{a.e. in } (1, \infty) \times (1, \infty).$$

Then we have $ax^{p-1}f^p(x) \ln x = by^{q-1}g^q(y) \ln y$ a.e. in $(1, \infty) \times (1, \infty)$. That is

$$ax^{p-1}f^p(x) \ln x = by^{q-1}g^q(y) \ln y = \text{constant} \quad \text{a.e. in } (1, \infty) \times (1, \infty)$$

which contradicts the fact that $0 < \int_1^\infty x^{p-1}f^p(x) dx < \infty$ and $0 < \int_1^\infty y^{q-1}g^q(y) dy < \infty$. It follows that (4.5) takes the form of strict inequality. By eqn. (4.1)

$$\begin{aligned} \int_1^\infty \int_1^\infty \frac{f(x)g(y)}{\ln xy} dx dy &< \alpha \sup_{x \geq 1} \left[\left\{ \int_1^\infty w_q(x)x^{p(x)-1}f^{p(x)}(x) dx \right\}^{\frac{1}{p(x)}} \right. \\ &\quad \times \left. \left\{ \int_1^\infty w_p(y)y^{q(y)-1}g^{q(y)}(y) dy \right\}^{\frac{1}{q(x)}} \right]. \end{aligned} \quad (4.6)$$

Since $\int_1^\infty x^{p(x)-1}f^{p(x)}(x) dx > 0$, then there exists $T_0 \geq 1$ such that for any $T > T_0$, $\int_1^T x^{p(x)-1}f^{p(x)}(x) dx > 0$, and $g(y, T) = \frac{1}{y} \left(\int_1^T \frac{f(x)}{\ln xy} dx \right)^{p(y)-1} > 0$ ($y \in (1, \infty)$),

by (4.3) we have

$$\begin{aligned}
 0 < \int_1^T y^{q(y)-1} g^{q(y)}(y, T) dy &= \int_1^T \frac{1}{y} \left(\int_1^T \frac{f(x)}{\ln xy} dx \right)^{p(y)} dy \\
 &= \int_1^T \int_1^T \frac{f(x)g(y, T)}{\ln xy} dxdy \\
 &< \alpha \sup_{x \geq 1} \left[\frac{\pi}{\sin(\pi/p(x))} \left\{ \int_1^T x^{p(x)-1} f^{p(x)}(x) dx \right\}^{1/p(x)} \right. \\
 &\quad \times \left. \left\{ \int_1^T x^{q(x)-1} g^{q(x)}(x, T) dx \right\}^{1/q(x)} \right], \tag{4.7}
 \end{aligned}$$

Let us denote

$$B = B(T, q, g) = \int_1^T y^{q(y)-1} g^{q(y)}(y, T) dy, \quad A = A(T, p, f) = \int_1^T x^{p(x)-1} f^{p(x)}(x, T) dx.$$

Then (4.7) is

$$B < \sup_{x \geq 1} \left[\frac{\pi \alpha}{\sin(\frac{\pi}{p(x)})} A^{1/p(x)} B^{1/q(x)} \right]$$

If we denote $C = A/B$, then this is

$$1 < \sup_{x \geq 1} \left[\frac{\pi \alpha}{\sin(\frac{\pi}{p(x)})} C^{1/p(x)} \right]$$

hence $\exists x_0$ such that

$$1 < \left[\frac{\pi \alpha}{\sin(\frac{\pi}{p(x_0)})} C^{1/p(x_0)} \right]$$

and hence

$$1 < \left[\frac{\pi \alpha}{\sin(\frac{\pi}{p(x_0)})} \right]^{p(x_0)} C,$$

So

$$B < \alpha^{\sup p(x)} \sup_{x \geq 1} \left[\frac{\pi}{\sin(\frac{\pi}{p(x)})} \right]^{p(x)} A$$

(4.4) follows. □

REFERENCES

- [1] YANG BICHENG, AND GAO MINGZHE, *On a best value of Hardy-Hilbert's Inequality*, Adv. Math. **26**, No. 2(1997), 159–164.
- [2] YANG BICHENG, AND L. DEBNATH, *On new strengthened Hardy-Hilbert's Inequality*, Int. J. Math. Math. Sci. **21**, No. 1(1998), 403–408.
- [3] YANG BICHENG, *On an extension of Hardy-Hilbert's Inequality*, Chin. Annal. Math. 23A (2002), 247–254.
- [4] YANG BICHENG, *On a New Inequality Similar to Hardy-Hilbert's Inequality*, Math. Inequ. Appl. **6**, No. 1(2003), 37–34.

- [5] G. DAS, AND S. NANDA, *Absolute almost convergence*, Indian J. Maths, **34** (1992), 99–110.
- [6] G. H. HARDY, J. E. LITTLEWOOD AND G. POLYA, *Inequalities*, Cambridge University Press, Cambridge, MA, (1952).
- [7] I. J. MADDOX, *Elements of Functional Analysis*, Cambridge University Press, (1970).
- [8] GAO MINGZHE, AND YANG BICHENG, *On the extended Hilbert's Inequality*, Proc. Amer. Math. Soc. **126**, 3(1998), 751–759.
- [9] S. SIMONS, *The sequence spaces $l(p_\gamma)$ and $m(p_\nu)$* , Proc. London Math. Soc. **13** (1965), 422–436.
- [10] S. K. SUNANDA, C. NAHAK AND S. NANDA, *Some new generalizations of Hardy's integral inequality*, Int. J. Math. and Math. Sci. Vol. 2006, Article ID 19013, 1–15.

(Received May 1, 2006)

S. K. Sunanda
*Department of Mathematics
 Indian Institute of Technology
 Kharagpur, India*
e-mail: sunanda@maths.iitkgp.ernet.in

C. Nahak
*Department of Mathematics
 Indian Institute of Technology
 Kharagpur, India*
e-mail: cnahak@maths.iitkgp.ernet.in

S. Nanda
*Department of Mathematics
 Indian Institute of Technology
 Kharagpur, India*