

## ON A GENERALIZATION OF THE LIZORKIN THEOREM ON FOURIER MULTIPLIERS

L. O. SARYBEKOVA, T. V. TARARYKOVA AND N. T. TLEUKHANOVA

(Communicated by V. Burenkov)

*Abstract.* A generalization of the Lizorkin theorem on Fourier multipliers is proved. The proofs are based on using the so-called net spaces and interpolation theorems. An example is given of a Fourier multiplier which satisfies the assumptions of the generalized theorem but does not satisfy the assumptions of the Lizorkin theorem.

Let  $F$  and  $F^{-1}$  be the direct and the inverse Fourier transforms respectively, namely,

$$(Ff)(\xi) = \int_{-\infty}^{+\infty} e^{-ix\xi} f(x) dx,$$

$$(F^{-1}g)(x) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{ix\xi} g(\xi) d\xi.$$

Let  $1 \leq p \leq q \leq \infty$ . It is said that  $\varphi$  is a Fourier multiplier from  $L_p$  to  $L_q$ , briefly,  $\varphi \in m_p^q$ , if there exists  $c_1 > 0$  such that for every function  $f$  in Schwartz space  $S$  the following inequality holds

$$\|T_\varphi(f)\|_{L_q} \leq c_1 \|f\|_{L_p},$$

where  $T_\varphi(f) = F^{-1}\varphi Ff$ . This inequality allows defining by continuity  $T_\varphi(f)$  for every  $f$  in  $L_p$ .

The set  $m_p^q$  of all Fourier multipliers from  $L_p$  to  $L_q$  is a normed space with the norm

$$\|\varphi\|_{m_p^q} = \|T_\varphi\|_{L_p \rightarrow L_q}.$$

Our aim is to find smoothness and metric characteristics for a function  $\varphi$  to be a Fourier multiplier from  $L_p$  to  $L_q$ .

We recall the following properties of  $m_p^q$  spaces for  $1 < p \leq q < \infty$  (see, for example, [1]):

1.  $m_p^q = m_q^{p'}$ , where  $\frac{1}{p} + \frac{1}{p'} = 1$  and  $\frac{1}{q} + \frac{1}{q'} = 1$ ;

---

Mathematics subject classification (2010): 42B15, 42B35.

Keywords and phrases: Fourier multipliers, Lizorkin theorem.

2.  $m_p^p \hookrightarrow m_r^r$ , if  $\left| \frac{1}{r} - \frac{1}{2} \right| < \left| \frac{1}{p} - \frac{1}{2} \right|$ ; here the symbol  $\hookrightarrow$  denotes continuous embedding, i.e.  $X \hookrightarrow Y$ , where  $X$  and  $Y$  are normed spaces, means that  $X \subset Y$  and there exists  $c_2 > 0$  such that for every  $x \in X$

$$\|x\|_Y \leq c_2 \|x\|_X;$$

3.  $m_2^2 = L_\infty$ .

A function  $f$  is absolutely continuous on  $[a, b]$ , briefly  $f \in AC([a, b])$ , if for any  $\varepsilon > 0$  there exists  $\delta > 0$  such that for any finite system of disjoint intervals  $(a_k, b_k), k = 1, \dots, n$ , such that  $\sum_{k=1}^n (b_k - a_k) < \delta$

$$\sum_{k=1}^n |f(b_k) - f(a_k)| < \varepsilon.$$

If  $\Omega \subset \mathbb{R}$  is an open set, then  $AC^{loc}(\Omega)$  is the set of functions  $f$ , defined on  $\Omega$ , such that  $f \in AC([a, b])$  for every  $[a, b] \subset \Omega$ .

**THEOREM.** (Lizorkin [3]). *Let  $1 < p \leq q < \infty$ ,  $A > 0$  and a function  $\varphi \in AC^{loc}(\mathbb{R} \setminus \{0\})$  satisfy the following conditions*

$$\sup_{y \in \mathbb{R} \setminus \{0\}} |y|^{\frac{1}{p} - \frac{1}{q}} |\varphi(y)| \leq A$$

and

$$\text{ess sup}_{y \in \mathbb{R} \setminus \{0\}} |y|^{1 + \frac{1}{p} - \frac{1}{q}} |\varphi'(y)| \leq A.$$

Then the function  $\varphi \in m_p^q$  and

$$\|\varphi\|_{m_p^q} \leq c_3 A,$$

where  $c_3 > 0$  depends only on  $p$  and  $q$ .

Let  $0 \leq \alpha < \infty, 0 < \beta < \infty, a, b \in \mathbb{R}, 0 < q, r \leq \infty$ ,  $G_t = \{[a, b] : b - a \geq t\}$ .

The net space  $N^{\alpha, \beta, r}(L_q)$  is the set of all functions  $f$  in  $L_1$  for which

$$\|f\|_{N^{\alpha, \beta, r}(L_q)} = \left( \int_0^\infty \left( t^\alpha \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \right)^r \frac{dt}{t} \right)^{\frac{1}{r}} < \infty,$$

if  $0 < r < \infty$ , and if  $r = \infty$

$$\|f\|_{N^{\alpha, \beta, \infty}(L_q)} = \sup_{t>0} t^\alpha \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} < \infty,$$

where  $\chi_Q$  is the characteristic function of  $Q$ ,  $|Q|$  is the length of  $Q$ .

Similar spaces were considered in [4].

LEMMA 1. Let  $0 < \alpha < 1, 0 < \beta < 1, 0 < q \leq \infty$ .

a) If  $0 < r \leq r_1 \leq \infty$ , then

$$N^{\alpha,\beta,r}(L_q) \hookrightarrow N^{\alpha,\beta,r_1}(L_q).$$

b) If  $0 < \sigma < \min\{1 - \alpha, 1 - \beta\}, 0 < r \leq \infty$ , then

$$N^{\alpha,\beta,r}(L_q) \hookrightarrow N^{\alpha+\sigma,\beta+\sigma,r}(L_q). \quad (1)$$

*Proof.* First we prove that  $N^{\alpha,\beta,r}(L_q) \hookrightarrow N^{\alpha,\beta,\infty}(L_q)$ . Indeed,

$$\begin{aligned} \|f\|_{N^{\alpha,\beta,\infty}(L_q)} &= \sup_{t>0} t^\alpha \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \\ &= (\alpha r)^{\frac{1}{r}} \sup_{t>0} \left( \int_0^t x^{\alpha r - 1} dx \right)^{\frac{1}{r}} \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \\ &\leq (\alpha r)^{\frac{1}{r}} \left( \int_0^\infty \left( x^\alpha \sup_{Q \in G_x} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \right)^r \frac{dx}{x} \right)^{\frac{1}{r}} \\ &= (\alpha r)^{\frac{1}{r}} \|f\|_{N^{\alpha,\beta,r}(L_q)}. \end{aligned}$$

Using this inequality and the multiplicative inequality for spaces  $L_p$ , we obtain

$$\|f\|_{N^{\alpha,\beta,r_1}(L_q)} \leq \|f\|_{N^{\alpha,\beta,r}(L_q)}^{\frac{r}{r_1}} \|f\|_{N^{\alpha,\beta,\infty}(L_q)}^{1-\frac{r}{r_1}} \leq (\alpha r)^{\frac{1}{r}-\frac{1}{r_1}} \|f\|_{N^{\alpha,\beta,r}(L_q)}.$$

Moreover,

$$\begin{aligned} \|f\|_{N^{\alpha+\sigma,\beta+\sigma,r}(L_q)} &= \left( \int_0^\infty \left( t^{\alpha+\sigma} \sup_{Q \in G_t} \frac{1}{|Q|^{\beta+\sigma}} \|F^{-1} \chi_Q F f\|_{L_q} \right)^r \frac{dt}{t} \right)^{\frac{1}{r}} \\ &\leq \left( \int_0^\infty \left( t^\alpha \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \right)^r \frac{dt}{t} \right)^{\frac{1}{r}} = \|f\|_{N^{\alpha,\beta,r}(L_q)}. \end{aligned}$$

Lemma 1 is proved.  $\square$

Let  $(A_0, A_1)$  be a compatible pair of Banach spaces [1], and

$$K(\tau, a; A_0, A_1) = \inf_{a=a_0+a_1} (\|a_0\|_{A_0} + \tau \|a_1\|_{A_1}), \quad a \in A_0 + A_1, \quad \tau > 0$$

be the Peetre functional.

Moreover, let for  $0 < q < \infty, 0 < \theta < 1$

$$(A_0, A_1)_{\theta,q} = \left\{ a \in A_0 + A_1 : \|a\|_{(A_0, A_1)_{\theta,q}} = \left( \int_0^\infty (\tau^{-\theta} K(\tau, a; A_0, A_1))^q \frac{d\tau}{\tau} \right)^{\frac{1}{q}} < \infty \right\},$$

and for  $q = \infty$

$$(A_0, A_1)_{\theta, \infty} = \left\{ a \in A_0 + A_1 : \|a\|_{(A_0, A_1)_{\theta, \infty}} = \sup_{0 < \tau < \infty} \tau^{-\theta} K(\tau, a; A_0, A_1) < \infty \right\}.$$

LEMMA 2. Let  $0 < \alpha_1 < 1, 0 < \beta < 1, 0 < r, q \leq \infty$ . Then

$$(N^{\alpha_1, \beta, \infty}(L_q), N^{0, \beta, \infty}(L_q))_{\theta, r} \hookrightarrow N^{\alpha, \beta, r}(L_q)$$

for  $0 < \theta < 1$  and  $\alpha = (1 - \theta)\alpha_1$ .

*Proof.* Let  $f = f_1 + f_0$ ,  $f_0 \in N^{0, \beta, \infty}(L_q)$ ,  $f_1 \in N^{\alpha_1, \beta, \infty}(L_q)$ . It is clear that

$$\begin{aligned} & \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \\ & \leq 2^{\left(\frac{1}{q}-1\right)_+} \left( \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f_1\|_{L_q} + \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f_0\|_{L_q} \right), \end{aligned}$$

where  $x_+ = x$ , if  $x > 0$  and  $x_+ = 0$ , if  $x \leq 0$ . Denoting  $v(\tau) = \tau^{\frac{1}{\alpha_1}}$ ,  $\tau > 0$ , we obtain

$$\begin{aligned} & \sup_{0 < t \leq v(\tau)} t^{\alpha_1} \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \\ & \leq 2^{\left(\frac{1}{q}-1\right)_+} \left( \sup_{0 < t \leq v(\tau)} t^{\alpha_1} \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f_1\|_{L_q} + \sup_{0 < t \leq v(\tau)} t^{\alpha_1} \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f_0\|_{L_q} \right) \\ & \leq 2^{\left(\frac{1}{q}-1\right)_+} \left( \sup_{t > 0} t^{\alpha_1} \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f_1\|_{L_q} + \tau \sup_{t > 0} \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f_0\|_{L_q} \right). \end{aligned}$$

Since the representation  $f = f_0 + f_1$  is arbitrary

$$\sup_{0 < t \leq v(\tau)} t^{\alpha_1} \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \leq 2^{\left(\frac{1}{q}-1\right)_+} K(\tau, f; N^{\alpha_1, \beta, \infty}, N^{0, \beta, \infty}).$$

Therefore for  $0 < r < \infty$  we get

$$\begin{aligned} & \left( \int_0^\infty \left( \tau^{-\theta} K(\tau, f; N^{\alpha_1, \beta, \infty}, N^{0, \beta, \infty}) \right)^r \frac{d\tau}{\tau} \right)^{\frac{1}{r}} \\ & \geq 2^{-\left(\frac{1}{q}-1\right)_+} \left( \int_0^\infty \left( \tau^{-\theta} \sup_{t \leq v(\tau)} t^{\alpha_1} \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \right)^r \frac{d\tau}{\tau} \right)^{\frac{1}{r}} \\ & = 2^{-\left(\frac{1}{q}-1\right)_+} \left( \alpha_1 \int_0^\infty \left( u^{-\theta \alpha_1} \sup_{t \leq u} t^{\alpha_1} \sup_{Q \in G_u} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \right)^r \frac{du}{u} \right)^{\frac{1}{r}} \\ & \geq 2^{-\left(\frac{1}{q}-1\right)_+} \alpha_1^{\frac{1}{r}} \left( \int_0^\infty \left( u^{-\theta \alpha_1} \sup_{t \leq u} t^{\alpha_1} \sup_{Q \in G_u} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q F f\|_{L_q} \right)^r \frac{du}{u} \right)^{\frac{1}{r}} \end{aligned}$$

$$\geq 2^{-(\frac{1}{q}-1)_+} \alpha_1^{\frac{1}{r}} \|f\|_{N^{\alpha,\beta,r}(L_q)},$$

i.e.

$$(N^{\alpha_1,\beta,\infty}(L_q), N^{0,\beta,\infty}(L_q))_{\theta,r} \hookrightarrow N^{\alpha,\beta,r}(L_q).$$

The argument for  $r = \infty$  is similar. Lemma 2 is proved.  $\square$

Let  $f$  be a Lebesgue measurable function on  $\mathbb{R}$ . The distribution function of  $f$  is defined by

$$m(\sigma, f) = |\{x \in \mathbb{R} : |f(x)| > \sigma\}|.$$

The function

$$f^*(t) = \inf\{\sigma \geq 0 : m(\sigma, f) \leq t\}$$

is the non-increasing rearrangement of  $f$ .

Let  $1 \leq p < \infty, 0 < \tau \leq \infty$ . A function  $f$  belongs to the Lorentz space  $L_{p\tau}$ , if  $f$  is measurable on  $\mathbb{R}$  and for  $\tau < \infty$

$$\|f\|_{L_{p\tau}} = \left( \int_0^\infty \left( t^{\frac{1}{p}} f^*(t) \right)^\tau \frac{dt}{t} \right)^{\frac{1}{\tau}} < \infty$$

and for  $\tau = \infty$

$$\|f\|_{L_{p\infty}} = \sup_{t>0} t^{\frac{1}{p}} f^*(t) < \infty.$$

**THEOREM 1.** Let  $1 < p < q \leq \infty, 0 < \tau \leq \infty$ , and  $0 \leq \alpha < 1 - \frac{1}{p} + \frac{1}{q}$ . Then

$$L_{p\tau} \hookrightarrow N^{\alpha,\beta,\tau}(L_q),$$

where  $\beta = \alpha + \frac{1}{p} - \frac{1}{q}$

*Proof.* Let  $1 \leq r < q \leq \infty$ . It is well known that the characteristic function  $\chi_Q$  of a segment  $Q$  is a Fourier multiplier from  $L_r$  to  $L_q$ . Moreover, there exists  $c_4(r, q) > 0$ , depending only on  $r$  and  $q$ , such that

$$\|F^{-1}\chi_Q Ff\|_{L_q} \leq c_4(r, q) |Q|^{\frac{1}{r} - \frac{1}{q}} \|f\|_{L_r}, \quad (2)$$

for every  $f \in L_r$ .

Let  $0 \leq \alpha \leq 1 - \frac{1}{p}$ . Then  $\frac{1}{p} - \frac{1}{q} < \beta \leq 1 - \frac{1}{q}$  and there exists  $p_0$  such that  $1 < p_0 < p$  and  $\beta = \frac{1}{p_0} - \frac{1}{q}$ . By (2) with  $r = p_0$

$$\|f\|_{N^{0,\beta,\infty}(L_q)} = \sup_{t>0} \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1}\chi_Q Ff\|_{L_q} \leq c_5 \|f\|_{L_{p_0}} \quad (3)$$

for every  $f \in L_{p_0}$ , where  $c_5 = c_4(p_0, q)$ .

Next let  $p < p_1 < q$  and  $\alpha_1 = \frac{1}{p_0} - \frac{1}{p_1}$ . Taking into account (2) with  $r = p_1$ , we get

$$\sup_{t>0} \sup_{Q \in G_t} \frac{1}{|Q|^{\frac{1}{p_1} - \frac{1}{q}}} \|F^{-1}\chi_Q Ff\|_{L_q} \leq c_6 \|f\|_{L_{p_1}}$$

for every  $f \in L_{p_1}$ , where  $c_6 = c_4(p_1, q)$ .

Since

$$\begin{aligned} \sup_{t>0} \sup_{Q \in G_t} \frac{1}{|Q|^{\frac{1}{p_1}-\frac{1}{q}}} \|F^{-1}\chi_Q Ff\|_{L_q} &= \sup_{t>0} \sup_{Q \in G_t} \frac{|Q|^{\alpha_1}}{|Q|^\beta} \|F^{-1}\chi_Q Ff\|_{L_q} \\ &\geq \sup_{t>0} t^{\alpha_1} \sup_{Q \in G_t} \frac{1}{|Q|^\beta} \|F^{-1}\chi_Q Ff\|_{L_q} = \|f\|_{N^{\alpha_1, \beta, \infty}(L_q)}, \end{aligned}$$

we have

$$\|f\|_{N^{\alpha_1, \beta, \infty}(L_q)} \leq c_6 \|f\|_{L_{p_1}} \quad (4)$$

for every  $f \in L_{p_1}$ . Inequalities (3) and (4) mean that

$$L_{p_0} \hookrightarrow N^{0, \beta, \infty}(L_q),$$

$$L_{p_1} \hookrightarrow N^{\alpha_1, \beta, \infty}(L_q),$$

and consequently

$$L_{p_0} + L_{p_1} \hookrightarrow N^{0, \beta, \infty}(L_q) + N^{\alpha_1, \beta, \infty}(L_q).$$

Let us denote by  $I$  the corresponding embedding operator. By (3) and (4) we have

$$I : L_{p_0} \rightarrow N^{0, \beta, \infty}(L_q),$$

and

$$I : L_{p_1} \rightarrow N^{\alpha_1, \beta, \infty}(L_q).$$

Moreover, in both cases the operator  $I$  is bounded.

Let  $\theta \in (0, 1)$  such that  $\frac{1}{p} = \frac{1-\theta}{p_1} + \frac{\theta}{p_0}$ . Since

$$\alpha = \beta - \frac{1}{p} + \frac{1}{q} = \frac{1}{p_0} - \frac{1}{p} = (1-\theta)\alpha_1.$$

By the interpolation properties of  $L_p$  we get

$$I : L_{p\tau} = (L_{p_1}, L_{p_0})_{\theta\tau} \rightarrow \left( N^{\alpha_1, \beta, \infty}(L_q), N^{0, \beta, \infty}(L_q) \right)_{\theta\tau},$$

and the operator  $I$  is bounded. Thus,

$$L_{p\tau} \hookrightarrow \left( N^{\alpha_1, \beta, \infty}(L_q), N^{0, \beta, \infty}(L_q) \right)_{\theta\tau},$$

and hence the statement of theorem follows by Lemma 2.

If  $1 - \frac{1}{p} \leq \alpha < 1 - \frac{1}{p} + \frac{1}{q}$ , then the statement of theorem follows from (1). Indeed, let  $0 < \tilde{\alpha} \leq 1 - \frac{1}{p}$ , then by proved above and by (1) we have

$$L_{p\tau} \hookrightarrow N^{\tilde{\alpha}, \tilde{\alpha} + \frac{1}{p} - \frac{1}{q}, \tau}(L_q) \hookrightarrow N^{\alpha, \beta, \tau}(L_q).$$

Theorem 1 is proved.  $\square$

**THEOREM 2.** Let  $1 < p < q \leq \infty$ ,  $0 \leq \alpha < 1 - \frac{1}{p} + \frac{1}{q}$ ,  $\beta = \alpha + \frac{1}{p} - \frac{1}{q}$ . If a function  $\varphi \in AC^{loc}(\mathbb{R} \setminus \{0\})$  satisfies to the following conditions

$$\sup_{y \in \mathbb{R}} |y|^{\frac{1}{p} - \frac{1}{q}} |\varphi(y)| \leq A$$

and

$$\sup_{t > 0} t^{1-\alpha} \left( y^\beta \varphi'(y) \right)^* (t) \leq A,$$

then  $\varphi \in m_p^q$  and

$$\|\varphi\|_{m_p^q} \leq c_7 A,$$

where  $c_7 > 0$  depends only on  $p, q$  and  $\alpha$ .

*Proof.* Let  $f \in S$ ,  $0 < a < b < \infty$ . Since  $\varphi \in AC([a, b])$ , by applying integration by parts and the Minkowski inequalities for sums and integrals, we get

$$\begin{aligned} \left\| \int_a^b \varphi(y) (Ff)(y) e^{iyz} dy \right\|_{L_q} &= \left\| \int_a^b \varphi(y) \left( \int_0^y (Ff)(\xi) e^{i\xi z} d\xi \right)' dy \right\|_{L_q} \\ &= \left\| \varphi(y) \int_0^y (Ff)(\xi) e^{i\xi z} d\xi |_a^b - \int_a^b \varphi'(y) \left( \int_0^y (Ff)(\xi) e^{i\xi z} d\xi \right) dy \right\|_{L_q} \\ &= 2\pi \left\| \varphi(y) (F^{-1} \chi_{[0,y]} Ff)(z) |_a^b - \int_a^b \varphi'(y) (F^{-1} \chi_{[0,y]} Ff)(z) dy \right\|_{L_q} \\ &\leq 2\pi (\|\varphi(a) (F^{-1} \chi_{[0,a]} Ff)(z)\|_{L_q} + \|\varphi(b) (F^{-1} \chi_{[0,b]} Ff)(z)\|_{L_q} \\ &\quad + \int_a^b |\varphi'(y)| \|F^{-1} \chi_{[0,y]} Ff\|_{L_q} dy) \equiv 2\pi (I_1 + I_2 + I_3). \end{aligned}$$

Taking into account (2) we have

$$I_1 = a^{\frac{1}{p} - \frac{1}{q}} |\varphi(a)| \frac{1}{a^{\frac{1}{p} - \frac{1}{q}}} \|F^{-1} \chi_{[0,a]} Ff\|_{L_q} \leq A c_8 \|f\|_{L_p},$$

where  $c_8 = c_4(p, q)$ .

Similarly,

$$I_2 \leq A c_8 \|f\|_{L_p}.$$

Furthermore,

$$\begin{aligned} I_3 &= \int_a^b y^\beta |\varphi'(y)| \frac{1}{y^\beta} \|F^{-1} \chi_{[0,y]} Ff\|_{L_q} dy \\ &\leq \int_0^\infty y^\beta |\varphi'(y)| \left( \sup_{Q \in G_y} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q Ff\|_{L_q} \right) dy. \end{aligned}$$

Since

$$\sup_{Q \in G_y} \frac{1}{|Q|^\beta} \|F^{-1} \chi_Q Ff\|_{L_q}$$

is a nonincreasing function, by using the inequality  $\int_0^\infty FGdy \leq \int_0^\infty F^*G^*dy$ , we get

$$\begin{aligned} I_3 &\leq \int_0^\infty \left( z^\beta \varphi'(z) \right)^*(y) \left( \sup_{Q \in G_y} \frac{1}{|Q|^\beta} \|F^{-1}\chi_Q Ff\|_{L_q} \right) dy \\ &= \int_0^\infty y^{1-\alpha} \left( z^\beta \varphi'(z) \right)^*(y) y^\alpha \left( \sup_{Q \in G_y} \frac{1}{|Q|^\beta} \|F^{-1}\chi_Q Ff\|_{L_q} \right) \frac{dy}{y} \\ &\leq A \|f\|_{N^{\alpha, \beta, 1}(L_q)}. \end{aligned}$$

By Theorem 1 there exists  $c_9 > 0$ , which depends only on  $p, q, \alpha$ , such that

$$I_3 \leq A c_9 \|f\|_{L_{p1}}.$$

Since  $L_{p1} \hookrightarrow L_p$ , there exists  $c_{10} > 0$ , depending only on  $p$ , such that

$$\|f\|_{L_p} \leq c_{10} \|f\|_{L_{p1}}.$$

Hence, there exists  $c_{11} > 0$ , depending only on  $p, q$  and  $\alpha$ , such that

$$\left\| \int_a^b \varphi(y)(Ff)(y) e^{iyz} dy \right\|_{L_q} \leq c_{11} A \|f\|_{L_{p1}}$$

for every  $f \in S$  and any  $0 < a < b < \infty$ .

Let couples of numbers  $(p_0, q_0)$  and  $(p_1, q_1)$  be such that  $1 < p_0 < p < p_1 < \infty$ ,  $1 < q_0 < q < q_1 < \infty$  and

$$\frac{1}{p_0} - \frac{1}{q_0} = \frac{1}{p_1} - \frac{1}{q_1} = \frac{1}{p} - \frac{1}{q}. \quad (5)$$

Similarly, one can prove that there exists  $c_{12} > 0$  such that

$$\left\| \int_a^b \varphi(y)(Ff)(y) e^{iyz} dy \right\|_{L_{q_0}} \leq c_{12} A \|f\|_{L_{p_0}}$$

and

$$\left\| \int_a^b \varphi(y)(Ff)(y) e^{iyz} dy \right\|_{L_{q_1}} \leq c_{12} A \|f\|_{L_{p_1}}$$

for every  $f \in S$ . Choose  $\theta \in (0, 1)$  such that  $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ . According to (5) we have

$$\begin{aligned} \frac{1}{q} &= \frac{1}{p} - \frac{1}{p_0} + \frac{1}{q_0} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1} - \frac{1}{p_0} + \frac{1}{q_0} \\ &= \theta \left( \frac{1}{p_1} - \frac{1}{p_0} \right) + \frac{1}{q_0} = \theta \left( \frac{1}{q_1} - \frac{1}{q_0} \right) + \frac{1}{q_0} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}. \end{aligned}$$

By the Marcinkiewicz interpolation theorem we have that for every  $0 < r < \infty$

$$\left\| \int_a^b \varphi(y)(Ff)(y) e^{iyz} dy \right\|_{L_{qr}} \leq c_{12} A \|f\|_{L_{pr}},$$

where  $c_{12} > 0$  depends only on  $p, q, \alpha$  and  $r$ .

If, in particular,  $r = p$ , then for some  $c_{13} > 0$ , which depends only on  $p, q$  and  $\alpha$ ,

$$\left\| \int_a^b \varphi(y)(Ff)(y)e^{iyz} dy \right\|_{L_q} \leq c_{13}A \|f\|_{L_p}$$

for all  $f \in S$  and for all  $0 < a < b < \infty$ , because  $L_{qp} \hookrightarrow L_q$  if  $p < q$ .

Since  $|\varphi(y)| \leq A|y|^{-\frac{1}{p} + \frac{1}{q}}$ ,  $y \in \mathbb{R} \setminus \{0\}$  and  $Ff \in S$ , the integral

$$\int_0^\infty \varphi(y)(Ff)(y)e^{iyz} dy$$

absolutely converges for every  $z \in \mathbb{R}$ .

Hence, for every  $z \in \mathbb{R}$

$$\lim_{n \rightarrow \infty} \left| \int_{\frac{1}{n}}^n \varphi(y)(Ff)(y)e^{iyz} dy \right| = \left| \int_0^\infty \varphi(y)(Ff)(y)e^{iyz} dy \right|,$$

and by the Fatou theorem

$$\left\| \int_0^\infty \varphi(y)(Ff)(y)e^{iyz} dy \right\|_{L_q} \leq \sup_{n \in N} \left\| \int_{\frac{1}{n}}^n \varphi(y)(Ff)(y)e^{iyz} dy \right\|_{L_q} \leq c_{13}A \|f\|_{L_p}.$$

Similarly,

$$\left\| \int_{-\infty}^0 \varphi(y)(Ff)(y)e^{iyz} dy \right\|_{L_q} \leq c_{13}A \|f\|_{L_p},$$

and therefore

$$\|T_\varphi(f)\|_{L_q} = \|F^{-1}\varphi Ff\|_{L_q} = \frac{1}{2\pi} \left\| \int_{-\infty}^\infty \varphi(y)(Ff)(y)e^{iyz} dy \right\|_{L_q} \leq \frac{c_{13}A}{\pi} \|f\|_{L_p}$$

for every  $f \in S$ . This means that

$$\|T_\varphi(f)\|_{L_q} \leq \frac{c_{13}A}{\pi} \|f\|_{L_p}$$

for every  $f \in L_p$ . Hence  $\varphi \in m_p^q$  and

$$\|\varphi\|_{m_p^q} \leq \frac{c_{13}A}{\pi}.$$

Theorem 2 is proved.  $\square$

The function

$$f^*(t) = \inf\{\sigma > 0 : \tilde{m}(\sigma, f) \geq t\},$$

where

$$\tilde{m}(\sigma, f) = |\{x \in \mathbb{R} : |f(x)| < \sigma\}|,$$

is the non-decreasing rearrangement of  $f$ .

LEMMA 3. Let  $f, g$  be measurable functions on  $\mathbb{R}$  such that  $f^*(t), g^*(t) < \infty$  for all  $t > 0$ . Then

$$\sup_{t>0} f^*(t)g^*(t) \leq \text{ess sup}_{x \in \mathbb{R}} |f(x)g(x)|.$$

*Proof.* Let  $1 \leq p < \infty$ . Since  $(|g|^p)^*(t) = (g^*(t))^p$ ,  $(|f|^p)^*(t) = (f^*(t))^p$ , we have by the well-known inequality [2] that for every  $a > 0$

$$\|f^*g^*\|_{L_p(0,2a)} = \|(f\chi_{[-a,a]})^*g^*\|_{L_p(0,\infty)} \leq \|f\chi_{(-a,a)}g\|_{L_p(-\infty,\infty)} = \|fg\|_{L_p(-a,a)}.$$

Passing to the limit as  $p \rightarrow +\infty$  by the Riesz lemma, we get

$$\text{ess sup}_{0 < t < 2a} f^*(t)g^*(t) \leq \text{ess sup}_{-a < x < a} |f(x)g(x)|.$$

Passing to the limit as  $a \rightarrow +\infty$ , we obtain the required inequality because

$$\text{ess sup}_{t>0} f^*(t)g^*(t) = \sup_{t>0} f^*(t)g^*(t).$$

Lemma 3 is proved.  $\square$

REMARK 1. Note that the assumptions of Theorem 2 are weaker than the assumptions of the Lizorkin theorem, since

$$\sup_{t \in \mathbb{R}^+} t^{1-\alpha} \left( y^\beta \varphi'(y) \right)^*(t) \leq 2^{1-\alpha} \text{ess sup}_{y \in \mathbb{R}} |y|^{1+\frac{1}{p}-\frac{1}{q}} |\varphi'(y)|,$$

and there exists a function  $\varphi$  satisfying the assumptions of Theorem 2, but not satisfying the assumptions of the Lizorkin theorem, i.e.

$$\sup_{y \in \mathbb{R}} |y|^{\frac{1}{p}-\frac{1}{q}} |\varphi(y)| < \infty,$$

$$\sup_{t \in \mathbb{R}^+} t^{1-\alpha} \left( y^\beta \varphi'(y) \right)^*(t) < \infty,$$

but

$$\text{ess sup}_{y \in \mathbb{R}} |y|^{1+\frac{1}{p}-\frac{1}{q}} |\varphi'(y)| = \infty.$$

Since  $(|y|^{1-\alpha})^*(t) = (\frac{t}{2})^{1-\alpha}$ , by Lemma 3 the first statement of the remark follows.

Let us prove the second statement. Let  $1 > \gamma > \beta$  and

$$\mu_k = \left( k^{1-\beta} + k^{-\gamma} - (k+1)^{-\gamma} \right)^{\frac{1}{1-\beta}}, \quad k \in N.$$

We define the function  $\varphi$  by the following formula

$$\varphi(x) = \begin{cases} k^{1-\beta} + k^{-\gamma} - x^{1-\beta}, & x \in [k, \mu_k], \\ (k+1)^{-\gamma}, & x \in [\mu_k, k+1], \end{cases} \quad k \in N,$$

for  $x \geq 1$ . If  $0 \leq x \leq 1$ ,

$$\varphi(x) = 1,$$

and if  $x < 0$ ,

$$\varphi(x) = \varphi(-x).$$

The function  $\varphi \in AC^{loc}(\mathbb{R} \setminus \{0\})$ , because it is continuous on  $\mathbb{R}$ , is even by definition and is absolutely continuous on any interval  $[0, 1]$ ,  $[k, \mu_k]$ ,  $[\mu_k, k+1]$ , where  $k \in N$ .

We show that the function  $\varphi$  satisfies the first condition. Let  $x \in [k, k+1]$ . Since  $\gamma > \frac{1}{p} - \frac{1}{q}$  we have

$$|x|^{\frac{1}{p}-\frac{1}{q}} |\varphi(x)| \leq (k+1)^{\frac{1}{p}-\frac{1}{q}} \frac{1}{k^\gamma} \leq \frac{(k+1)^{\frac{1}{p}-\frac{1}{q}}}{k^{\frac{1}{p}-\frac{1}{q}}} \leq 2.$$

Note that

$$|x^\beta \varphi'(x)| = \begin{cases} 1-\beta, & x \in (k, \mu_k), \\ 0, & x \in (\mu_k, k+1), \end{cases} \quad k \in N.$$

If

$$I = \left\{ x : |x^\beta \varphi'(x)| = 1-\beta \right\},$$

then

$$|I| = \sum_{k=1}^{\infty} (\mu_k - k) = \sum_{k=1}^{\infty} k \left( \left( 1 + \frac{1}{k^{1-\beta}} (k^{-\gamma} - (k+1)^{-\gamma}) \right)^{\frac{1}{1-\beta}} - 1 \right) < \infty.$$

Indeed, by using inequality  $(1+x)^\nu - 1 \leq \nu 2^{\nu-1} x$  for  $\nu \geq 1$  and  $0 \leq x \leq 1$ , we have

$$|I| \leq \frac{2^{\frac{\beta}{1-\beta}}}{1-\beta} \sum_{k=1}^{\infty} \frac{k}{k^{1-\beta}} (k^{-\gamma} - (k+1)^{-\gamma}).$$

Since the series  $\sum_{k=1}^{\infty} k^{-(1-\beta+\gamma)}$  converges, by the limit comparison test the series

$$\sum_{k=1}^{\infty} \frac{k}{k^{1-\beta}} (k^{-\gamma} - (k+1)^{-\gamma})$$

also converges.

Since the function  $|x^\beta \varphi'(x)|$  takes only two values 0 and  $1-\beta$ , it follows that

$$(x^\beta \varphi'(x))^*(t) = \begin{cases} 1-\beta, & x \in [0, |I|], \\ 0, & x > |I|. \end{cases}$$

Therefore,

$$\sup_{t>0} t^{1-\alpha} \cdot (x^\beta \varphi'(x))^*(t) = \sup_{0 < t \leq 2d} t^{1-\alpha} \cdot (1-\beta) = (1-\beta) (|I|)^{1-\alpha} < \infty,$$

i.e. the second condition holds.

On the other hand, if we choose an arbitrary sequence of points  $\{x_k\}_{k=1}^{\infty}$  such that  $x_k \in (k, \mu_k)$ , then

$$\begin{aligned}\lim_{k \rightarrow \infty} |x_k|^{1+\frac{1}{p}-\frac{1}{q}} \cdot |\varphi'(x_k)| &= \lim_{k \rightarrow \infty} |x_k|^{1+\frac{1}{p}-\frac{1}{q}-\beta} \cdot |x_k|^{\beta} |\varphi'(x_k)| \\ &= \lim_{k \rightarrow \infty} |x_k|^{1+\frac{1}{p}-\frac{1}{q}-\beta} \cdot (1-\beta) = \infty.\end{aligned}$$

Hence, the last condition holds.

## REFERENCES

- [1] J. BERGH, J. LEFSTREM, *Interpolation spaces. An Introduction*, Springer-Verlag, Berlin Heidelberg New York, 1976.
- [2] G. H. HARDY, J. E. LITTLEWOOD, G. POLYA, *Inequalities*, Cambridge University Press, 1934.
- [3] P. I. LIZORKIN, *On multipliers of Fourier integrals in  $L_{p,\theta}$  spaces*, Trudy Matematicheskogo Instituta imeni V. A. Steklova, 1967, Vol. 139 (in Russian). English transl. in Proceeding of the Steklov institute of mathematics, 1967, Vol. 139.
- [4] E. D. NURSULTANOV, *Nikol'skii's inequality for different metrics and properties of the sequence of norms of the Fourier sums of a function in the Lorentz space*, Trudy Matematicheskogo Instituta imeni V. A. Steklova, 2006, Vol. 255, pp. 1970–215 (in Russian). English transl. in Proceeding of the Steklov institute of mathematics, 2006, Vol. 255.

(Received October 17, 2008)

*L. O. Sarybekova,  
L. N. Gumilyov Eurasian National University  
5 Munaitpasov St  
Astana 010008  
Kazakhstan  
e-mail: lsarybekova@yandex.ru*

*T. V. Tararykova  
L. N. Gumilyov Eurasian National University  
5 Munaitpasov St  
Astana 010008  
Kazakhstan  
e-mail: tararykovat@cardiff.ac.uk*

*N. T. Tleukhanova  
L. N. Gumilyov Eurasian National University  
5 Munaitpasov St  
Astana 010008  
Kazakhstan  
e-mail: tleukhanova@yandex.ru*