

AN OPERATOR EXTENSION OF THE PARALLELOGRAM LAW AND RELATED NORM INEQUALITIES

MOHAMMAD SAL MOSLEHIAN

*Dedicated to my dear teacher
 Aziz Atai Langroudi
 with respect and affection*

Abstract. We establish a general operator parallelogram law concerning a characterization of inner product spaces, get an operator extension of Bohr's inequality and present several norm inequalities. More precisely, let \mathfrak{A} be a C^* -algebra, T be a locally compact Hausdorff space equipped with a Radon measure μ and let $(A_t)_{t \in T}$ be a continuous field of operators in \mathfrak{A} such that the function $t \mapsto A_t$ is norm continuous on T and the function $t \mapsto \|A_t\|$ is integrable. If $\alpha : T \times T \rightarrow \mathbb{C}$ is a measurable function such that $\overline{\alpha(t,s)}\alpha(s,t) = 1$ for all $t, s \in T$, then we show that

$$\int_T \int_T |\alpha(t,s)A_t - \alpha(s,t)A_s|^2 d\mu(t)d\mu(s) + \int_T \int_T |\alpha(t,s)B_t - \alpha(s,t)B_s|^2 d\mu(t)d\mu(s) \\ = 2 \int_T \int_T |\alpha(t,s)A_t - \alpha(s,t)B_s|^2 d\mu(t)d\mu(s) - 2 \left| \int_T (A_t - B_t) d\mu(t) \right|^2.$$

1. Introduction

Let \mathfrak{A} be a C^* -algebra and let T be a locally compact Hausdorff space. A field $(A_t)_{t \in T}$ of operators in \mathfrak{A} is called a continuous field of operators if the function $t \mapsto A_t$ is norm continuous on T . If $\mu(t)$ is a Radon measure on T and the function $t \mapsto \|A_t\|$ is integrable, one can form the Bochner integral $\int_T A_t d\mu(t)$, which is the unique element in \mathfrak{A} such that

$$\varphi \left(\int_T A_t d\mu(t) \right) = \int_T \varphi(A_t) d\mu(t)$$

for every linear functional φ in the norm dual \mathfrak{A}^* of \mathfrak{A} ; see [8, Section 4.1] and [7].

Let $\mathbb{B}(\mathcal{H})$ be the algebra of all bounded linear operators on a separable complex Hilbert space \mathcal{H} endowed with inner product $\langle \cdot, \cdot \rangle$. We denote the absolute value of $A \in \mathbb{B}(\mathcal{H})$ by $|A| = (A^*A)^{1/2}$. For $x, y \in \mathcal{H}$, the rank one operator $x \otimes y$ is defined on \mathcal{H} by $(x \otimes y)(z) = \langle z, y \rangle x$.

Mathematics subject classification (2010): Primary 47A63; Secondary 46C15, 47A30, 47B10, 47B15, 15A60.

Keywords and phrases: Bochner integral, Schatten p -norm, norm inequality, parallelogram law, unitarily invariant norm, convex function, Bohr's inequality, inner product space.

Let $A \in \mathbb{B}(\mathcal{H})$ be a compact operator and let $0 < p < \infty$. The Schatten p -norm (p -quasi-norm) for $1 \leq p < \infty$ ($0 < p < 1$) is defined by $\|A\|_p = (\text{tr}|A|^p)^{1/p}$, where tr is the usual trace functional. Clearly

$$\| |A|^p \|_1 = \|A\|_p^p \tag{1.1}$$

for $p > 0$. For $p > 0$, the Schatten p -class, denoted by \mathcal{C}_p , is defined to be the two-sided ideal in $\mathbb{B}(\mathcal{H})$ of those compact operators A for which $\|A\|_p$ is finite. In particular, \mathcal{C}_1 and \mathcal{C}_2 are the trace class and the Hilbert-Schmidt class, respectively. For $1 \leq p < \infty$, \mathcal{C}_p is a Banach space; in particular the triangle inequality holds. However, for $0 < p < 1$, the quasi-norm $\|\cdot\|_p$ does not satisfy the triangle inequality. In addition to Schatten p -norms and operator norm, there are other interesting norms defined on some ideals contained in the ideal of compact operators. A unitarily invariant norm $\|\cdot\|$ is defined only on a norm ideal $\mathcal{C}_{\|\cdot\|}$ associated with it and has the property $\| |UAV| \| = \| |A| \|$, where U and V are unitaries and $A \in \mathcal{C}_{\|\cdot\|}$. For more information on the theory of the unitarily invariant norms the reader is referred to [15].

One can show that

$$\sum_{i,j=1}^n \|x_i - x_j\|^2 + \sum_{i,j=1}^n \|y_i - y_j\|^2 = 2 \sum_{i,j=1}^n \|x_i - y_j\|^2 - 2 \left\| \sum_{i=1}^n (x_i - y_i) \right\|^2 \tag{1.2}$$

holds in an inner product space, which is indeed a generalization of the classical *parallelogram law*:

$$|z + w|^2 + |z - w|^2 = 2|z|^2 + 2|w|^2 \quad (z, w \in \mathbb{C}).$$

There are several extensions of parallelogram law among them we could refer the interested reader to [4, 5, 12, 17, 6]. Generalizations of the parallelogram law for the Schatten p -norms have been given in the form of the celebrated Clarkson inequalities (see [10] and references therein). Since \mathcal{C}_2 is a Hilbert space under the inner product $\langle A, B \rangle = \text{tr}(B^*A)$, it follows from (1.2) that if $A_1, \dots, A_n, B_1, \dots, B_n \in \mathcal{C}_2$ with $\sum_{i=1}^n (A_i - B_i) = 0$, then

$$\sum_{i,j=1}^n \|A_i - A_j\|_2^2 + \sum_{i,j=1}^n \|B_i - B_j\|_2^2 = 2 \sum_{i,j=1}^n \|A_i - B_j\|_2^2. \tag{1.3}$$

The classical *Bohr's inequality* states that for any $z, w \in \mathbb{C}$ and any positive real numbers r, s with $\frac{1}{r} + \frac{1}{s} = 1$,

$$|z + w|^2 \leq r|z|^2 + s|w|^2.$$

Many interesting operator generalizations of this inequality have been obtained; cf. [1, 3, 9, 14, 16].

In this paper, we establish an extended operator parallelogram law and get a generalization of Bohr's inequality. We also present several unitarily invariant and Schatten p -norm inequalities. Our results can be regarded as extensions of main results of [11].

2. Joint extensions of the parallelogram law and Bohr’s inequality

We start our work with following clear lemma. The first equality is called the operator parallelogram law.

LEMMA 2.1. *Let $A, B \in \mathbb{B}(\mathcal{H})$. Then*

$$|A + B|^2 + |A - B|^2 = 2|A|^2 + 2|B|^2 \quad (a);$$

and

$$|A + B|^2 - |A - B|^2 = 4\text{Re}(A^*B) \quad (b)$$

We now state our main result, which is an operator version of equality (1.2). As we will see later, it is indeed a joint operator extension of Bohr and parallelogram inequalities.

THEOREM 2.2. *Let \mathfrak{A} be a C^* -algebra, T be a locally compact Hausdorff space equipped with a Radon measure μ and let $(A_t)_{t \in T}$ be a continuous field of operators in \mathfrak{A} such that the function $t \mapsto A_t$ is norm continuous on T and the function $t \mapsto \|A_t\|$ is integrable. Let $\alpha : T \times T \rightarrow \mathbb{C}$ be a measurable function such that $\overline{\alpha(t, s)}\alpha(s, t) = 1$ for all $t, s \in T$. Then*

$$\begin{aligned} & \int_T \int_T |\alpha(t, s)A_t - \alpha(s, t)A_s|^2 d\mu(t)d\mu(s) + \int_T \int_T |\alpha(t, s)B_t - \alpha(s, t)B_s|^2 d\mu(t)d\mu(s) \\ &= 2 \int_T \int_T |\alpha(t, s)A_t - \alpha(s, t)B_s|^2 d\mu(t)d\mu(s) - 2 \left| \int_T (A_t - B_t) d\mu(t) \right|^2. \end{aligned}$$

Proof.

$$\begin{aligned} & \int_T \int_T |\alpha(t, s)A_t - \alpha(s, t)A_s|^2 d\mu(t)d\mu(s) + \int_T \int_T |\alpha(t, s)B_t - \alpha(s, t)B_s|^2 d\mu(t)d\mu(s) \\ &= \int_T \int_T (|\alpha(t, s)A_t - \alpha(s, t)A_s|^2 + |\alpha(t, s)B_t - \alpha(s, t)B_s|^2) d\mu(t)d\mu(s) \\ &= \int_T \int_T \left(\frac{1}{2} |\alpha(t, s)A_t - \alpha(s, t)A_s + \alpha(t, s)B_t - \alpha(s, t)B_s|^2 \right. \\ & \quad \left. + \frac{1}{2} |\alpha(t, s)A_t - \alpha(s, t)A_s - \alpha(t, s)B_t + \alpha(s, t)B_s|^2 \right) d\mu(t)d\mu(s) \\ & \hspace{15em} \text{(by Lemma 2.1(a))} \\ &= \int_T \int_T \left(\frac{1}{2} |(\alpha(t, s)A_t - \alpha(s, t)B_s) - (\alpha(s, t)A_s - \alpha(t, s)B_t)|^2 \right. \\ & \quad \left. + \frac{1}{2} |(\alpha(t, s)A_t - \alpha(t, s)B_t) - (\alpha(s, t)A_s - \alpha(s, t)B_s)|^2 \right) d\mu(t)d\mu(s) \\ &= \int_T \int_T \left[(|\alpha(t, s)A_t - \alpha(s, t)B_s|^2 + |\alpha(s, t)A_s - \alpha(t, s)B_t|^2 \right. \end{aligned}$$

$$\begin{aligned}
 & -\frac{1}{2} \left| (\alpha(t,s)A_t - \alpha(s,t)B_s) + (\alpha(s,t)A_s - \alpha(t,s)B_t) \right|^2 \\
 & + \frac{1}{2} \left| (\alpha(t,s)A_t - \alpha(t,s)B_t) - (\alpha(s,t)A_s - \alpha(s,t)B_s) \right|^2 \Big] d\mu(t)d\mu(s) \\
 & \hspace{15em} \text{(by Lemma 2.1(a))} \\
 & = \int_T \int_T |\alpha(t,s)A_t - \alpha(s,t)B_s|^2 d\mu(t)d\mu(s) + \int_T \int_T |\alpha(t,s)A_t - \alpha(s,t)B_s|^2 d\mu(t)d\mu(s) \\
 & - \frac{1}{2} \int_T \int_T \left(|(\alpha(t,s)A_t - \alpha(t,s)B_t) + (\alpha(s,t)A_s - \alpha(s,t)B_s)|^2 \right. \\
 & \quad \left. - |(\alpha(t,s)A_t - \alpha(t,s)B_t) - (\alpha(s,t)A_s - \alpha(s,t)B_s)|^2 \right) d\mu(t)d\mu(s) \\
 & = 2 \int_T \int_T |\alpha(t,s)A_t - \alpha(s,t)B_s|^2 d\mu(t)d\mu(s) \\
 & \quad - 2 \operatorname{Re} \int_T \int_T (\alpha(t,s)A_t - \alpha(t,s)B_t)^* (\alpha(s,t)A_s - \alpha(s,t)B_s) d\mu(t)d\mu(s) \\
 & \hspace{15em} \text{(by Lemma 2.1(b))} \\
 & = 2 \int_T \int_T |\alpha(t,s)A_t - \alpha(s,t)B_s|^2 d\mu(t)d\mu(s) \\
 & \quad - 2 \operatorname{Re} \left[\left(\int_T (A_t - B_t) d\mu(t) \right)^* \left(\int_T (A_s - B_s) d\mu(s) \right) \right] \\
 & = 2 \int_T \int_T |\alpha(t,s)A_t - \alpha(s,t)B_s|^2 d\mu(t)d\mu(s) - 2 \left| \int_T (A_t - B_t) d\mu(t) \right|^2. \quad \square
 \end{aligned}$$

If we let $T = \{1, \dots, n\}$, μ be the counting measure on T and $\alpha(i, j) = \sqrt{\frac{r_i}{r_j}}$, where $r_i > 0$ ($1 \leq i \leq n$), in Theorem 2.2, then we get

COROLLARY 2.3. (Generalized Parallelogram Law) *Let $A_1, \dots, A_n, B_1, \dots, B_n \in \mathbb{B}(\mathcal{H})$ and let r_1, \dots, r_n be positive numbers. Then*

$$\begin{aligned}
 & \sum_{1 \leq i < j \leq n} \left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right|^2 + \sum_{1 \leq i < j \leq n} \left| \sqrt{\frac{r_i}{r_j}} B_i - \sqrt{\frac{r_j}{r_i}} B_j \right|^2 \\
 & = \sum_{i,j=1}^n \left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} B_j \right|^2 - \left| \sum_{i=1}^n (A_i - B_i) \right|^2. \tag{2.1}
 \end{aligned}$$

If we set $B_1 = \dots = B_n = 0$ in Corollary 2.3, then the following extension of parallelogram law is obtained

COROLLARY 2.4. [6, Theorem 4.2] *Suppose that $A_1, \dots, A_n \in \mathbb{B}(\mathcal{H})$ and r_1, \dots, r_n are positive numbers with $\sum_{i=1}^n \frac{1}{r_i} = 1$. Then*

$$(0 \leq) \sum_{1 \leq i < j \leq n} \left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right|^2 = \sum_{i=1}^n r_i |A_i|^2 - \left| \sum_{i=1}^n A_i \right|^2. \tag{2.2}$$

REMARK 2.5. If $n = 2$ and $t := \frac{r_1}{r_2} > 0$, then operator equality (2.2) can be re-stated as the following form which is, as noted in [6], a generalization of [9, Theorem 1] and [3, Theorem 3].

$$|A_1 + A_2|^2 + \frac{1}{t}|tA_1 - A_2|^2 = (1 + t)|A_1|^2 + \left(1 + \frac{1}{t}\right)|A_2|^2.$$

We also infer the following extension of Bohr’s inequality [16, Theorem 7] from (2.2).

$$\left| \sum_{i=1}^n A_i \right|^2 \leq \sum_{i=1}^n r_i |A_i|^2.$$

The preceding inequality is, in turn, a special case of the following general form of the Bohr inequality being easily deduced from the fact that the left hand side of the operator equality in Theorem 2.2 is a positive element of the C^* -algebra \mathfrak{A} .

COROLLARY 2.6. (Generalized Operator Bohr’s Inequality) *Let \mathfrak{A} be a C^* -algebra, T be a locally compact Hausdorff space equipped with a Radon measure μ and let $(A_t)_{t \in T}$ be a continuous field of operators in \mathfrak{A} such that the function $t \mapsto A_t$ is norm continuous on T and the function $t \mapsto \|A_t\|$ is integrable. Let $\alpha : T \times T \rightarrow \mathbb{C}$ be a measurable function such that $\alpha(t, s)\alpha(s, t) = 1$ for all $t, s \in T$. Then*

$$\left| \int_T (A_t - B_t) d\mu(t) \right|^2 \leq \int_T \int_T |\alpha(t, s)A_t - \alpha(s, t)B_s|^2 d\mu(t) d\mu(s).$$

A weighted extension of norm equality (1.2) can be deduced from (2.1) as follows:

COROLLARY 2.7. *Let $x_1, \dots, x_n, y_1, \dots, y_n \in \mathcal{H}$ and let r_1, \dots, r_n be positive numbers. Then*

$$\begin{aligned} \sum_{1 \leq i < j \leq n} \left\| \sqrt{\frac{r_i}{r_j}} x_i - \sqrt{\frac{r_j}{r_i}} x_j \right\|^2 + \sum_{1 \leq i < j \leq n} \left\| \sqrt{\frac{r_i}{r_j}} y_i - \sqrt{\frac{r_j}{r_i}} y_j \right\|^2 \\ = \sum_{i,j=1}^n \left\| \sqrt{\frac{r_i}{r_j}} x_i - \sqrt{\frac{r_j}{r_i}} y_j \right\|^2 - \left\| \sum_{i=1}^n (x_i - y_i) \right\|^2. \end{aligned}$$

Proof. Let e be a non-zero vector of \mathcal{H} and set $A_i = x_i \otimes e, B_i = y_i \otimes e$ for $i = 1, \dots, n$. It follows from the elementary properties of rank one operators and equality (2.1) that

$$\begin{aligned} \left(\sum_{i,j=1}^n \left\| \sqrt{\frac{r_i}{r_j}} x_i - \sqrt{\frac{r_j}{r_i}} x_j \right\|^2 + \sum_{i,j=1}^n \left\| \sqrt{\frac{r_i}{r_j}} y_i - \sqrt{\frac{r_j}{r_i}} y_j \right\|^2 \right) e \otimes e \\ = \sum_{i,j=1}^n \left| \left(\sqrt{\frac{r_i}{r_j}} x_i - \sqrt{\frac{r_j}{r_i}} x_j \right) \otimes e \right|^2 + \sum_{i,j=1}^n \left| \left(\sqrt{\frac{r_i}{r_j}} y_i - \sqrt{\frac{r_j}{r_i}} y_j \right) \otimes e \right|^2 \end{aligned}$$

$$\begin{aligned}
 &= \sum_{i,j=1}^n \left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right|^2 + \sum_{i,j=1}^n \left| \sqrt{\frac{r_i}{r_j}} B_i - \sqrt{\frac{r_j}{r_i}} B_j \right|^2 \\
 &= 2 \sum_{i,j=1}^n \left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} B_j \right|^2 - 2 \left| \sum_{i=1}^n (A_i - B_i) \right|^2 \\
 &= 2 \sum_{i,j=1}^n \left| \left(\sqrt{\frac{r_i}{r_j}} x_i - \sqrt{\frac{r_j}{r_i}} y_j \right) \otimes e \right|^2 - 2 \left| \sum_{i=1}^n (x_i - y_i) \otimes e \right|^2 \\
 &= \left(2 \sum_{i,j=1}^n \left\| \sqrt{\frac{r_i}{r_j}} x_i - \sqrt{\frac{r_j}{r_i}} y_j \right\|^2 - 2 \left\| \sum_{i=1}^n (x_i - y_i) \right\|^2 \right) e \otimes e,
 \end{aligned}$$

from which we conclude the result. \square

3. A general parallelogram law

An extension of a result of [2] for n -tuples may be stated as follows.

- If $f : [0, \infty) \rightarrow [0, \infty)$ is a convex function, then for any positive operators A_1, \dots, A_n , any nonnegative numbers $\alpha_1, \dots, \alpha_n$ with $\sum_{j=1}^n \alpha_j = 1$ and any unitarily invariant norm $\|\cdot\|$ on $\mathbb{B}(\mathcal{H})$

$$\left\| \sum_{j=1}^n \alpha_j f(A_j) \right\| \geq \left\| f \left(\sum_{j=1}^n \alpha_j A_j \right) \right\|. \tag{3.1}$$

The following result is also known [13]:

- If $f : [0, \infty) \rightarrow [0, \infty)$ is a convex function with $f(0) = 0$, then for any positive operators A_1, \dots, A_n and any unitarily invariant norm $\|\cdot\|$ on $\mathbb{B}(\mathcal{H})$

$$\left\| f \left(\sum_{j=1}^n A_j \right) \right\| \geq \left\| \sum_{j=1}^n f(A_j) \right\|. \tag{3.2}$$

The reverse inequalities hold for concave functions; see [10] for more details. We now prove another significant theorem.

THEOREM 3.1. *Let $A_1, \dots, A_n \in \mathcal{C}_{\|\cdot\|}$, r_1, \dots, r_n be positive real numbers with $\sum_{i=1}^n \frac{1}{r_i} = 1$, let g be a nonnegative convex function on $[0, \infty)$ such that $g(0) = 0$ and let $f(t) = g(t^2)$. Then*

$$\left\| \sum_{i=1}^n \frac{1}{r_i} f(|r_i A_i|) \right\| \geq \left\| \sum_{1 \leq i < j \leq n} f \left(\left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right| \right) + f \left(\left| \sum_{i=1}^n A_i \right| \right) \right\| \tag{3.3}$$

for all unitarily invariant norm. If g is concave, then the reverse of (3.3) holds.

Proof.

$$\begin{aligned}
 \left\| \left\| \sum_{i=1}^n \frac{1}{r_i} f(|r_i A_i|) \right\| \right\| &= \left\| \left\| \sum_{i=1}^n \frac{1}{r_i} g(|r_i A_i|^2) \right\| \right\| \\
 &\geq \left\| \left\| g \left(\sum_{i=1}^n \frac{1}{r_i} |r_i A_i|^2 \right) \right\| \right\| && \text{(by (3.1))} \\
 &= \left\| \left\| g \left(\sum_{1 \leq i < j \leq n} \left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right|^2 + \sum_{i=1}^n |A_i|^2 \right) \right\| \right\| && \text{(by (2.2))} \\
 &\geq \left\| \left\| \sum_{1 \leq i < j \leq n} g \left(\left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right|^2 \right) + g \left(\sum_{i=1}^n |A_i|^2 \right) \right\| \right\| && \text{(by (3.2))} \\
 &= \left\| \left\| \sum_{1 \leq i < j \leq n} f \left(\left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right| \right) + f \left(\sum_{i=1}^n |A_i| \right) \right\| \right\|. \quad \square
 \end{aligned}$$

The function $g(t) = t^p$ for $1 \leq p < \infty$ ($g(t) = t^p$ for $0 < p \leq 1$, resp.) is convex (concave, resp.) on $[0, \infty)$. Hence we get the following corollary.

COROLLARY 3.2. *Let $A_1, \dots, A_n \in \mathcal{C}_p$ and r_1, \dots, r_n be positive real numbers with $\sum_{i=1}^n \frac{1}{r_i} = 1$. Then*

$$\sum_{i=1}^n r_i^{p-1} \|A_i\|_p^p \geq \sum_{1 \leq i < j \leq n} \left\| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right\|_p^p + \left\| \sum_{i=1}^n A_i \right\|_p^p$$

for any $2 \leq p < \infty$. The reverse inequality holds for any $0 < p \leq 2$.

Proof. Let $2 \leq p < \infty$.

$$\begin{aligned}
 \sum_{i=1}^n r_i^{p-1} \|A_i\|_p^p &= \sum_{i=1}^n r_i^{p-1} \| |A_i|^p \|_1 && \text{(by (1.1))} \\
 &= \text{tr} \left(\sum_{i=1}^n r_i^{p-1} |A_i|^p \right) \\
 &= \left\| \left\| \sum_{i=1}^n \frac{1}{r_i} |r_i A_i|^p \right\| \right\|_1 \\
 &\geq \left\| \left\| \sum_{1 \leq i < j \leq n} \left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right|^p + \sum_{i=1}^n |A_i|^p \right\| \right\| \\
 &\quad \text{(by Theorem 3.1 for } g(t) = t^{\frac{p}{2}}; 2 \leq p < \infty)
 \end{aligned}$$

$$\begin{aligned}
&= \operatorname{tr} \left(\sum_{1 \leq i < j \leq n} \left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right|^p + \left| \sum_{i=1}^n A_i \right|^p \right) \\
&= \sum_{1 \leq i < j \leq n} \operatorname{tr} \left(\left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right|^p \right) + \operatorname{tr} \left(\left| \sum_{i=1}^n A_i \right|^p \right) \\
&= \sum_{1 \leq i < j \leq n} \left\| \left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right|^p \right\|_1 + \left\| \left| \sum_{i=1}^n A_i \right|^p \right\|_1 \\
&= \sum_{1 \leq i < j \leq n} \left\| \left| \sqrt{\frac{r_i}{r_j}} A_i - \sqrt{\frac{r_j}{r_i}} A_j \right|^p \right\|_p + \left\| \left| \sum_{i=1}^n A_i \right|^p \right\|_p. \tag{by (1.1)}
\end{aligned}$$

The proof for the reverse inequality is similar. \square

Acknowledgement. The author would like to sincerely thank the anonymous referee for carefully reading the manuscript and for several useful comments.

This research was supported by a grant from Ferdowsi University of Mashhad (No. MP88082MOS).

REFERENCES

- [1] S. ABRAMOVICH, J. BARIĆ AND J.E. PEČARIĆ, *A new proof of an inequality of Bohr for Hilbert space operators*, Linear Algebra Appl. **430**, 4 (2009), 1432–1435.
- [2] J.S. AUJLA AND F.C. SILVA, *Weak majorization inequalities and convex functions*, Linear Algebra Appl. **369** (2003), 217–233.
- [3] W.-S. CHEUNG AND J. PEČARIĆ, *Bohr's inequalities for Hilbert space operators*, J. Math. Anal. Appl. **323**, 1 (2006), 403–412.
- [4] Y.J. CHO, V. ČULJAK, M. MATIĆ AND J.E. PEČARIĆ, *On parallelogram law and Bohr's inequality in n -inner product spaces*, Inequality theory and applications, Vol. 2 (Chinju/Masan, 2001), 79–90, Nova Sci. Publ., Hauppauge, NY, 2003.
- [5] O. EĞECIOĞLU, *Parallelogram-law-type identities*, Linear Algebra Appl. **225** (1995), 1–12.
- [6] M. FUJII AND H. ZHOU, *Matrix order in Bohr inequality for operators*, Banach J. Math. Anal., **4**, 1 (2010), 21–27.
- [7] F. HANSEN, J.E. PEČARIĆ AND I. PERIĆ, *Jensen's operator inequality and its converses*, Math. Scand. **100**, 1 (2007), 61–73.
- [8] F. HANSEN AND G.K. PEDERSEN, *Jensen's operator inequality*, Bull. London Math. Soc. **35** (2003), 553–564.
- [9] O. HIRZALLAH, *Non-commutative operator Bohr inequality*, J. Math. Anal. Appl. **282** (2003), 578–583.
- [10] O. HIRZALLAH AND F. KITTANEH, *Non-commutative Clarkson inequalities for n -tuples of operators*, Integral Equations Operator Theory **60**, 3 (2008), 369–379.
- [11] O. HIRZALLAH, F. KITTANEH AND M.S. MOSLEHIAN, *Schatten p -norm inequalities related to a characterization of inner product spaces*, Math. Inequal. Appl. **13**, 2 (2010), 235–241.
- [12] M. KATO, *A note on a generalized parallelogram law and the Littlewood matrices*, Bull. Kyushu Inst. Tech. Math. Natur. Sci. **33** (1986), 37–39.

- [13] T. KOSEM, *Inequalities between $\|f(A+B)\|$ and $\|f(A) + f(B)\|$* , Linear Algebra Appl. **418**, 1 (2006), 153–160.
- [14] M.S. MOSLEHIAN, J.E. PEČARIĆ AND I. PERIĆ, *An operator extension of Bohr's inequality*, Bull. Iranian Math. Soc. **35**, 2 (2009), 67–74.
- [15] B. SIMON, *Trace Ideals and their Applications*, Cambridge University Press, Cambridge, 1979.
- [16] F. ZHANG, *On the Bohr inequality of operators*, J. Math. Anal. Appl. **333** (2007), 1264–1271.
- [17] W.L. ZENG, *An extension of the parallelogram characterization of inner product spaces*, J. Math. Res. Exposition **2**, 3 (1982), 23–25.

(Received March 5, 2011)

Mohammad Sal Moslehian
Department of Mathematics
Center of Excellence in Analysis on Algebraic Structures (CEAAS)
Ferdowsi University of Mashhad
P.O. Box 1159, Mashhad 91775
Iran
e-mail: moslehian@ferdowsi.um.ac.ir and moslehian@ams.org