

## FUGLEDE–PUTNAM’S THEOREM FOR $w$ -HYPONORMAL OPERATORS

A. BACHIR AND F. LOMBARKIA

**Abstract.** An asymmetric Fuglede-Putnam’s Theorem for  $w$ -hyponormal operators and dominant operators is proved, as a consequence of this result, we obtain that the range of the generalized derivation induced by the above classes of operators is orthogonal to its kernel.

*Mathematics subject classification (2010):* 47B47, 47A30, 47B20.

*Keywords and phrases:* Fuglede-Putnam theorem, hyponormal operator, dominant operator,  $w$ -hyponormal operator.

### REFERENCES

- [1] A. ALUTHGE, *On  $p$ -hyponormal operators for  $0 < p < 1$* , Integral Equations Operator Theory **13** (1990), 307–315.
- [2] A. ALUTHGE AND D. WANG,  *$w$ -hyponormal operators*, Integral Equations Operator Theory **36** (2000), 1–10.
- [3] A. ALUTHGE AND D. WANG, *An operator inequality which implies paranormality*, Math. Inequal. App. **2** (1999), 113–119.
- [4] J. H. ANDERSON AND C. FOIAS, *properties which normal operators share with normal derivations and related operators*, Pacific J. Math. **61** (1975), 313–325.
- [5] A. BACHIR, *Generalized Derivation*, SUT Journal of Mathematics **40**, 2 (2004), 111–116.
- [6] M. CHO, *Spectral properties of  $p$ -hyponormal operators for  $0 < p < \frac{1}{2}$* , Glasgow. Math. J. **36** (1992), 117–122.
- [7] M. CHO AND T. HURUYA,  *$P$ -hyponormal operators for  $0 < p < \frac{1}{2}$* , Comment. Math. Prace Math. **33** (1993), 23–29.
- [8] B. P. DUGGAL, *Quasi-similar  $p$ -hyponormal operators*, Integral Equation and Operator Theory **26** (1996), 338–345.
- [9] B. FUGLEDE, *A commutativity theorem for normal operators*, Proc. Nat. Acad. Sci. U. S. A. **36** (1950), 35–40.
- [10] F. HANSEN, *An inequality*, Math. Ann. **246** (1980), 249–250.
- [11] E. HEINZ, *Beiträge zur Störungstheorie der Spektralzerlegung*, Math. Ann. **123** (1951), 415–438.
- [12] M. ITO, T. YAMAZAKI, *Relation between two inequalities*  $\left(B^{\frac{r}{2}} A^p B^{\frac{r}{2}}\right)^{\frac{r}{p+r}} \geqslant B^r$  and  $A^p \geqslant \left(A^{\frac{p}{2}} B^r A^{\frac{p}{2}}\right)^{\frac{p}{p+r}}$  *and their applications*, Integral Equations Operator Theory **44** (2002), 442–450.
- [13] I. H. JEON, K. TANAHASHI AND A. UCHIYAMA, *On Quasimilarity for Log-Hyponormal operators*, Glasgow Math. J. **46** (2004), 169–176.
- [14] C. R. PUTNAM, *On normal operators in Hilbert space*, Amer. J. Math **73** (1951), 357–362.
- [15] J. G. STAMPFLI AND B. WADHWA, *on dominant operators*, Monatshefte. Für. Math. **84** (1977), 33–36.
- [16] K. TANAHASHI, *On log-hyponormal operators*, Integr. Equat. Oper. Th. **34** (1999), 364–372.
- [17] A. UCHIYAMA AND K. TANAHASHI, *Fuglede-Putnam’s theorem for  $p$ -hyponormal operators*, Glasgow. Math. J. **44** (2002), 397–416.
- [18] A. UCHIYAMA, *Berger-Shaw’s theorem fo  $p$ -hyponormal operators*, Integr. Equat. Oper. Th. **33** (1997), 307–315.
- [19] M. YANAGIDA, *Power of class  $wA(s,t)$  Operators Associated with Generalized Aluthge Transformation*, J. of Inequal. & Appl. **7** (2002), 143–168.