

HARDY TYPE INEQUALITIES FOR FRACTIONAL AND q -FRACTIONAL INTEGRAL OPERATORS

FETHI BOUZAFFOUR

(Communicated by J. Pečarić)

Abstract. Hardy type inequality for different fractional integral operators with sharp constants on finite intervals are given.

1. Introduction

The Hardy inequality is

$$\int_0^\infty x^{-\alpha p} |I^\alpha f(x)|^p dx \leq \left\{ \frac{\Gamma(1/p')}{\Gamma(\alpha + 1/p')} \right\}^p \int_0^\infty |f(x)|^p dx, \quad (1.1)$$
$$1 < p < \infty, \quad 1/p + 1/p' = 1,$$

where the Riemann-Liouville transform I^α is defined by

$$I^\alpha f(x) = \int_0^x \frac{(x-t)^{\alpha-1}}{\Gamma(\alpha)} f(t) dt.$$

Several generalization of this inequality for I^α with sharp constants on finite and infinite intervals are given in ([6], [7], ...). This paper is devoted to give analogues of the previous inequality for different fractional integral operators of Erdelyi-Kober operator type, discrete transform and basic analogue of Erdelyi-Kober operator. The method used is the same as in [6].

2. Hardy type inequality for Erdelyi-Kober fractional integral operator

The following lemma is used hereafter to establish Hardy type inequalities.

LEMMA 2.1. *Let U, V be Hilbert spaces and assume $A : U \rightarrow V$ to be an infinite-dimensional linear compact operator. If there exists an orthonormal basis $\{u_n\}_{n \in \mathbb{N}}$ of*

Mathematics subject classification (2010): 26A33, 26D10.

Keywords and phrases: Hardy inequality, fractional integral operator, q -fractional integral operator.

U, an orthonormal system (not necessary a basis) $\{v_n\}_{n \in \mathbb{N}}$ of V , a sequence of non-increasing positive numbers s_n such that

$$Au = \sum_{n \in \mathbb{N}} s_n (u, u_n)_U v_n,$$

for all $u \in U$, where $(., .)_U$ is inner product in U . Then s_0 is the norm of the operator A .

The proof of this lemma can be found in [6].

We consider the Erdelyi-Kober operator [3]

$$T_{\lambda, \mu} f(x) = \frac{x^{-\lambda-\mu} \Gamma(\lambda + \mu + 1)}{\Gamma(\lambda + 1) \Gamma(\mu)} \int_0^x t^\lambda (x-t)^{\mu-1} f(t) dt,$$

where f is a locally integrable function and $\lambda > -1$, $\mu > 0$.

The Jacobi polynomials are defined by [12]

$$P_n^{(\alpha, \beta)}(x) = \frac{(\alpha+1)_n}{n!} {}_2F_1\left(\begin{matrix} -n, n+\alpha+\beta+1 \\ \alpha+1 \end{matrix}; \frac{1-x}{2}\right),$$

and satisfy the orthogonality relations

$$\begin{aligned} & \int_{-1}^1 P_m^{(\alpha, \beta)}(x) P_n(\alpha, \beta)(x) (1-x)^\alpha (1+x)^\beta dx \\ &= \frac{2^{\alpha+\beta+1}}{2n+\alpha+\beta+1} \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\Gamma(n+\alpha+\beta+1)n!} \delta_{mn}, \end{aligned}$$

where $\alpha > -1$ and $\beta > -1$.

It will be convenient to use the slightly different Jacobi polynomials $J_n(x; \alpha, \beta)$, which are defined by

$$J_n(x; \alpha, \beta) = {}_2F_1\left(\begin{matrix} -n, n+\alpha+\beta+1 \\ \alpha+1 \end{matrix}; x\right).$$

THEOREM 2.1. *Let α , β and μ be real numbers such that*

$$\min(\alpha, \alpha+\beta+1, \beta-\mu, \mu-1) > -1.$$

Then for all $f \in L^2((0, 1), x^\alpha (1-x)^\beta)$ the following inequality

$$\int_0^1 x^{\alpha+\mu} (1-x)^{\beta-\mu} |T_{\alpha, \mu} f(x)|^2 dx \leq \frac{\Gamma(\alpha+\mu+1)\Gamma(\beta-\mu+1)}{\Gamma(\alpha+1)\Gamma(\beta+1)} \int_0^1 x^\alpha (1-x)^\beta |f(x)|^2 dx$$

is valid, with equality holds when $f(x) = 1$.

Proof. Let

$$\begin{aligned} u_n(x) &= \sqrt{\frac{(2n+\alpha+\beta+1)\Gamma(n+\alpha+\beta+1)\Gamma(n+\alpha+1)}{n!\Gamma(\alpha+1)^2\Gamma(n+\beta+1)}} J_n(x; \alpha, \beta), \\ v_n(x) &= \sqrt{\frac{(2n+\alpha+\beta+1)\Gamma(n+\alpha+\beta+1)\Gamma(n+\alpha+\mu+1)}{n!\Gamma(\alpha+\mu+1)^2\Gamma(n+\beta-\mu+1)}} J_n(x; \alpha+\mu, \beta-\mu), \\ s_n &= \frac{\Gamma(\alpha+\mu+1)}{\Gamma(\alpha+1)} \sqrt{\frac{\Gamma(n+\alpha+1)\Gamma(n+\beta-\mu+1)}{\Gamma(n+\alpha+\mu+1)\Gamma(n+\beta+1)}}. \end{aligned}$$

Then $\{u_n\}_{n \in \mathbb{N}}$ and $\{v_n\}_{n \in \mathbb{N}}$ are orthonormal basis in the spaces $L^2((0, 1), x^\alpha (1-x)^\beta)$ and $L^2((0, 1), x^{\alpha+\mu} (1-x)^{\beta-\mu})$.

From the following asymptotic expansions:

$$\frac{\Gamma(z+a)}{\Gamma(z+b)} \sim z^{a-b}, \text{ as } z \rightarrow \infty, \quad |\arg(z)| < \pi,$$

we obtain

$$s_n \sim \frac{\Gamma(\alpha+\mu+1)}{\Gamma(\alpha+1)} n^{-\mu}, \text{ as } n \rightarrow \infty.$$

On the other hand a simple calculation shows that

$$\left(\frac{s_{n+1}}{s_n} \right)^2 - 1 = -\mu \frac{2n+\alpha+\beta+1}{(n+\alpha+\mu+1)(n+\beta+1)} < 0 \quad (\mu > 0).$$

Hence, the sequence $\{s_n\}_{n \in \mathbb{N}}$ is positive decreasing and approaching 0.

The beta integral yields

$$T_{\alpha,\mu} x^n = \frac{(\alpha)_n}{(\alpha+\mu+1)_n} x^n.$$

Therefore, we have

$$T_{\alpha,\mu} J_n(\cdot; \alpha, \beta)(x) = J_n(x; \alpha+\mu, \beta-\mu),$$

and

$$T_{\beta,\mu} u_n = s_n v_n.$$

So the result is a consequence of Lemma 2.1. \square

An equivalent inequality can be obtained for the fractional integral operators $S_{\lambda,\mu}$, which are defined by [3]

$$S_{\lambda,\mu} f(x) = \frac{x^{-\lambda-\mu}\Gamma(\lambda+\mu+1)}{\Gamma(\lambda+1)\Gamma(\mu)} \int_x^1 (1-t)^\lambda (x-t)^{\mu-1} f(1-t) dt,$$

where f is a locally integrable function and $\lambda > -1$, $\mu > 0$.

Put

$$\begin{aligned} u_n(x) &= \sqrt{\frac{(2n+\alpha+\beta+1)\Gamma(n+\alpha+\beta+1)\Gamma(n+\alpha+1)}{n!\Gamma(\alpha+1)^2\Gamma(n+\beta+1)}} J_n(x; \alpha, \beta), \\ v_n(x) &= \sqrt{\frac{(2n+\alpha+\beta+1)\Gamma(n+\alpha+\beta+1)\Gamma(n+\alpha-\mu+1)}{n!\Gamma(\alpha-\mu+1)^2\Gamma(n+\beta-\mu+1)}} J_n(1-x; \alpha+\mu, \beta-\mu), \\ s_n &= \frac{\Gamma(\alpha-\mu+1)}{\Gamma(\alpha+1)} \sqrt{\frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+\mu+1)}{\Gamma(n+\alpha-\mu+1)\Gamma(n+\beta+1)}}, \end{aligned}$$

and using the beta integral formula to obtain

$$S_{\alpha,\mu} u_n = s_n v_n.$$

Therefore, from the Lemma 2.1, we get under the following condition

$$\min(\alpha, \alpha+\beta+1, \beta-\mu, \mu-1) > -1,$$

the inequality for $S_{\alpha,\mu}$

$$\int_0^1 x^{\beta+\mu} (1-x)^{\alpha-\mu} |S_{\beta,\mu} f(x)|^2 dx \leq \frac{\Gamma(\alpha-\mu+1)\Gamma(\beta+\mu+1)}{\Gamma(\alpha+1)\Gamma(\beta+1)} \int_0^1 x^\alpha (1-x)^\beta |f(x)|^2 dx.$$

Note that the equality holds when $f(x) = 1$.

3. Hardy type inequality for discrete transform

We consider the polynomials

$$\phi_N(x) = \sum_{k=0}^N \alpha_k x^k, \text{ with } \alpha_k \neq 0, N = 0, 1, \dots,$$

and we define a transform S_N on finite sequences $\{f(n), n = 0, \dots, N\}$ by

$$S_N[f; \phi_N, x] = \sum_{n=0}^N \frac{(-1)^n}{n!} \phi_N^{(n)}(x) f(n).$$

Then from the Taylor series it is easy to see

$$S_N\left[\binom{n}{j}, \phi_N, x\right] = (-1)^j \alpha_j x^j, \quad j = 0, \dots, N.$$

The transform $S_N[., \phi_N, x]$ (3.1) with $\phi_N(x) = (1-x)^N$, has the property

$$S_N[Q_j(x; \alpha, \beta, N), (1-x)^N, x] = J_j(x; \beta, \alpha), \quad j = 0, 1, \dots, N$$

where $Q_j(x, \alpha, \beta, N)$ are the Hahn polynomials [12] given by

$$Q_j(x, \alpha, \beta, N) = {}_3F_2\left(\begin{matrix} -j, j+\alpha+\beta+1, -x \\ \alpha+1, -N \end{matrix}; 1\right),$$

and satisfying for $\alpha > -1$ and $\beta > -1$, the orthogonality relations

$$\begin{aligned} \sum_{x=0}^N \binom{\alpha+x}{x} \binom{\beta+N-\alpha}{N-x} Q_m(x, \alpha, \beta, N) Q_n(x, \alpha, \beta, N) \\ = \frac{(-1)^n (n+\alpha+\beta+1)_{N+1} (\beta+1)_n n!}{(2n+\alpha+\beta+1)(\alpha+1)_n (-N)_n N!} \delta_{mn}. \end{aligned}$$

Let

$$\begin{aligned} u_j(x) &= \sqrt{\frac{N! (2j+\alpha+\beta+1) (-N)_j (\alpha+1)_j}{(-1)^j j! (\beta+1)_j (j+\alpha+\beta+1)_{N+1}}} Q_j(x, \alpha, \beta, N) \\ v_j(x) &= \sqrt{\frac{(2j+\alpha+\beta+1) \Gamma(j+\alpha+\beta+1) (\beta+1)_j}{j! \Gamma(\beta+1) \Gamma(j+\alpha+1)}} J_j(x, \beta, \alpha) \\ s_j &= \sqrt{\frac{N! (-N)_j (\alpha+1)_j \Gamma(\beta+1) \Gamma(j+\alpha+1)}{(-1)^j (\beta+1)_j^2 (j+\alpha+\beta+1)_{N+1} \Gamma(j+\alpha+\beta+1)}}. \end{aligned}$$

For $j = 0, 1, \dots, N$, we have

$$\begin{aligned} \frac{s_{j+1}}{s_j} &= \sqrt{\frac{N-j}{N+\alpha+\beta+2+j}} \frac{\alpha+j+1}{\beta+j+1} \\ &\leqslant \sqrt{\frac{N}{N+\alpha+\beta+2}} \left(1 + \frac{\alpha-\beta}{\beta+j+1}\right). \end{aligned}$$

Then, for $\beta \geq \alpha > -1$, we see that the sequence $\{s_j\}_{j=0, \dots, N}$ is decreasing and from Lemma 2.1, we obtain

$$\begin{aligned} \int_0^1 x^\beta (1-x)^\alpha \left| S_N[f, (1-x)^N, x] \right|^2 dx &\leq \frac{N! \Gamma(\beta+1) \Gamma(\alpha+1)}{(\alpha+\beta+1)_{N+1} \Gamma(\alpha+\beta+1)} \\ &\quad \times \sum_{n=0}^N \binom{\alpha+n}{n} \binom{N+\beta-\alpha}{N-n} |f(n)|^2. \end{aligned}$$

4. Hardy type inequality for q -fractional integral operator

Basic analogue of the Erdelyi-Kober fractional integral operator, defined by Al-Salam and Ismail (19.4.1, [8]) as

$$I^{\alpha, \eta} f(x) = \frac{\Gamma_q(\alpha+\eta)}{\Gamma_q(\alpha) \Gamma_q(\eta)} \int_0^1 t^{\alpha-1} \frac{(qt;q)_\infty}{(tq^\eta;q)_\infty} f(xt) d_q t, \quad \eta > 0, \alpha > 0,$$

where the q -integral of Jackson is defined by [4]

$$\int_0^a f(x) d_q x := (1-q)a \sum_{k=0}^{\infty} f(aq^k) q^k.$$

For complex a and $0 < q < 1$, the q -shift factorial are defined by [4]

$$(a;q)_0 := 1, \quad (a;q)_n := \prod_{k=0}^{n-1} (1 - aq^k), \quad n = 1, 2, \dots,$$

$$(a;q)_{\infty} := \lim_{n \rightarrow \infty} (a;q)_n.$$

The q -gamma function is defined by

$$\Gamma_q(z) := \frac{(q;q)_{\infty}}{(q^z;q)_{\infty}} (1-q)^{1-z}.$$

The basic hypergeometric series are defined by [4]

$${}_r\varphi_s \left(\begin{matrix} a_1, a_2, \dots, a_r \\ b_1, b_2, \dots, b_s \end{matrix} \middle| q, z \right) := \sum_{k=0}^{+\infty} \frac{(a_1, a_2, \dots, a_r; q)_k}{(b_1, b_2, \dots, b_s; q)_k} \left((-1)^k q^{\binom{k}{2}} \right)^{1+s-r} z^k,$$

where

$$(a_1, \dots, a_r; q)_n = (a_1; q)_n \dots (a_r; q)_n.$$

The little q -Jacobi polynomials are defined by

$$p_n(x; q^{\alpha}, q^{\beta}; q) := {}_2\phi_1 \left(\begin{matrix} q^{-n}, q^{\alpha+\beta+n+1} \\ q^{\alpha+1} \end{matrix} \middle| q, x \right),$$

which satisfy the orthogonality relations

$$\begin{aligned} \frac{(q^{\alpha+1}, q^{\beta+1}; q)_{\infty}}{(q^{\alpha+\beta+2}, q; q)_{\infty}} \sum_{k=0}^{\infty} p_m(q^k; q^{\alpha}, q^{\beta}; q) p_n(q^k; q^{\alpha}, q^{\beta}; q) q^{k(\alpha+1)} \frac{(q^{k+1}; q)_{\infty}}{(q^{\beta+k+1})_{\infty}} \\ = \frac{q^{n(\alpha+1)} (1 - q^{\alpha+\beta+1}) (q^{\beta+1}, q; q)_n}{(1 - q^{\alpha+\beta+2n+1}) (q^{\alpha+1}, q^{\alpha+\beta+1}; q)_n} \delta_{m,n}, \end{aligned}$$

where $\alpha > -1$ and $\beta > -1$, see[12].

THEOREM 4.1. Let α , β and η be real numbers such that

$$\min(\alpha, \beta - \eta, \eta - 1) > -1.$$

Then for all $f \in L^2 \left((0, 1), x^{\alpha} \frac{(qx; q)_{\infty}}{(xq^{\beta}; q)_{\infty}} d_q x \right)$, the inequality

$$\begin{aligned} & \int_0^1 x^{\alpha+\eta} \frac{(qx; q)_{\infty}}{(xq^{\beta-\eta}; q)_{\infty}} |I^{\alpha+1, \eta} f(x)|^2 d_q x \\ & \leq \frac{(q^{\alpha+1}, q^{\beta+1}; q)_{\infty}}{(q^{\alpha+\eta+1}, q^{\beta-\eta+1}; q)_{\infty}} \int_0^1 x^{\alpha} \frac{(qx; q)_{\infty}}{(xq^{\beta}; q)_{\infty}} |f(x)|^2 d_q x \end{aligned}$$

is valid, with equality holds when $f(x) = 1$.

Proof. When $f(x) = 1$ we have $I^{\alpha+1,\eta}f(x) = 1$ and the equality is easily verified.

Let

$$\begin{aligned} u_n(x) &= \sqrt{\frac{(q^{\alpha+1}, q^{\beta+1}; q)_\infty}{(1-q)(q, q^{\alpha+\beta+2}; q)_\infty} \frac{1-q^{\alpha+\beta+2n+1}}{(1-q^{\alpha+\beta+1})q^{n(\alpha+1)}} \frac{(q^{\alpha+1}, q^{\alpha+\beta+1}; q)_n}{(q, q^{\beta+1}; q)_n}} \\ &\quad \times p_n(x; q^\alpha, q^\beta; q) \\ v_n(x) &= \sqrt{\frac{(q^{\alpha+\eta+1}, q^{\beta-\eta+1}; q)_\infty}{(1-q)(q, q^{\alpha+\beta+2}; q)_\infty} \frac{1-q^{\alpha+\beta+2n+1}}{(1-q^{\alpha+\beta+1})q^{n(\alpha+\eta+1)}} \frac{(q^{\alpha+\eta+1}, q^{\alpha+\beta+1}; q)_n}{(q, q^{\beta-\eta+1}; q)_n}} \\ &\quad \times p_n(x; q^{\alpha+\eta}, q^{\beta-\eta}; q) \\ s_n &= \sqrt{\frac{(q^{\alpha+1}, q^{\beta+1}; q)_\infty q^{n\eta}}{(q^{\alpha+\eta+1}, q^{\beta-\eta+1}; q)_\infty} \frac{(q^{\alpha+1}, q^{\beta-\eta+1}; q)_n}{(q^{\beta+1}, q^{\alpha+\eta+1}; q)_n}}. \end{aligned}$$

Using now the formula (19.4.24, [8])

$$I^{\alpha+1,\eta} \left(p_n \left(\cdot; q^\alpha, q^\beta; q \right) \right) (x) = p_n \left(x; q^{\alpha+\eta}, q^{\beta-\eta}; q \right),$$

we get

$$I^{\alpha+1,\eta} u_n(x) = s_n v_n(x).$$

For $n = 0, 1, 2, \dots$, we have

$$\frac{s_{n+1}}{s_n} = \sqrt{q^\eta \frac{(1-q^{\alpha+n+1})(1-q^{\beta-\eta+n+1})}{(1-q^{\beta+n+1})(1-q^{\alpha+\eta+n+1})}},$$

and

$$\lim_{n \rightarrow \infty} s_n = 0.$$

Then, the condition

$$\min(\alpha, \beta - \eta, \eta - 1) > -1,$$

shows that the sequence $\{s_n\}$ is positive, decreasing and tending to 0.

Hence, from the Lemma 2.1, we have $\sqrt{\frac{(q^{\alpha+1}, q^{\beta+1}; q)_\infty}{(q^{\alpha+\eta+1}, q^{\beta-\eta+1}; q)_\infty}}$ is the norm of the operator $I^{\alpha+1,\eta}$. \square

Consider the family of operators, see ([8], 19.5.1)

$$S_r(f)(\cos \theta) = \frac{(q, r^2; q)_\infty}{2\pi} \int_0^\pi \frac{(e^{2i\phi}, e^{-2i\phi}; q)_\infty f(\cos \phi)}{(re^{i(\phi+\theta)}, re^{-i(\phi+\theta)}, re^{i(\phi-\theta)}, re^{-i(\phi+\theta)}; q)_\infty} d\phi, \quad r \in (0, 1).$$

The operators S_r satisfy the semigroup property [8]

$$S_r \circ S_t = S_{rt} \text{ for } r, t, rt \in (0, 1).$$

Let

$$\begin{aligned} u_n(x) &= \sqrt{\frac{(q, t_1 t_2; q)_\infty (t_1 t_2; q)_n}{2\pi (q; q)_n t_1^{2n}}} \frac{p_n(x; t_1, t_2 | q)}{(t_1 e^{i\theta}, t_1 e^{-i\theta}; q)_\infty}, \quad x = \cos \theta \\ v_n(x) &= \sqrt{\frac{(q, t_1 t_2; q)_\infty (t_1 t_2; q)_n}{2\pi (q; q)_n (r t_1)^{2n}}} \frac{p_n(x; r t_1, t_2/r | q)}{(r t_1 e^{i\theta}, r t_1 e^{-i\theta}; q)_\infty} \\ s_n &= r^n, \end{aligned}$$

where $p_n(x; t_1, t_2 | q)$ is the Al-Salam-Chihara polynomials defined by

$$p_n(x; t_1, t_2; q) := {}_3\phi_2 \left(\begin{matrix} q^{-n}, t_1 e^{i\theta}, t_2 e^{-i\theta} \\ t_1 t_2, 0 \end{matrix} \middle| q, q \right).$$

If $\max(|t_1|, |t_2|) < 1$, then we have the orthogonality relations [12]

$$\frac{1}{2\pi} \int_{-1}^1 \frac{w(x; t_1, t_2 | q)}{\sqrt{1-x^2}} p_m(x; t_1, t_2 | q) p_n(x; t_1, t_2 | q) dx = \frac{t_1^{2n}}{(t_1 t_2; q)_n (q^{n+1}, t_1 t_2; q)_\infty} \delta_{mn},$$

where

$$w(x; t_1, t_2 | q) := \frac{(e^{2i\theta}, e^{-2i\theta}; q)_\infty}{(t_1 e^{i\theta}, t_1 e^{-i\theta}, t_2 e^{i\theta}, t_2 e^{-i\theta}; q)_\infty}, \quad x = \cos \theta.$$

Using Theorem 19.5.1, [8], we obtain

$$S_r \left(\frac{p_n(x; t_1, t_2 | q)}{(t_1 e^{i\theta}, t_1 e^{-i\theta}; q)_\infty} \right) = \frac{p_n(x; r t_1, t_2/r | q)}{(r t_1 e^{i\theta}, r t_1 e^{-i\theta}; q)_\infty},$$

and

$$S_r(u_n)(x) = s_n v_n(x).$$

Let

$$0 < r < 1, \text{ and } \max(|t_1|, |t_2|) < 1.$$

By Lemma 2.1, we get

$$\begin{aligned} &\int_0^\pi \frac{(e^{2i\theta}, e^{-2i\theta}, t_1 r e^{i\theta}, t_1 r e^{-i\theta}; q)_\infty}{(t_2 / r e^{i\theta}, t_2 / r e^{-i\theta}; q)_\infty} |S_r(f)(\cos \theta)|^2 d\theta \\ &\leq \int_0^\pi \frac{(e^{2i\theta}, e^{-2i\theta}, t_1 e^{i\theta}, t_1 e^{-i\theta}; q)_\infty}{(t_2 e^{i\theta}, t_2 e^{-i\theta}; q)_\infty} |f(\cos \theta)|^2 d\theta. \end{aligned}$$

Note that, the equality holds when

$$f(x) = \frac{1}{(t_1 e^{i\theta}, t_1 e^{-i\theta}; q)_\infty}.$$

Acknowledgments

This research is supported by NPST Program of King Saud University, project number 10-math 1293-02.

REFERENCES

- [1] K. F. ANDERSEN AND H. P. HEINIG, *Weighted norm inequalities for certain integral operators*, SIAM J. Math. Anal. **14**, 4 (1983), 834–844.
- [2] J. BAUMEISTER, *Stable Solution of Inverse Problems*, Vieweg und Sohn, Braunschweig/Wiesbaden, 1987.
- [3] A. ERDELYI ET AL., *Higher Transcendental Functions*, vol. 2, McGraw–Hill, New York–Toronto–London, 1953.
- [4] G. GASPER AND M. RAHMAN, *Basic Hypergeometric Series*, second edition, Cambridge University Press, Cambridge, 2004.
- [5] I. C. GOHBERG AND M. G. KREIN, *Introduction to the Theory of Linear Nonselfadjoint Operators*, Amer. Math. Soc., Providence, 1969.
- [6] R. GORENFLO AND VU KIM TUAN, *Singular value decomposition of fractional integration operators in L^2 -spaces with weights*, J. Inv. Ill-Posed Problems **3**, 1 (1995), 1–9.
- [7] G. H. HARDY, J. E. LITTLEWOOD AND G. POLYA, *Inequalities*, Cambridge Univ. Press, Cambridge, 1959.
- [8] M. E. H. ISMAIL, *Classical and Quantum Orthogonal Polynomials in one variable*, Cambridge University Press, Cambridge, 2005.
- [9] A. P. PRUDNIKOV, YU. A. BRYCHKOV AND O. I. MARICHEV, *Integrals and Series, Vol. 2: Special Functions*, Gordon and Breach Science Publishers, New York–London–Paris–Montreux–Tokyo–Melbourne, 1990.
- [10] S. G. SAMKO, A. A. KILBAS AND O. I. MARICHEV, *Fractional Integrals and Derivatives. Theory and Applications*, Gordon and Breach Science Publishers, Switzerland–Australia–Belgium–France–Germany–Great Britain–India–Japan–Malaysia–Netherlands–Russia–Singapore–USA, 1993.
- [11] V. D. STEPANOV, *Two-weighted estimates of Riemann–Liouville integrals*, Math. USSR Izvestiya **36**, 3 (1991), 669–681. Translated from the Russian.
- [12] R. KOEKOEK AND R. SWARTTOUW, *The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogues*, Reports of the Faculty of Technical Mathematics and Informatics no. 98–17, Delft University of Technology, Delft, 1998.

(Received November 5, 2011)

Fethi Bouzaffour
*Department of mathematics, College of Sciences
King Saud University
P. O. Box 2455
Riyadh 11451
Saudi Arabia*
e-mail: fbouzaffour@ksu.edu.sa