

ON k -QUASI- M -HYPONORMAL OPERATORS

SALAH MECHERI

Abstract. In this present article we introduce a new class of operators which we will be called the class of k -quasi- M -hyponormal operators that includes hyponormal and M -hyponormal operators. A part from other results, we show that following results hold for a k -quasi M -hyponormal operator T :

- (i) T has the Bishop's property (β) .
- (ii) The spectral mapping theorem holds for the essential approximate point spectrum of T .
- (iii) Every non-zero isolated point in the spectrum of T is a simple pole of the resolvent of T .

Mathematics subject classification (2010): Primary 47B47, 47A30, 47B20; Secondary 47B10.

Keywords and phrases: k -quasi- M -hyponormal operator, M -hyponormal operator, Bishop's property (β) .

REFERENCES

- [1] S. C. ARORA AND R. KUMAR, *M -hyponormal operators*, Yokoham Math. J **28** (1980), 41–43.
- [2] P. AIENA AND F. VILLAFANE, *Weyl's theorem for some classes of operators*, Integral equations and Operator Theory **53** (2005), 453–466.
- [3] A. DEFANT AND K. FLORET, *Tensor Norms and Operator Ideals*, North-Holland-Amsterdam. Elsevier Science Publishers, 1993.
- [4] T. FURUTA, *Invitation to linear operatorss-From Matrices to bounded Linear Operators in Hilbert space*, Taylor and Francis. London, 2001.
- [5] P. R. HALMOS, *A Hilbert space problem book*, Van Nostrand, Princeton, 1967.
- [6] YOUNG MIN HAN AND JU HEE SON, *On Quasi- M -Hyponormal operators*, Filomat **25** (2011), 37–52.
- [7] IN HO JEAN AND IN HYOUM KIM, *On operators satisfying $T^*|T^2|T \geq T^*|T|^2T$* , Linear Algebra Appl. **418** (2006), 854–862.
- [8] E. KO, *Remark on generalized k -quasihyponormal operators*, Bull. Korean Math. Soc **35** (1998), 701–707.
- [9] S. MECHERI, *Weyl's theorem for algebraically (p,k) -quasihyponormal operators*, Georgian Math. J. **13** (2006), 1998–2007.
- [10] S. MECHERI AND S. MAKHLOUF, *Weyl Type theorems for posinormal operators*, Math. Proc. Royal Irish. Acad. **108(A)** (2008), 68–79.
- [11] M. OUDGHIRI, *Weyl's and Browder's theorem for operators satisfying the SVEP*, Studia Math **163** (2004), 85–101.
- [12] M. PUTINAR, *Hyponormal operators are subscalar*, J. Operator Theory **12** (1984), 385–395.
- [13] M. A. ROSENBLUM, *On the operator equation $BX - XA = Q$* , Duke Math. J. **23** (1956), 263–269.
- [14] J. STAMPFLI, *Hyponormal operators and spectral density*, Trans. Amer. Math. Soc. **117** (1965), 469–476.
- [15] A. UCHIYAMA, Y. TAKASHI, *Weyl's theorem for p -hyponormal or M -hyponormal operators*, Glasg. Math. J. **43** (2001), 375–381.