

A NEW MULTIPLE HALF-DISCRETE HILBERT-TYPE INEQUALITY

BING HE AND BICHENG YANG

(Communicated by I. Perić)

Abstract. By using the way of weight functions and technique of real analysis, a new multiple half-discrete Hilbert-type inequality with the best constant factor is given. As applications, the equivalent forms, operator expressions as well as some reverse inequalities are also considered.

1. Introduction

Suppose that $a_m, b_n \geq 0$, $a = \{a_m\}_{m=1}^{\infty}$, $b = \{b_n\}_{n=1}^{\infty} \in l^2$, $\|a\| = \{\sum_{m=1}^{\infty} a_m^2\}^{\frac{1}{2}} > 0$, $\|b\| > 0$, we have the following well known discrete Hilbert's inequality (cf. [1])

$$\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{a_m b_n}{m+n} < \pi \|a\| \|b\|, \quad (1)$$

where the constant factor π is the best possible. Moreover, for $f(x), g(y) \geq 0$, $f, g \in L^2(\mathbf{R}_+)$, $\|f\| = \{\int_0^{\infty} f^2(x) dx\}^{\frac{1}{2}} > 0$, $\|g\| > 0$, we still have the following Hilbert's integral inequality

$$\int_0^{\infty} \int_0^{\infty} \frac{f(x)g(y)}{x+y} dx dy < \pi \|f\| \|g\|, \quad (2)$$

with the same best constant factor π . Inequalities (1) and (2) are important in analysis and its applications (cf. [2]–[8]). Hilbert's inequality may be classified into several types (discrete, integral, half-discrete etc.). On half-discrete Hilbert-type inequality, Yang [9] gave the following inequality with the best factor π recently:

$$\int_0^{\infty} f(x) \sum_{n=1}^{\infty} \frac{a_n}{x+n} dx < \pi \|f\| \|a\|, \quad (3)$$

It is significant to generalize Hilbert-type inequalities into multiple forms. In recent years, some results have been obtained ([10]–[17]). In 2005, Yang [18] gave a multiple Hilbert-type integral inequality as follows: If $n \in \mathbf{N} \setminus \{1\}$, $p_i, r_i > 1$, $\sum_{i=1}^n (1/p_i) =$

Mathematics subject classification (2010): 26D15, 47A07.

Keywords and phrases: Multiple half-discrete Hilbert-type inequality, weight function, operator expression, equivalent form, reverse inequality.

$\sum_{i=1}^n (1/r_i) = 1$, $\lambda > 0$, $0 < \int_0^\infty x_i^{p_i(1-\lambda/r_i)-1} f_i^{p_i}(x_i) dx_i < \infty$ ($i = 1, \dots, n$), then

$$\begin{aligned} & \int_0^\infty \cdots \int_0^\infty \frac{1}{(\sum_{i=1}^n x_i)^\lambda} \prod_{i=1}^n f_i(x_i) dx_1 \cdots dx_n \\ & < \frac{1}{\Gamma(\lambda)} \prod_{i=1}^n \Gamma\left(\frac{\lambda}{r_i}\right) \left\{ \int_0^\infty x^{p_i(1-\frac{\lambda}{r_i})-1} f_i^{p_i}(x) dx \right\}^{\frac{1}{p_i}}, \end{aligned} \quad (4)$$

where the constant factor $\frac{1}{\Gamma(\lambda)} \prod_{i=1}^n \Gamma\left(\frac{\lambda}{r_i}\right)$ is the best possible.

Until now, we have obtained only few multiple Hilbert-type inequalities. It is not an easy job to find new forms of such inequalities. In this paper, by using the way of weight functions and the technique of real analysis, a new multiple half-discrete Hilbert-type inequality is given. As applications, the equivalent forms, operator expressions as well as some reverse inequalities are also considered.

2. Some Lemmas

LEMMA 1. (cf. [19], (9.1.1)) If $m \in \mathbb{N}$, $p_i \neq 0, 1$, ($i = 1, \dots, m+1$), $\sum_{i=1}^{m+1} \frac{1}{p_i} = 1$, then

$$A := \prod_{i=1}^{m+1} \left[x_i^{p_i-1} \prod_{j=1(j \neq i)}^{m+1} x_j^{-1} \right]^{\frac{1}{p_i}} = 1. \quad (5)$$

Proof. We find

$$\begin{aligned} A &= \prod_{i=1}^{m+1} \left(x_i^{p_i-1+1} \prod_{j=1}^{m+1} x_j^{-1} \right)^{\frac{1}{p_i}} = \prod_{i=1}^{m+1} \left(x_i^{p_i} \right)^{\frac{1}{p_i}} \left(\prod_{j=1}^{m+1} x_j^{-1} \right)^{\frac{1}{p_i}} \\ &= \prod_{i=1}^{m+1} x_i \left(\prod_{j=1}^{m+1} x_j^{-1} \right)^{\sum_{i=1}^{m+1} \frac{1}{p_i}} = \prod_{i=1}^{m+1} x_i \prod_{j=1}^{m+1} x_j^{-1} = 1. \end{aligned}$$

Hence we have (5). \square

LEMMA 2. If $c > 0$, $n \in \mathbb{N}$, then

$$\int_0^c (\ln x)^n dx = c \sum_{i=0}^n (-1)^{n-i} \frac{n!}{i!} (\ln c)^i. \quad (6)$$

Proof. We prove (6) by mathematical induction. For $n = 1$,

$$\int_0^c \ln x dx = (x \ln x - x) \Big|_0^c = c(\ln c - 1).$$

Assume that (6) is valid for $n = k - 1$, i.e.

$$\int_0^c (\ln x)^{k-1} dx = c \sum_{i=0}^{k-1} (-1)^{k-1-i} \frac{(k-1)!}{i!} (\ln c)^i.$$

Then for $n = k$, we have

$$\begin{aligned} \int_0^c (\ln x)^k dx &= x(\ln x)^k|_{0^+}^c - k \int_0^c (\ln x)^{k-1} dx \\ &= c(\ln c)^k - ck \sum_{i=0}^{k-1} (-1)^{k-1-i} \frac{(k-1)!}{i!} (\ln c)^i \\ &= c \sum_{i=0}^k (-1)^{k-i} \frac{k!}{i!} (\ln c)^i. \end{aligned}$$

Hence (6) is valid. \square

LEMMA 3. Suppose that $s, t \in \mathbf{R}_+$, $k \in \mathbf{N}$, $a = \min\{s, 1\}$, $b = \max\{s, 1\}$, then

$$\begin{aligned} &\int_0^\infty \frac{\min\{s, t, 1\}}{\max\{s, t, 1\}} \left(\ln \frac{\min\{s, t, 1\}}{\max\{s, t, 1\}} \right)^k t^{-1} dt \\ &= \frac{a}{b} \left[- \left(\ln \frac{a}{b} \right)^{k+1} + 2 \sum_{i=0}^k (-1)^{k-i} \frac{k!}{i!} \left(\ln \frac{a}{b} \right)^i \right]. \end{aligned} \quad (7)$$

Proof. In view of (6), we have

$$\begin{aligned} &\int_0^\infty \frac{\min\{s, t, 1\}}{\max\{s, t, 1\}} \left(\ln \frac{\min\{s, t, 1\}}{\max\{s, t, 1\}} \right)^k t^{-1} dt \\ &= \int_0^{\min\{s, 1\}} \frac{t}{\max\{s, 1\}} \left(\ln \frac{t}{\max\{s, 1\}} \right)^k t^{-1} dt \\ &\quad + \int_{\min\{s, 1\}}^{\max\{s, 1\}} \frac{\min\{s, 1\}}{\max\{s, 1\}} \left(\ln \frac{\min\{s, 1\}}{\max\{s, 1\}} \right)^k t^{-1} dt \\ &\quad + \int_{\max\{s, 1\}}^\infty \frac{\min\{s, 1\}}{t} \left(\ln \frac{\min\{s, 1\}}{t} \right)^k t^{-1} dt \\ &= \frac{1}{b} \int_0^a \left(\ln \frac{t}{b} \right)^k dt + \frac{a}{b} \left(\ln \frac{a}{b} \right)^k \int_a^b t^{-1} dt + a \int_b^\infty \left(\ln \frac{a}{t} \right)^k t^{-2} dt \\ &= \frac{2}{b} \int_0^a \left(\ln \frac{t}{b} \right)^k dt - \frac{a}{b} \left(\ln \frac{a}{b} \right)^{k+1} \\ &= 2 \int_0^{a/b} (\ln x)^k dx - \frac{a}{b} \left(\ln \frac{a}{b} \right)^{k+1} \\ &= \frac{a}{b} \left[- \left(\ln \frac{a}{b} \right)^{k+1} + 2 \sum_{i=0}^k (-1)^{k-i} \frac{k!}{i!} \left(\ln \frac{a}{b} \right)^i \right]. \end{aligned}$$

Hence (7) is valid. \square

LEMMA 4. For $m, n \in \mathbf{N}$, define the weight functions $\omega_i(x_i)$ ($i = 1, \dots, m$) and

$\omega_{m+1}(n)$ as follows:

$$\begin{aligned}\omega_i(x_i) &:= \sum_{n=1}^{\infty} n^{-1} \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \\ &\quad \times \prod_{j=1(j \neq i)}^m x_j^{-1} dx_1 \cdots dx_{i-1} dx_{i+1} \cdots dx_m,\end{aligned}\tag{8}$$

$$\omega_{m+1}(n) := \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{j=1}^m x_j^{-1} dx_1 \cdots dx_m.\tag{9}$$

Then we have

$$\omega_{m+1}(n) = (m+1)!,\tag{10}$$

$$0 < (m+1)! (1 - \theta_i(x_i)) < \omega_i(x_i) < (m+1)!,\tag{11}$$

$$\begin{aligned}\theta_i(x_i) &:= \frac{1}{(m+1)!} \int_0^{1/x_i} u_{m+1}^{-1} \left[\int_0^{\infty} \cdots \int_0^{\infty} \frac{\min\{u_1, \dots, u_{i-1}, 1, u_{i+1}, \dots, u_{m+1}\}}{\max\{u_1, \dots, u_{i-1}, 1, u_{i+1}, \dots, u_{m+1}\}} \right. \\ &\quad \left. \times \prod_{j=1(j \neq i)}^m x_j^{-1} du_1 \cdots du_{i-1} du_{i+1} \cdots du_m \right] du_{m+1} > 0 \quad (i = 1, \dots, m).\end{aligned}\tag{12}$$

Proof. Setting $x_j = nu_j$ ($j = 1, \dots, m$) in the integral of (9), we find

$$\omega_{m+1}(n) = \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{u_j, 1\}}{\max_{1 \leq j \leq m} \{u_j, 1\}} \prod_{j=1}^m u_j^{-1} du_1 \cdots du_m.$$

We prove (10) by mathematical induction. For $m = 1$,

$$\begin{aligned}\omega_2(n) &= \int_0^{\infty} \frac{\min\{u_1, 1\}}{\max\{u_1, 1\}} u_1^{-1} du_1 \\ &= \int_0^1 du_1 + \int_1^{\infty} u_1^{-2} du_1 = 2 \int_0^1 du_1 = 2!.\end{aligned}$$

Assuming that for $m = k - 1$ ($k \geq 3$), we have

$$\begin{aligned}\omega_k(n) &= \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq k-1} \{u_j, 1\}}{\max_{1 \leq j \leq k-1} \{u_j, 1\}} \prod_{j=1}^{k-1} u_j^{-1} du_1 \cdots du_{k-1} \\ &= \int_0^{\infty} \left(\int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq k-1} \{u_j, 1\}}{\max_{1 \leq j \leq k-1} \{u_j, 1\}} \prod_{j=2}^{k-1} u_j^{-1} du_2 \cdots du_{k-1} \right) u_1^{-1} du_1 \\ &= (k-1)! \int_0^{\infty} \frac{\min\{u_1, 1\}}{\max\{u_1, 1\}} \left[1 + \frac{(-1)^{k-2}}{(k-1)!} \left(\ln \frac{\min\{u_1, 1\}}{\max\{u_1, 1\}} \right)^{k-2} \right. \\ &\quad \left. + \sum_{i=0}^{k-4} \frac{(-1)^{i+1}}{i!} \left(\frac{1}{i+1} - \frac{1}{k-1} \right) \left(\ln \frac{\min\{u_1, 1\}}{\max\{u_1, 1\}} \right)^{i+1} \right] \frac{du_1}{u_1}\end{aligned}$$

$$\begin{aligned}
&= 2(k-1)! \int_0^1 \left[1 + \frac{(-1)^{k-2}}{(k-1)!} (\ln x)^{k-2} + \sum_{i=0}^{k-4} \frac{(-1)^{i+1}}{i!} \left(\frac{1}{i+1} - \frac{1}{k-1} \right) (\ln x)^{i+1} \right] dx \\
&= 2(k-1)! \left[1 + \frac{(k-2)!}{(k-1)!} + \sum_{i=0}^{k-4} \frac{(i+1)!}{i!} \left(\frac{1}{i+1} - \frac{1}{k-1} \right) \right] = k!,
\end{aligned}$$

then for $m = k$, in view of the assumption and (7), we have

$$\begin{aligned}
\omega_{k+1}(n) &= \int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq k} \{u_j, 1\}}{\max_{1 \leq j \leq k} \{u_j, 1\}} \prod_{j=1}^k u_j^{-1} du_1 \cdots du_k \\
&= \int_0^\infty \left(\int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq k} \{u_j, 1\}}{\max_{1 \leq j \leq k} \{u_j, 1\}} \prod_{j=2}^k u_j^{-1} du_2 \cdots du_k \right) u_1^{-1} du_1 \\
&= \int_0^\infty \left[\int_0^\infty \left(\int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq k} \{u_j, 1\}}{\max_{1 \leq j \leq k} \{u_j, 1\}} \prod_{j=3}^k u_j^{-1} du_3 \cdots du_k \right) \frac{du_2}{u_2} \right] \frac{du_1}{u_1} \\
&= \int_0^\infty \left\{ \int_0^\infty (k-1)! \frac{\min\{u_1, u_2, 1\}}{\max\{u_1, u_2, 1\}} \left[1 + \frac{(-1)^{k-2}}{(k-1)!} \left(\ln \frac{\min\{u_1, u_2, 1\}}{\max\{u_1, u_2, 1\}} \right)^{k-2} \right. \right. \\
&\quad \left. \left. + \sum_{i=0}^{k-4} \frac{(-1)^{i+1}}{i!} \left(\frac{1}{i+1} - \frac{1}{k-1} \right) \left(\ln \frac{\min\{u_1, u_2, 1\}}{\max\{u_1, u_2, 1\}} \right)^{i+1} \right] \frac{du_2}{u_2} \right\} \frac{du_1}{u_1} \\
&= (k-1)! \int_0^\infty \frac{\min\{u_1, 1\}}{\max\{u_1, 1\}} \left\{ \left(2 - \ln \frac{\min\{u_1, 1\}}{\max\{u_1, 1\}} \right) + \frac{(-1)^{k-2}}{(k-1)!} \right. \\
&\quad \times \left. \left[-\left(\ln \frac{\min\{u_1, 1\}}{\max\{u_1, 1\}} \right)^{k-1} + 2 \sum_{i=0}^{k-2} (-1)^{k-i} \frac{(k-2)!}{i!} \left(\ln \frac{\min\{u_1, 1\}}{\max\{u_1, 1\}} \right)^i \right] \right. \\
&\quad \left. + \sum_{i=0}^{k-4} \frac{(-1)^{i+1}}{i!} \left(\frac{1}{i+1} - \frac{1}{k-1} \right) \left[-\left(\ln \frac{\min\{u_1, 1\}}{\max\{u_1, 1\}} \right)^{i+2} \right. \right. \\
&\quad \left. \left. + 2 \sum_{j=0}^{i+1} (-1)^{i+1-j} \frac{(i+1)!}{j!} \left(\ln \frac{\min\{u_1, 1\}}{\max\{u_1, 1\}} \right)^j \right] \right\} \frac{du_1}{u_1} \\
&= 2(k-1)! \int_0^1 \left\{ \left(2 - \ln x \right) + \frac{(-1)^{k-2}}{(k-1)!} \left[-(\ln x)^{k-1} \right. \right. \\
&\quad \left. \left. + 2 \sum_{i=0}^{k-2} (-1)^{k-i} \frac{(k-2)!}{i!} (\ln x)^i \right] + \sum_{i=0}^{k-4} \frac{(-1)^{i+1}}{i!} \left(\frac{1}{i+1} - \frac{1}{k-1} \right) \right. \\
&\quad \times \left. \left[-(\ln x)^{i+2} + 2 \sum_{j=0}^{i+1} (-1)^{i+1-j} \frac{(i+1)!}{j!} (\ln x)^j \right] \right\} dx \\
&= 2(k-1)! \left\{ 3 + \frac{(-1)^{k-2}}{(k-1)!} \left[(-1)^k (k-1)! + 2 \sum_{i=0}^{k-2} (-1)^k (k-2)! \right] \right. \\
&\quad \left. + \sum_{i=0}^{k-4} \frac{(-1)^{i+1}}{i!} \left(\frac{1}{i+1} - \frac{1}{k-1} \right) \left[(-1)^{i+1} (i+2)! + 2 \sum_{j=0}^{i+1} (-1)^{i+1} (i+1)! \right] \right\}
\end{aligned}$$

$$\begin{aligned}
&= 2(k-1)! \left[6 + 3 \sum_{i=0}^{k-4} (i+1)(i+2) \left(\frac{1}{i+1} - \frac{1}{k-1} \right) \right] \\
&= (k+1)!.
\end{aligned}$$

and then (10) follows. \square

Since $\frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \cdot n^{-1}$ is decreasing with respect to $n \in \mathbb{N}$, then for any $i = 1, \dots, m$, it follows

$$\begin{aligned}
\omega_i(x_i) &< \int_0^\infty x_{m+1}^{-1} \int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m+1} \{x_j\}}{\max_{1 \leq j \leq m+1} \{x_j\}} dx_{m+1} \\
&\quad \times \prod_{j=1(j \neq i)}^m x_j^{-1} dx_1 \cdots dx_{i-1} dx_{i+1} \cdots dx_m \\
&\stackrel{u_j=x_j/x_i(j \neq i)}{=} \omega_{m+1}(n) = (m+1)!.
\end{aligned}$$

$$\begin{aligned}
\omega_i(x_i) &> \int_1^\infty x_{m+1}^{-1} \left[\int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m+1} \{x_j\}}{\max_{1 \leq j \leq m+1} \{x_j\}} \right. \\
&\quad \times \left. \prod_{j=1(j \neq i)}^m x_j^{-1} dx_1 \cdots dx_{i-1} dx_{i+1} \cdots dx_m \right] dx_{m+1} \\
&\stackrel{u_j=x_j/x_i(j \neq i)}{=} \int_{1/x_i}^\infty u_{m+1}^{-1} \left[\int_0^\infty \cdots \int_0^\infty \frac{\min \{u_1, \dots, u_{i-1}, 1, u_{i+1}, \dots, u_{m+1}\}}{\max \{u_1, \dots, u_{i-1}, 1, u_{i+1}, \dots, u_{m+1}\}} \right. \\
&\quad \times \left. \prod_{j=1(j \neq i)}^m u_j^{-1} du_1 \cdots du_{i-1} du_{i+1} \cdots du_m \right] du_{m+1} \\
&= (k+1)! (1 - \theta_i(x_i)) > 0,
\end{aligned}$$

where $\theta_i(x_i)$ is defined by (12). \square

LEMMA 5. If $m \in \mathbb{N}, p_i \neq 0, 1 (i = 1, \dots, m+1), \frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i} = 1 - \frac{1}{p_{m+1}}$, then for $0 < \varepsilon < |p_{m+1}|$,

$$\begin{aligned}
I(\varepsilon) &:= \varepsilon \sum_{n=1}^\infty n^{-\frac{\varepsilon}{p_{m+1}}-1} \int_1^\infty \cdots \int_1^\infty \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{j=1}^m x_j^{-\frac{\varepsilon}{p_j}-1} dx_1 \cdots dx_m \\
&= (m+1)! + o(1) (\varepsilon \rightarrow 0^+).
\end{aligned} \tag{13}$$

Proof. Since $\frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} n^{-\frac{\varepsilon}{p_{m+1}}-1}$ is strictly decreasing with respect to $n \in \mathbb{N}$, then by the decreasing property and setting $u_j = x_j/x_{m+1}$ ($j = 1, \dots, m$), we find

$$\begin{aligned}
I(\varepsilon) &\geq \varepsilon \int_1^\infty x_{m+1}^{-\frac{\varepsilon}{p_{m+1}}-1} \int_1^\infty \cdots \int_1^\infty \frac{\min_{1 \leq j \leq m+1} \{x_j\}}{\max_{1 \leq j \leq m+1} \{x_j\}} \prod_{j=1}^m x_j^{-\frac{\varepsilon}{p_j}-1} \\
&\quad \times dx_1 \cdots dx_m dx_{m+1} = \varepsilon \int_1^\infty x_{m+1}^{-\varepsilon-1} \left[\int_{\frac{1}{x_{m+1}}}^\infty \cdots \int_{\frac{1}{x_{m+1}}}^\infty \frac{\min_{1 \leq j \leq m} \{u_j, 1\}}{\max_{1 \leq j \leq m} \{u_j, 1\}} \right.
\end{aligned}$$

$$\begin{aligned}
& \times \left[\int_{\frac{1}{x_{m+1}}}^{\infty} \cdots \int_{\frac{1}{x_{m+1}}}^{\infty} \frac{\min_{1 \leq j \leq m} \{u_j, 1\}}{\max_{1 \leq j \leq m} \{u_j, 1\}} \prod_{j=1}^m u_j^{-\frac{\varepsilon}{p_j}-1} du_1 \cdots du_m \right] dx_{m+1} \\
& \geq \varepsilon \int_1^{\infty} x_{m+1}^{-\varepsilon-1} \left[\int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{u_j, 1\}}{\max_{1 \leq j \leq m} \{u_j, 1\}} \prod_{j=1}^m u_j^{-\frac{\varepsilon}{p_j}-1} du_1 \cdots du_m \right] dx_{m+1} \\
& - \varepsilon \sum_{i=1}^m \int_1^{\infty} x_{m+1}^{-1} A_i(x_{m+1}) dx_{m+1} \\
& = (m+1)! + o(1) - \varepsilon \sum_{i=1}^m \int_1^{\infty} x_{m+1}^{-1} A_i(x_{m+1}) dx_{m+1} (\varepsilon \rightarrow 0^+), \\
A_i(x_{m+1}) & := \int_0^{\infty} \cdots \int_0^{\infty} \int_0^{1/x_{m+1}} \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{u_j, 1\}}{\max_{1 \leq j \leq m} \{u_j, 1\}} \\
& \quad \prod_{j=1}^m u_j^{-\frac{\varepsilon}{p_j}-1} du_1 \cdots du_{i-1} du_i du_{i+1} \cdots dx_m.
\end{aligned}$$

Without loss of generality, we estimate $\int_1^{\infty} x_{m+1}^{-1} A_m(x_{m+1}) dx_{m+1}$ as follows:

$$\begin{aligned}
& \int_1^{\infty} x_{m+1}^{-1} A_m(x_{m+1}) dx_{m+1} \\
& = \int_1^{\infty} x_{m+1}^{-1} \left[\int_0^{1/x_{m+1}} \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{u_j, 1\}}{\max_{1 \leq j \leq m} \{u_j, 1\}} \right. \\
& \quad \left. \times \prod_{j=1}^m u_j^{-\frac{\varepsilon}{p_j}-1} du_1 \cdots du_{m-1} du_m \right] dx_{m+1} \\
& = \int_0^1 \left(\int_1^{1/u_m} x_{m+1}^{-1} dx_{m+1} \right) \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{u_j, 1\}}{\max_{1 \leq j \leq m} \{u_j, 1\}} \\
& \quad \times \prod_{j=1}^m u_j^{-\frac{\varepsilon}{p_j}-1} du_1 \cdots du_{m-1} du_m \\
& = \int_0^1 (-\ln u_m) \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{u_j, 1\}}{\max_{1 \leq j \leq m} \{u_j, 1\}} \\
& \quad \times \prod_{j=1}^m u_j^{-\frac{\varepsilon}{p_j}-1} du_1 \cdots du_{m-1} du_m.
\end{aligned}$$

Setting $\alpha > 0$ small enough, since $\lim_{u_m \rightarrow 0^+} u_m^\alpha (-\ln u_m) = 0$, there exists a constant $M_m > 0$, and $0 < u_m^\alpha (-\ln u_m) < M_m, u_m \in (0, 1]$. We obtain

$$\begin{aligned}
0 & < \int_1^{\infty} x_{m+1}^{-1} A_m(x_{m+1}) dx_{m+1} \\
& \leq M_m \int_0^{\infty} \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{u_j, 1\}}{\max_{1 \leq j \leq m} \{u_j, 1\}} \\
& \quad \times \prod_{j=1}^{m-1} u_j^{-\frac{\varepsilon}{p_j}-1} u_m^{-\frac{\alpha p_m + \varepsilon}{p_m}-1} du_1 \cdots du_{m-1} du_m \\
& \rightarrow M_m O_m(\alpha) < \infty (\varepsilon \rightarrow 0^+),
\end{aligned}$$

namely, $\sum_{i=1}^{\infty} \int_1^{\infty} x_{m+1}^{-1} A_i(x_{m+1}) dx_{m+1} = O(1)$. Hence

$$I(\varepsilon) \geq (m+1)! + o(1) - \varepsilon O(1) (\varepsilon \rightarrow 0^+).$$

We still have

$$\begin{aligned} I(\varepsilon) &\leq \varepsilon \sum_{n=1}^{\infty} n^{-\varepsilon-1} \left[n^{\varepsilon - \frac{\varepsilon}{p_{m+1}}} \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{j=1}^m x_j^{-\frac{\varepsilon}{p_j}-1} dx_1 \cdots dx_m \right] \\ &= \varepsilon \left(1 + \sum_{n=2}^{\infty} n^{-\varepsilon-1} \right) [(m+1)! + o(1)] \\ &\leq \varepsilon \left(1 + \int_1^{\infty} y^{-\varepsilon-1} dy \right) [(m+1)! + o(1)] \\ &= (\varepsilon + 1) [(m+1)! + o(1)] (\varepsilon \rightarrow 0^+). \end{aligned}$$

Hence (13) is valid. \square

LEMMA 6. If $m \in \mathbb{N}$, $p_i \neq 0, 1$ ($i = 1, \dots, m+1$), $\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i} = 1 - \frac{1}{p_{m+1}}$, $f_i(x_i)$ ($i = 1, \dots, m$) are non-negative measurable functions in \mathbf{R}_+ , then (i) for $p_i > 1$ ($i = 1, \dots, m+1$), we have the following inequality

$$\begin{aligned} J &:= \left\{ \sum_{n=1}^{\infty} n^{-1} \left(\int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \right)^p \right\}^{1/p} \\ &\leq [(m+1)!]^{\frac{1}{p_{m+1}}} \prod_{i=1}^m \left\{ \int_0^{\infty} \omega_i(x_i) x_i^{p_i-1} f_i(x_i) dx_i \right\}^{1/p_i}; \end{aligned} \quad (14)$$

(ii) for $0 < p_1 < 1$, $p_i < 0$ ($i = 2, \dots, m+1$), or $p_i < 0$ ($i = 1, \dots, m$), $0 < p_{m+1} < 1$, we have the reverse of (14).

Proof. For $p_i > 1$ ($i = 1, \dots, m+1$) ($p > 1$), by Hölder's inequality with weight (cf. [21]) and (6), we have

$$\begin{aligned} &\left(\int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \right)^p \\ &= \left\{ \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m \left[x_i^{p_i-1} n^{-1} \prod_{j=1(j \neq i)}^m x_j^{-1} \right]^{1/p_i} f_i(x_i) \right. \\ &\quad \times \left. \left[n^{p_{m+1}-1} \prod_{j=1}^m x_j^{-1} \right]^{1/p_{m+1}} dx_1 \cdots dx_m \right\}^p \\ &\leq \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m \left[x_i^{p_i-1} n^{-1} \prod_{j=1(j \neq i)}^m x_j^{-1} \right]^{\frac{p}{p_i}} f_i^p(x_i) dx_1 \cdots dx_m \\ &\quad \times \left\{ n^{p_{m+1}-1} \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{j=1}^m x_j^{-1} dx_1 \cdots dx_m \right\}^{\frac{p}{p_{m+1}}} \end{aligned}$$

$$= \frac{[\omega_{m+1}(n)]^{\frac{p}{p_{m+1}}}}{n^{-1}} \int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} n^{-1} \\ \times \prod_{i=1}^m \left[x_i^{p_i-1} \prod_{j=1(j \neq i)}^m x_j^{-1} \right]^{\frac{p}{p_i}} f_i^p(x_i) dx_1 \cdots dx_m.$$

Since $\omega_{m+1}(n) = (m+1)!$, we obtain

$$J \leq [(m+1)!]^{\frac{1}{p_{m+1}}} \left\{ \sum_{n=1}^\infty \int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} n^{-1} \right. \\ \left. \times \prod_{i=1}^m \left[x_i^{p_i-1} \prod_{j=1(j \neq i)}^m x_j^{-1} \right]^{\frac{p}{p_i}} f_i^p(x_i) dx_1 \cdots dx_m \right\}^{\frac{1}{p}} \\ = [(m+1)!]^{\frac{1}{p_{m+1}}} \left\{ \int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \sum_{n=1}^\infty n^{-1} \right. \\ \left. \times \prod_{i=1}^m \left[x_i^{p_i-1} \prod_{j=1(j \neq i)}^m x_j^{-1} \right]^{\frac{p}{p_i}} f_i^p(x_i) dx_1 \cdots dx_m \right\}^{\frac{1}{p}}.$$

Since $\sum_{n=1}^\infty \frac{1}{p_i/p} = 1$, applying Hölder's inequality, it follows

$$J \leq [(m+1)!]^{\frac{1}{p_{m+1}}} \prod_{i=1}^m \left\{ \int_0^\infty \left[\sum_{n=1}^\infty n^{-1} \int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \right] \right. \\ \left. \times \prod_{j=1(j \neq i)}^m x_j^{-1} dx_1 \cdots dx_{i-1} dx_{i+1} \cdots dx_m \right] x_i^{p_i-1} f_i^p(x_i) dx_i \right\}^{\frac{1}{p_i}} \\ = [(m+1)!]^{\frac{1}{p_{m+1}}} \prod_{i=1}^m \left\{ \int_0^\infty \omega_i(x_i) x_i^{p_i-1} f_i^p(x_i) dx_i \right\}^{\frac{1}{p_i}},$$

and (14) follows.

(ii) For $0 < p_1 < 1$, $p_i < 0$ ($i = 2, \dots, m+1$), or $p_i < 0$ ($i = 1, \dots, m$), $0 < p_{m+1} < 1$, in view of the assumptions and by the same way, we obtain the reverse of (14). \square

3. Main results and applications

In this section, for $p_i > 1$ ($i = 1, \dots, m+1$), we define the following functions and spaces:

$$\varphi_i(x_i) := x_i^{p_i-1} (x_i \in \mathbf{R}_+), \quad \psi(n) := n^{p_{m+1}-1} (n \in \mathbf{N}),$$

$$L_{p, \varphi_i}(\mathbf{R}_+) := \left\{ f; \|f\|_{p, \varphi_i} = \left\{ \int_0^\infty x_i^{p_i-1} |f(x_i)|^{p_i} dx \right\}^{\frac{1}{p_i}} < \infty \right\}, \quad (x_i \in \mathbf{R}_+, i = 1, \dots, m),$$

$$l_{p_{m+1}, \psi} := \left\{ a = \{a_n\}_{n=1}^{\infty}; \|a\|_{p_{m+1}, \psi} = \left\{ \sum_{n=1}^{\infty} n^{p_{m+1}-1} |a_n|^{p_{m+1}} \right\}^{\frac{1}{p_{m+1}}} < \infty \right\}.$$

THEOREM 1. If $m \in \mathbf{N}$, $p_i > 1$ ($i = 1, \dots, m+1$), $\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i} = 1 - \frac{1}{p_{m+1}}$, $f_i(x_i)$, $a_n \geq 0$, $f_i \in L_{p_i, \varphi_i}(\mathbf{R}_+)$ ($i = 1, \dots, m$), $a = \{a_n\}_{n=1}^{\infty} \in l_{p_{m+1}, \psi}$, $\|f_i\|_{p_i, \varphi_i}$, $\|a\|_{p_{m+1}, \psi} > 0$, then we have the following equivalent inequalities

$$\begin{aligned} I &:= \sum_{n=1}^{\infty} a_n \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \\ &< (m+1)! \|a\|_{p_{m+1}, \psi} \prod_{i=1}^m \|f_i\|_{p_i, \varphi_i}, \end{aligned} \quad (15)$$

$$\begin{aligned} J &= \left\{ \sum_{n=1}^{\infty} n^{-1} \left(\int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \right)^p \right\}^{1/p} \\ &< (m+1)! \prod_{i=1}^m \|f_i\|_{p_i, \varphi_i}, \end{aligned} \quad (16)$$

where the constant factor $(m+1)!$ in the above inequalities are the best possible.

Proof. By (14), for $\omega_i(x_i) < (m+1)!$ and the assumptions of Theorem, we have (16).

By Hölder's inequality, we obtain

$$\begin{aligned} I &= \sum_{n=1}^{\infty} \left[(\psi(n))^{\frac{1}{p_{m+1}}} \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \right. \\ &\quad \times \left. \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \right] \left[(\psi(n))^{\frac{1}{p_{m+1}}} a_n \right] \leq J \|a\|_{p_{m+1}, \psi}. \end{aligned} \quad (17)$$

Then by (16), we have (15). Assuming (15) is valid, setting

$$a_n := n^{-1} \left(\int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \right)^{p-1}, \quad n \in \mathbf{N},$$

then we find $J^{p-1} = \|a\|_{p_{m+1}, \psi}$. By (14), we have $J < \infty$. If $J = 0$, then (16) is valid trivially; if $J > 0$, then by (15), it follows

$$\begin{aligned} \|a\|_{p_{m+1}, \psi}^{p_{m+1}} &= J^p = I < (m+1)! \|a\|_{p_{m+1}, \psi} \prod_{i=1}^m \|f_i\|_{p_i, \varphi_i}, \\ \|a\|_{p_{m+1}, \psi}^{p_{m+1}-1} &= J < (m+1)! \prod_{i=1}^m \|f_i\|_{p_i, \varphi_i}, \end{aligned}$$

and (16) follows. Hence (15) and (16) are equivalent.

It remains to prove that the constants appearing in inequalities (15) and (16) are the best possible. For $\varepsilon > 0$, define $\tilde{f}_i(x_i)$ and \tilde{a}_n as follows:

$$\tilde{f}_i(x_i) := \begin{cases} 0, & 0 < x_i < 1, \\ x_i^{-\frac{\varepsilon}{p_i}-1}, & x_i \geq 1 \end{cases} \quad (i = 1, \dots, m),$$

$\tilde{a}_n := n^{-\frac{\varepsilon}{pm+1}-1}$, $n \in \mathbb{N}$. If there exists a constant $K \leq (m+1)!$, such that (15) is still valid as we replace $(m+1)!$ by K , then by (13), we have

$$\begin{aligned} (m+1)! + o(1) &= I(\varepsilon) \\ &= \varepsilon \sum_{n=1}^{\infty} \tilde{a}_n \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m \tilde{f}_i(x_i) dx_1 \cdots dx_m \\ &< \varepsilon K \|\tilde{a}\|_{p_{m+1}, \psi} \prod_{i=1}^m \|\tilde{f}_i\|_{p_i, \varphi_i} \\ &= \varepsilon K \left(1 + \sum_{n=2}^{\infty} \frac{1}{n^{1+\varepsilon}} \right)^{\frac{1}{pm+1}} \prod_{i=1}^m \left(\frac{1}{\varepsilon} \right)^{\frac{1}{p_i}} \\ &< \varepsilon K \left(1 + \int_1^{\infty} \frac{dy}{y^{1+\varepsilon}} \right)^{\frac{1}{pm+1}} \prod_{i=1}^m \left(\frac{1}{\varepsilon} \right)^{\frac{1}{p_i}} = K(1+\varepsilon)^{\frac{1}{pm+1}}, \end{aligned}$$

and then $(m+1)! \leq K(\varepsilon \rightarrow 0^+)$. Hence $(m+1)! = K$ is the best constant factor in (15).

By the equivalence, the constant factor $(m+1)!$ in (16) is still the best possible. Otherwise, we can come to a contradiction by (17) that the constant in (15) is not the best possible. \square

REMARK 1. With the assumptions of Theorem 1, we define a half-discrete Hilbert-type operator $T : \prod_{i=1}^m L_{p_i, \varphi_i}(\mathbf{R}_+) \rightarrow l_{p, \psi^{1-p}}$ as follows:

For any $f = (f_1, \dots, f_m) \in \prod_{i=1}^m L_{p_i, \varphi_i}(\mathbf{R}_+)$, we define a Tf , satisfying for any $n \in \mathbb{N}$,

$$(Tf)(n) = \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m. \quad (18)$$

Then by (16), it follows

$$\|Tf\|_{p, \psi^{1-p}} \leq (m+1)! \prod_{i=1}^m \|f_i\|_{p_i, \varphi_i},$$

and consequently $Tf \in l_{p, \psi^{1-p}}$. Hence T is a bounded linear operator with $\|T\| \leq (m+1)!$. Since the constant factor in (16) is the best possible, we have

$$\|T\| := \sup_{f(\neq 0) \in \prod_{i=1}^m L_{p_i, \varphi_i}(\mathbf{R}_+)} \frac{\|Tf\|_{p, \psi^{1-p}}}{\prod_{i=1}^m \|f_i\|_{p_i, \varphi_i}} = (m+1)!. \quad (19)$$

In order to obtain reverse inequalities that correspond to those in Theorem 1, we define the function $\Phi_1(x_1) := (1 - \theta_1(x_1))\varphi_1(x_1)$. For $p_i < 1$ ($p_i \neq 0$), the spaces $L_{p_i, \varphi_i}(\mathbf{R}_+)$ and $l_{p_{m+1}, \psi}$ are not normal spaces, but we still use them as the formal symbols in the following.

THEOREM 2. Suppose that $m \in \mathbf{N}$, $0 < p_1 < 1$, $p_i < 0$ ($i = 2, \dots, m+1$), $\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i} = 1 - \frac{1}{p_{m+1}}$, $f_i(x_i)$, $a_n \geq 0$, $f_i \in L_{p_i, \varphi_i}(\mathbf{R}_+)$ ($i = 1, \dots, m$), $a = \{a_n\}_{n=1}^\infty \in l_{p_{m+1}, \psi}$, $\|f_i\|_{p_i, \varphi_i}$, $\|a\|_{p_{m+1}, \psi} > 0$, then we have the following equivalent inequalities

$$\begin{aligned} I &= \sum_{n=1}^\infty a_n \int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \\ &> (m+1)! \|a\|_{p_{m+1}, \psi} \|f_1\|_{p_1, \Phi_1} \prod_{i=2}^m \|f_i\|_{p_i, \varphi_i}, \end{aligned} \quad (20)$$

$$\begin{aligned} J &= \left\{ \sum_{n=1}^\infty n^{-1} \left(\int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \right)^p \right\}^{1/p} \\ &> (m+1)! \|f_1\|_{p_1, \Phi_1} \prod_{i=2}^m \|f_i\|_{p_i, \varphi_i}, \end{aligned} \quad (21)$$

where the constant factor $(m+1)!$ in the above inequalities is the best possible.

Proof. By the reverse of (14), for $\omega_1(x_1) > (m+1)![1 - \theta_1(x_1)]$, $\omega_i(x_i) < (m+1)!$ and the assumptions of Theorem, we have (21). By Hölder's inequality, we obtain

$$\begin{aligned} I &= \sum_{n=1}^\infty \left[(\psi(n))^{\frac{1}{p_{m+1}}} \int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \right. \\ &\quad \left. \times \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \right] \left[(\psi(n))^{\frac{1}{p_{m+1}}} a_n \right] \geq J \|a\|_{p_{m+1}, \psi}. \end{aligned} \quad (22)$$

Then by (21), we have (20). Assuming that (20) is valid, setting

$$a_n := n^{-1} \left(\int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \right)^{p-1}, \quad n \in \mathbf{N},$$

we find that $J^{p-1} = \|a\|_{p_{m+1}, \psi}$. By the reverse of (14), we have $J > 0$. If $J = \infty$, then (21) is valid trivially; if $J < \infty$, then by (20), it follows

$$\begin{aligned} \|a\|_{p_{m+1}, \psi}^{p_{m+1}} &= J^p = I > (m+1)! \|a\|_{p_{m+1}, \psi} \|f_1\|_{p_1, \Phi_1} \prod_{i=2}^m \|f_i\|_{p_i, \varphi_i}, \\ \|a\|_{p_{m+1}, \psi}^{p_{m+1}-1} &= J > (m+1)! \|f_1\|_{p_1, \Phi_1} \prod_{i=2}^m \|f_i\|_{p_i, \varphi_i}, \end{aligned}$$

namely, (21) follows. Hence (20) and (21) are equivalent.

Now we prove that inequalities (20) and (21) include the best constants. For $0 < \varepsilon < |p_{m+1}|$, we set $\tilde{f}_i(x_i)$ and \tilde{a}_n as Theorem 1. If there exists a constant $K \geq (m+1)!$, such that (20) is still valid as we replace $(m+1)!$ by K , then by (13), we have

$$\begin{aligned} (m+1)! + o(1) &= I(\varepsilon) \\ &= \varepsilon \sum_{n=1}^\infty \tilde{a}_n \int_0^\infty \cdots \int_0^\infty \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m \tilde{f}_i(x_i) dx_1 \cdots dx_m \\ &> \varepsilon K \|\tilde{a}\|_{p_{m+1}, \psi} \|\tilde{f}_1\|_{p_1, \Phi_1} \prod_{i=2}^m \|\tilde{f}_i\|_{p_i, \varphi_i}. \end{aligned}$$

For $0 < \alpha < 1$, define

$$F(u_2, \dots, u_{m+1}) := u_{m+1}^{-\alpha} \frac{\min\{1, u_2, \dots, u_{m+1}\}}{\max\{1, u_2, \dots, u_{m+1}\}} \frac{\min\{1, u_2, \dots, u_m\}}{\max\{1, u_2, \dots, u_m\}}$$

$$(u = (u_2, \dots, u_{m+1}) \in \mathbf{R}_+^m).$$

Since we find

$$\lim_{u \rightarrow 0^+} F(u_2, \dots, u_{m+1}) = \lim_{u \rightarrow \infty} F(u_2, \dots, u_{m+1}) = 0,$$

there exists a constant $M_m > 0$, such that for any $(u_2, \dots, u_{m+1}) \in \mathbf{R}_+^m$, $F(u_2, \dots, u_{m+1}) \leq M_m$, namely,

$$\frac{\min\{1, u_2, \dots, u_{m+1}\}}{\max\{1, u_2, \dots, u_{m+1}\}} \leq M_m u_{m+1}^\alpha \frac{\min\{1, u_2, \dots, u_m\}}{\max\{1, u_2, \dots, u_m\}}.$$

Hence, in view of (12), it follows that

$$\begin{aligned} \theta_1(x_1) &= \frac{1}{(m+1)!} \int_0^{1/x_1} u_{m+1}^{-1} \left[\int_0^\infty \cdots \int_0^\infty \frac{\min\{1, u_2, \dots, u_{m+1}\}}{\max\{1, u_2, \dots, u_{m+1}\}} \right. \\ &\quad \times \left. \prod_{j=2}^m x_j^{-1} du_2 \cdots, du_m \right] du_{m+1} \\ &\leq \frac{M_m}{(m+1)!} \int_0^{1/x_1} u_{m+1}^{\alpha-1} \left[\int_0^\infty \cdots \int_0^\infty \frac{\min\{1, u_2, \dots, u_m\}}{\max\{1, u_2, \dots, u_m\}} \right. \\ &\quad \times \left. \prod_{j=2}^m x_j^{-1} du_2 \cdots, du_m \right] du_{m+1} \\ &= \frac{M_m}{m+1} \int_0^{1/x_1} u_{m+1}^{\alpha-1} du_{m+1} = O\left(\frac{1}{x_1^\alpha}\right). \end{aligned}$$

Hence we find

$$\begin{aligned} \varepsilon K \|\tilde{a}\|_{p_{m+1}, \Psi} \|\tilde{f}_1\|_{p_1, \Phi_1} \prod_{i=2}^m \|\tilde{f}_i\|_{p_i, \varphi_i} \\ = \varepsilon K \left(\sum_{n=1}^{\infty} \frac{1}{n^{1+\varepsilon}} \right)^{\frac{1}{p_{m+1}}} \left[\int_1^\infty \left(1 - O\left(\frac{1}{x_1^\alpha}\right)\right) \frac{dx_1}{x_1^{1+\varepsilon}} \right]^{\frac{1}{p_1}} \prod_{i=2}^m \left(\frac{1}{\varepsilon}\right)^{\frac{1}{p_i}} \\ > \varepsilon K \left(1 + \int_1^\infty \frac{dy}{y^{1+\varepsilon}} \right)^{\frac{1}{p_{m+1}}} \left[\frac{1}{\varepsilon} - O(1) \right]^{\frac{1}{p_1}} \prod_{i=2}^m \left(\frac{1}{\varepsilon}\right)^{\frac{1}{p_i}} \\ = K(1+\varepsilon)^{\frac{1}{p_{m+1}}} [1 - \varepsilon O(1)]^{\frac{1}{p_1}}, \end{aligned}$$

and then $(m+1)! \geq K(\varepsilon \rightarrow 0^+)$. Hence $(m+1)! = K$ is the best constant factor in (20).

By the equivalence, the constant factor $(m+1)!$ in (21) is still the best possible. Otherwise, we can come to a contradiction by (22) that the constant in (20) is not the best possible. \square

In the same way, we still have

THEOREM 3. Assuming that $m \in \mathbf{N}$, $p_i < 0$ ($i = 1, \dots, m$), $0 < p_{m+1} < 1$, $\frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i} = 1 - \frac{1}{p_{m+1}}$, $f_i(x_i), a_n \geq 0$, $f_i \in L_{p_i, \varphi_i}(\mathbf{R}_+)$ ($i = 1, \dots, m$), $a = \{a_n\}_{n=1}^\infty \in l_{p_{m+1}, \psi}$, $\|f_i\|_{p_i, \varphi_i}, \|a\|_{p_{m+1}, \psi} > 0$, then we have the following equivalent inequalities

$$\begin{aligned} & \sum_{n=1}^{\infty} a_n \int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \\ & > (m+1)! \|a\|_{p_{m+1}, \psi} \prod_{i=1}^m \|f_i\|_{p_i, \varphi_i}, \end{aligned} \quad (23)$$

$$\left\{ \sum_{n=1}^{\infty} n^{-1} \left(\int_0^{\infty} \cdots \int_0^{\infty} \frac{\min_{1 \leq j \leq m} \{x_j, n\}}{\max_{1 \leq j \leq m} \{x_j, n\}} \prod_{i=1}^m f_i(x_i) dx_1 \cdots dx_m \right)^p \right\}^{1/p} \\ > (m+1)! \prod_{i=1}^m \|f_i\|_{p_i, \varphi_i}, \quad (24)$$

where the constant factor $(m+1)!$ in the above inequalities is the best possible.

Acknowledgement.

This research was supported by the National Natural Science Foundation of China, under Research Grant 11271085 (first author), the Natural Science Foundation of Guangdong Province, China, under Research Grant S20120400068 65 (first author), and the 2012 Knowledge Construction Special Foundation Item of Guangdong Institution of Higher Learning College and University, under Research Grant 2012KJCX0079 (second author).

REFERENCES

- [1] G. H. HARDY, J. E. LITTLEWOOD, AND G. PÓLYA, *Inequalities*, Cambridge University Press, Cambridge, 1934.
- [2] D. S. MITRINOVIĆ, J. E. PEČARIĆ, AND A. M. FINK, *Inequalities involving functions and their integrals and derivatives*, Kluwer Academic Publishers, Boston, 1991.
- [3] B. YANG, *A Hilbert-type integral inequality with a non-homogeneous kernel*, Journal of Xiamen University (Natural Science), **48**(2)(2009), 165–169.
- [4] B. YANG, *On the norm of an integral operator and applications*, J. Math. Anal. Appl., **321** (2006), 182–192.
- [5] B. YANG, *On the norm of a certain self-adjoint integral operator and applications to bilinear integral inequalities*, Taiwanese Journal of Mathematics, **12** (2) (2008), 315–324.
- [6] B. HE, *On a Hilbert-type integral inequality with a homogeneous kernel in R^2 and its equivalent form*, Journal of Inequalities and Applications, 2012, **94** (2012) doi:10.1186/1029-242X-2012-94.
- [7] B. HE AND Y. LI, *On several new inequalities close to Hilbert-Pachpatte's inequality*, J. Ineq. in Pure and Applied Math., **7** (4) (2006), Art.154: 1–9.
- [8] B. HE AND B. YANG, *On a half-discrete inequality with a general homogeneous kernel*, Journal of Inequalities and Applications 2012, **30** (2012) doi:10.1186/1029-242X-2012-30.
- [9] B. YANG, *A survey of the study of Hilbert-type inequalities with parameters*, Advances in Math., 2009, **38** (3): 257–268.
- [10] Q. HUANG AND B. YANG, *On a multiple Hilbert-type integral operator and applications*, Journal of Inequalities and Applications, Vol. 2009, Art. ID 192197: 1–13.
- [11] Q. HUANG AND B. YANG, *A multiple Hilbert-type integral inequality with a non-homogeneous kernel*, Journal of Inequalities and Applications, 2013, **73** (2013) doi:10.1186/1029-242X-2013-73.

- [12] Y. HONG, *On multiple Hardy-Hilbert integral inequalities with some parameters*, Journal of Inequalities and Applications, Vol. 2006, Art. ID 94960: 1–11.
- [13] I. BRNETIĆ, M. KRNIĆ, AND J. PEČARIĆ, *Multiple Hilbert and Hardy-Hilbert inequalities with non-conjugate parameters*, Bull. Austral. Math. Soc. **71** (2005), 447–457.
- [14] M. KRNIĆ, *Multidimensional Hilbert-type inequality on the weighted Orlicz spaces*, Mediterranean J. Math. **9** (2012), 883–895.
- [15] I. PERIĆ AND P. VUKOVIĆ, *Hardy-Hilbert's inequalities with a general homogeneous kernel*, Math. Inequal. Appl. **12** (2009), 525–536.
- [16] I. PERIĆ AND P. VUKOVIĆ, *Hilbert-type inequalities with a product-type homogeneous kernel and Schur polynomials*, J. Math. Anal. Appl. **359** (2009), 786–793.
- [17] W. ZHONG AND B. YANG, *On a multiple Hilbert-type integral inequality with the symmetric kernel*, Journal of Inequalities and Applications, Vol. 2007, Art. ID 27962: 1–17.
- [18] B. YANG, I. BRNETIĆ, M. KRNIĆ, AND J. PEČARIĆ, *Generalization of Hilbert and Hardy-Hilbert integral inequalities*, Math. Inequal. Appl., **8**, 2 (2005): 259–272.
- [19] B. YANG, *The norm of operator and Hilbert-type inequalities*, Science Press, Beijin, 2009.
- [20] J. KUANG, *Introduction to real analysis*, Hunan Education Press, Changsha, 1996.
- [21] J. KUANG, *Applied inequalities*, Shangdong Science Technic Press, Jinan, 2004.

(Received March 7, 2013)

Bing He

Dept. of Math., Guangdong University of Education
Guangzhou, Guangdong 510303, P. R. China
e-mail: hzs314@163.com

Bicheng Yang

Dept. of Math., Guangdong University of Education
Guangzhou, Guangdong 510303, P. R. China
e-mail: bcyang@gdei.edu.cn