

ON THE STABILITY OF AN ALTERNATIVE QUADRATIC FUNCTIONAL EQUATION

PAISAN NAKMAHACHALASINT

Abstract. We will investigate the stability of the alternative quadratic functional equation

$$f(x+y) + f(x-y) + 2f(x) + 2f(y) \neq 0 \Rightarrow f(x+y) + f(x-y) = 2f(x) + 2f(y).$$

Mathematics subject classification (2010): 39B52, 39B82.

Keywords and phrases: Alternative functional equation; Quadratic functional equation; Stability of functional equation.

REFERENCES

- [1] B. BATKO, *On the stability of an alternative functional equation*, Math. Inequal. Appl. **8** (2005), 685–691.
- [2] B. BATKO AND J. TABOR, *Stability of an alternative Cauchy equation on a restricted domain*, Aequationes Math. **57** (1999), 221–232.
- [3] B. BATKO AND J. TABOR, *Stability of the generalized alternative Cauchy equation*, Abh. Math. Sem. Univ. Hamb. **69** (1999), 67–73.
- [4] S. CZERWIK, *Functional Equations and Inequalities in Several Variables*, World Scientific, Singapore, 2002
- [5] D. H. HYERS, *On the Stability of the Linear Functional Equations*, Proc. Natl. Acad. Sci. **27** (1941), 222–224.
- [6] J. M. RASSIAS, *On the General Quadratic Functional Equation*, Bol. Soc. Mat. Mexicana **11** (2005), 259–268.
- [7] TH. M. RASSIAS, *On the Stability of the Linear Mapping in Banach Spaces*, Proc. Amer. Math. Soc. **72** (1978), 297–300.
- [8] S. M. ULAM, *Problems in Modern Mathematics*, Wiley-Interscience, New York, 1964.