

SHARP ESTIMATES REGARDING THE REMAINDER OF THE ALTERNATING HARMONIC SERIES

ALINA SÎNTĂMĂRIAN

Abstract. In the present paper we obtain enhanced estimates regarding the remainder of the alternating harmonic series. More precisely, we show that

$$\frac{1}{4n^2+a} < \left| \sum_{k=1}^n (-1)^{k-1} \frac{1}{k} - (-1)^{n-1} \frac{1}{2n} - \ln 2 \right| \leqslant \frac{1}{4n^2+b},$$

for all $n \in \mathbb{N}$, with $a = 2$ and $b = \frac{2(3-4\ln 2)}{2\ln 2-1} = 1.177398899\dots$. In addition, the constants a and b are the best possible with the above-mentioned property.

Mathematics subject classification (2010): 11Y60, 40A05, 41A44, 33B15.

Keywords and phrases: Alternating harmonic series, harmonic number, digamma function, Euler–Mascheroni constant, best constant.

REFERENCES

- [1] H. ALZER, *Inequalities for the gamma and polygamma functions*, Abh. Math. Sem. Univ. Hamburg **68** (1998), 363–372.
- [2] C.-P. CHEN, *Inequalities for the Euler–Mascheroni constant*, Appl. Math. Lett. **23**, 2 (2010), 161–164.
- [3] C.-P. CHEN AND F. QI, *The best lower and upper bounds of harmonic sequence*, RGMIA **6**, 2 (2003), 303–308.
- [4] C.-P. CHEN AND C. MORTICI, *New sequence converging towards the Euler–Mascheroni constant*, Comput. Math. Appl. **64**, 4 (2012), 391–398.
- [5] C.-P. CHEN AND H. M. SRIVASTAVA, *New representations for the Lugo and Euler–Mascheroni constants*, Appl. Math. Lett. **24**, 7 (2011), 1239–1244.
- [6] O. FURDUI, *Limits, Series, and Fractional Part Integrals. Problems in Mathematical Analysis*, Springer, New York (2013).
- [7] I. S. GRADSHTEYN AND I. M. RYZHIK, *Table of Integrals, Series, and Products* (7th ed.), Elsevier/Academic Press, Amsterdam (2007).
- [8] B.-N. GUO AND F. QI, *Sharp bounds for harmonic numbers*, Appl. Math. Comput. **218**, 3 (2011), 991–995.
- [9] D. K. KAZARINOFF, *A simple derivation of the Leibniz–Gregory series for $\pi/4$* , Amer. Math. Monthly **62**, 10 (1955), 726–727.
- [10] F. W. J. OLVER (ED.), D. W. LOZIER (ED.), R. F. BOISVERT (ED.) AND C. W. CLARK (ED.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, Cambridge, 2010.
- [11] W. RAUTENBERG, *Zur Approximation von e durch $(1 + \frac{1}{n})^n$ (On the approximation of e by $(1 + \frac{1}{n})^n$)*, Math. Semesterber. **33** (1986), 227–236.
- [12] A. SÎNTĂMĂRIAN, *A new proof for estimating the remainder of the alternating harmonic series*, Creat. Math. Inform. **21**, 2 (2012), 221–225.
- [13] L. TÓTH, *Problem E3432*, Amer. Math. Monthly **98**, 3 (1991), 264.
- [14] L. TÓTH, *Problem E3432 (Solution)*, Amer. Math. Monthly **99**, 7 (1992), 684–685.
- [15] L. TÓTH, *On a class of Leibniz series*, Rev. Anal. Numér. Théor. Approx. **21**, 2 (1992), 195–199.
- [16] L. TÓTH AND J. BUKOR, *On the alternating series $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$* , J. Math. Anal. Appl. **282**, 1 (2003), 21–25.