

HARDY AND RELLICH TYPE INEQUALITIES WITH TWO WEIGHT FUNCTIONS

SEMRA AHMETOLAN AND ISMAIL KOMBE

Abstract. In the present paper we prove several sharp two-weight Hardy, Hardy-Poincaré, and Rellich type inequalities on the sub-Riemannian manifold $\mathbb{R}^{2n+1} = \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}$ defined by the vector fields:

$$X_j = \frac{\partial}{\partial x_j} + 2ky_j|z|^{2k-2}\frac{\partial}{\partial l}, \quad Y_j = \frac{\partial}{\partial y_j} - 2kx_j|z|^{2k-2}\frac{\partial}{\partial l}, \quad j = 1, 2, \dots, n$$

where $(z, y) = (x, y, l) \in \mathbb{R}^{2n+1}$, $|z| = (|x|^2 + |y|^2)^{1/2}$ and $k \geq 1$.

Mathematics subject classification (2010): 22E30, 26D10, 43A80.

Keywords and phrases: Hardy inequality with two weight functions, Rellich inequality with two weight functions, sharp constant.

REFERENCES

- [1] B. ABDELLAOUI AND I. PERAL, *Existence and nonexistence results for quasilinear elliptic equations involving the p -Laplacian with a critical potential*, Annali di Matematica **182** (2003), 247–270.
- [2] B. ABDELLAOUI, I. PERAL AND A. PRIMA, *Strong regularizing effect of a gradient term in the heat equation with the Hardy potential*, J. Func. Anal **258** (2010), 1247–1272.
- [3] S. AHMETOLAN AND S. CAVDAR, *On Nonlinear Degenerate Parabolic Equations with Time Dependent Singular Potential Functions*, Mathematische Nachrichten **284**, 10 (2011), 1219–1233.
- [4] S. AHMETOLAN AND I. KOMBE, *A sharp uncertainty principle and Hardy-Poincaré inequalities on sub-Riemannian manifolds*, Mathematical Inequalities & Applications **15**, 2 (2012), 457–467.
- [5] G. J. AZORERO AND I. PERAL, *Hardy inequalities and some critical elliptic and parabolic problems*, J. Diff. Equations **144** (1998), 441–476.
- [6] P. BARAS AND J. A. GOLDSTEIN, *The Heat Equation with a Singular Potential*, Transactions of the American Mathematical Society **284**, 1 (1984), 121–139.
- [7] R. BEALS, B. GAVEAU AND P. GREINER, *On a Geometric Formula for the Fundamental Solution of Subelliptic Laplacians*, Mathematische Nachrichten **181** (1996), 81–163.
- [8] R. BEALS, B. GAVEAU AND P. GREINER, *Green's Functions for Some Highly Degenerate Elliptic Operators*, Journal of Functional Analysis **165** (1999), 407–429.
- [9] H. BREZIS AND J. L. VÁZQUEZ, *Blow-up solutions of some nonlinear elliptic problems*, Rev. Mat. de la Universidad Complutense de Madrid **10**, 2 (1997), 443–469.
- [10] X. CABRÉ AND Y. MARTEL, *Existence versus explosion instantanée pour des équations de la chaleur linéaires avec potentiel singulier*, C. R. Acad. Sci. Paris **329**, 1 (1999), 973–978.
- [11] G. B. FOLLAND, *Fundamental solution for a subelliptic operator*, Bull. Amer. Math. Soc. **79**, 2 (1973), 373–376.
- [12] J. A. GOLDSTEIN AND I. KOMBE, *Nonlinear Degenerate Parabolic Equations with Singular Lower-Order Term*, Advances in Differential Equations **8**, 10 (2003), 1153–1192.
- [13] G. R. GOLDSTEIN, J. A. GOLDSTEIN AND A. RHANDI, *Weighted Hardy's inequality and the Kolmogorov equation perturbed by an inverse-square potential*, Appl. Anal. **91**, 11 (2012), 2057–2071.
- [14] P. C. GREINER, *A fundamental Solution for non-elliptic partial differential operator*, Canadian Journal of Mathematics **31** (1979), 1107–1120.
- [15] G. H. HARDY, *Note on a theorem of Hilbert*, Math. Zeitschr. **6** (1920), 314–317.

- [16] D. HAUER AND A. RHANDI, *A weighted Hardy inequality and nonexistence of positive solutions*, Arch. Math. (Basel) **100**, 3 (2013), 273–287.
- [17] L. HÖRMANDER, *Hypoelliptic Second Order Differential Equations*, Acta Mathematica **119** (1967), 147–171.
- [18] I. KOMBE, *The Linear Heat Equation with Highly Oscillating Potential*, Proceeding of the American Mathematical Society **132**, 9 (2004), 2683–2691.
- [19] I. KOMBE, *On the Nonexistence of Positive Solutions to Nonlinear Degenerate Parabolic Equations with Singular Coefficients*, Applicable Analysis **85**, 5 (2006), 467–478.
- [20] I. KOMBE, *Sharp Weighted Rellich and Uncertainty Principle inequalities on Carnot Groups*, Communications in Applied Analysis **14** (2010), no. 2, 251 **14**, 2 (2010), 251–272.
- [21] A. MORADIFAM AND N. GHOUSSOUB, *Bessel pairs and optimal Hardy and Hardy-Rellich inequalities*, Math. Ann. **349**, 1 (2011), 1–57.
- [22] P. NIU, Y. OU AND J. HAN, *Several Hardy type inequalities with Weights related to generalized greiner operator*, Canadian Math. Bulletin **53**, 1 (2010), 153–162.
- [23] P. NIU, H. ZHANG AND Y. WANG, *Hardy Thype and Rellich Type Inequalities on the Heisenberg Group*, Proceeding. Amer. Math. Soc. **129** (2001), 3623–3630.
- [24] I. PERAL AND J. L. VÁZQUEZ, *On the stability or insitability of the singular solution of the semilinear heat equation with exponential reaction term*, Arch. Rational Mech. Anal. **129** (1995), 201–224.
- [25] F. RELLICH, *Halbbeschränkte Differentialoperatoren höherer Ordnung*, Proceedings of the International Congress of Mathematicians, Edit by J. C. H. Gerretsen et all, Amsterdam **3** (1956), 243–250.
- [26] J. VANCOSTENOBLE AND E. ZUAZUA, *Hardy inequalities, observability, and control for the wave and Schrödinger equations with singular potentials*, SIAM journal on Math. Anal. **41**, 4 (2009), 1508–1532.
- [27] J. L. VÁZQUEZ AND E. ZUAZUA, *The Hardy constant and the asymptotic behaviour of the heat equation with an inverse-square potential*, J. Funct. Anal. **173** (2000), 103–153.
- [28] H. ZHANG AND P. NIU, *Hardy-type inequalities and Pohozaev-type identities for a class of p -degenerate subelliptic operators and applications*, Nonlinear Anal. **54**, 1 (2003), 165–186.