

A NEW SCHAUDER BASIS FOR $L^r((0,1)^n)$, $n = 2, 3$

JAN LANG, OSVALDO MÉNDEZ AND BEHZAD ROUHANI

Abstract. We show that under suitable conditions on p , q , and summability, the system of generalized trigonometric functions $\{\prod_{i=1}^d \sin_{p,q}(n_i \pi_{p,q} x_i)\}_{n_1, \dots, n_d}$ is a basis for $L^r((0,1)^d)$ for any $r \in (1, \infty)$ where $d = 2$ or $d = 3$.

Mathematics subject classification (2010): 33E30, 35P10, 35P30, 41A58.

Keywords and phrases: p -Laplacian, generalized trigonometric functions, Riesz theorem, Schauder basis, multi-dimensional Fourier series.

REFERENCES

- [1] E. AZROUL, A. BARBARA AND M. RHOUdaf, *On some nonhomogeneous nonlinear elliptic equations with nonstandard growth arising in image processing*, 1st Spring School on Numerical Methods for Partial Differential Equations, Tetouan, Morocco, 2010.
- [2] P. BUSHELL, P. AND D. EDMUNDS, *Remarks on generalised trigonometric functions*, Rocky Mountain J. Math. **42** 1 (2012), 25–57.
- [3] Y. CHEN, S. LEVINE AND M. RAO, *Variable exponent, linear growth functionals in image restoration*, SIAM J. Appl. Math. **66** 4 (2006), 1383–1406.
- [4] Y. CHEN, S. LEVINE AND M. RAO, *Functionals with $p(x)$ -growth in image processing*, Duquesne University, Department of Mathematics and Computer Science Technical Report 2004-01, available at www.mathcs.duq.edu/~sel/CLR05SIAPfinal.pdf.
- [5] D. EDMUNDS, P. GURKA AND J. LANGS, *Properties of generalized trigonometric functions*, J. Approx. Theory (2012), 47–56.
- [6] J. LANG AND D. EDMUNDS, *Eigenvalues, Embeddings and Generalised Trigonometric Functions*, Lecture Notes in Math. **2016**, Springer (2011).
- [7] V. LEVIN, *On a class of integral inequalities*, Rec. Math. Moscow **4** (1938), 309–331.
- [8] E. LUNDBERG, *Om hypergonometriska funktioner af komplexa variabla*, Stockholm (1879).
- [9] E. SCHMIDT, *Über die Ungleichung, welche die Integrale über eine Potenz einer Funktion und über eine andere Potenz ihrer Ableitung verbindet*, Math Ann. **117** (1940), 301–326.