

TWO-WEIGHT INEQUALITIES FOR GEOMETRIC MAXIMAL OPERATORS

ADAM OSĘKOWSKI

Abstract. We study one- and two-weight inequalities for the geometric maximal operator on probability spaces equipped with a tree-like structure. We provide a characterization of weights, in terms of Muckenhoupt and Sawyer-type conditions, for which the appropriate strong-type estimates hold. Our approach rests on Bellman function method, which allows us to identify sharp constants involved in the estimates.

Mathematics subject classification (2010): 42B25, 46E30, 60G42.

Keywords and phrases: Maximal, dyadic, Bellman function, best constants.

REFERENCES

- [1] R. BAÑUELOS AND A. OSĘKOWSKI, *Sharp weak type inequalities for fractional integral operators*, Potential Anal. **47** (2017), 103–121.
- [2] D. CRUZ-URIBE, *The minimal operator and the geometric maximal operator in \mathbb{R}^n* , Studia Math. **144** (2001), no. 1, 1–37.
- [3] D. CRUZ-URIBE AND C. J. NEUGEBAUER, *Weighted norm inequalities for the geometric maximal operator*, Publ. Mat. **42** (1998), 239–263.
- [4] C. FEFFERMAN AND E. M. STEIN, *Some maximal inequalities*, Amer. J. Math. **93** (1971), pp. 107–115.
- [5] A. D. MELAS, *The Bellman functions of dyadic-like maximal operators and related inequalities*, Adv. Math. **192** (2005), 310–340.
- [6] F. NAZAROV AND S. TREIL, *The hunt for Bellman function: applications to estimates of singular integral operators and to other classical problems in harmonic analysis*, Algebra i Analiz **8** (1997), 32–162.
- [7] P. ORTEGA SALVADOR AND C. RAMÍREZ TORREBLANCA, *Weighted inequalities for the one-sided geometric maximal operators*, Math. Nachr. **284** (2011), 1515–1522.
- [8] A. OSĘKOWSKI, *Sharp estimates for Lipschitz class*, J. Geom. Anal. **26** (2016), 1346–1369.
- [9] A. OSĘKOWSKI, *Sharp weighted bounds for geometric maximal operators*, Glasgow Math. J. **59** (2017), 533–547.
- [10] X. SHI, *Two inequalities related to geometric mean operators*, J. Zhejiang Teacher College **1** (1980), 21–25.
- [11] L. SLAVIN, A. STOKOLOS AND V. VASYUNIN, *Monge-Ampère equations and Bellman functions: The dyadic maximal operator*, C. R. Acad. Sci. Paris, Ser. I **346** (2008), 585–588.
- [12] L. SLAVIN AND V. VASYUNIN, *Sharp results in the integral-form John-Nirenberg inequality*, Trans. Amer. Math. Soc. **363** (2011), 4135–4169.
- [13] L. SLAVIN AND A. VOLBERG, *Bellman function and the H^1 -BMO duality*, Harmonic analysis, partial differential equations, and related topics, 113–126, Contemp. Math., 428, Amer. Math. Soc., Providence, RI, 2007.
- [14] V. VASYUNIN AND A. VOLBERG, *Monge-Ampère equation and Bellman optimization of Carleson embedding theorems*, Linear and complex analysis, pp. 195–238, Amer. Math. Soc. Transl. Ser. 2, 226, Amer. Math. Soc., Providence, RI, 2009.
- [15] V. VASYUNIN AND A. VOLBERG, *Burkholder's function via Monge-Ampère equation*, Illinois J. Math. **54** (2010), 1393–1428.

- [16] J. WITTWER, *Survey article: a user's guide to Bellman functions*, Rocky Mountain J. Math. **41** (2011), 631–661.
- [17] X. YIN AND B. MUCKENHOUPT, *Weighted inequalities for the maximal geometric mean operator*, Proc. Amer. Math. Soc. **124** (1996), 75–81.