

ON APPROXIMATING THE ERROR FUNCTION

ZHEN-HANG YANG, WEI-MAO QIAN, YU-MING CHU AND WEN ZHANG

(Communicated by J Pečarić)

Abstract. In the article, we find the best possible parameters p and q on the interval $(7/5, (7\pi - 20)/(5\pi - 15))$ such that the double inequality

$$\sqrt{1 - \lambda(p)e^{-px^2} - (1 - \lambda(p))e^{-\mu(p)x^2}} < \text{erf}(x) < \sqrt{1 - \lambda(q)e^{-qx^2} - (1 - \lambda(q))e^{-\mu(q)x^2}}$$

holds for all $x > 0$, where $\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$ is the error function, $\lambda(p) = 4[(7\pi - 20) - 5(\pi - 3)p]/[\pi(15p^2 - 40p + 28)]$, $\mu(p) = 4(5p - 7)/[5(3p - 4)]$.

1. Introduction

Let $x > 0$. Then the classical error function $\text{erf}(x)$ is given by

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt.$$

It is well known that the error function $\text{erf}(x)$ is one of the most important special functions, it has many important applications in probability, statistics and partial differential equations theory. Recently, the special functions have attracted the attention of many mathematicians [8, 10–16, 26–34, 38, 41–45]. In particular, many remarkable inequalities for the error function can be found in the literature [1, 5, 6, 9, 17–23, 35, 36].

Pólya [25] proved that the inequality

$$\text{erf}(x) < \sqrt{1 - e^{-4x^2/\pi}}$$

holds for all $x > 0$.

In [7], Chu proved that the double inequality

$$\sqrt{1 - e^{-px^2}} < \text{erf}(x) < \sqrt{1 - e^{-qx^2}}$$

holds for all $x > 0$ if and only if $p \in (0, 1]$ and $q \in [4/\pi, \infty)$.

Mathematics subject classification (2010): 33B20, 26D15, 26A48.

Keywords and phrases: error function, gamma function, monotonicity, bound.

The research was supported by the Natural Science Foundation of China under Grants 61673169, 11371125, 11701176 and 11401191, and the Tianyuan Special Funds of the National Natural Science Foundation of China under Grant 11626101.

Alzer [2] presented the double inequality

$$\left(1 - e^{-\beta(p)x^p}\right)^{1/p} < \frac{1}{\Gamma\left(1 + \frac{1}{p}\right)} \int_0^x e^{-t^p} dt < \left(1 - e^{-\alpha(p)x^p}\right)^{1/p}$$

for $x > 0$ and $p > 0$ with $p \neq 1$, where $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$ is the classical gamma function, and $\alpha(p)$ and $\beta(p)$ are respectively given by

$$\alpha(p) = \frac{1}{\Gamma^p\left(1 + \frac{1}{p}\right)} \quad (p > 1), \quad \alpha(p) = 1 \quad (0 < p < 1)$$

and

$$\beta(p) = \frac{1}{\Gamma^p\left(1 + \frac{1}{p}\right)} \quad (0 < p < 1), \quad \beta(p) = 1 \quad (p > 1).$$

Let $n \geq 2$, and $\alpha_n, \beta_n, \alpha_n^*, \beta_n^*$ be respectively defined by

$$\alpha_2 = 0.90686 \dots, \quad \alpha_n = 1 \quad (n \geq 3), \quad \beta_n = n - 1,$$

$$\alpha_n^* = n + 1 \quad (n = 2k), \quad \alpha_n^* = n - 1 \quad (n = 2k - 1), \quad \beta_n^* = 1.$$

In [3, 4], Alzer proved that the double inequalities

$$\lambda_n \operatorname{erf}\left(\sum_{i=1}^n x_i\right) \leq \sum_{i=1}^n \operatorname{erf}(x_i) - \prod_{i=1}^n \operatorname{erf}(x_i) \leq \mu_n \operatorname{erf}\left(\sum_{i=1}^n x_i\right), \quad (1.1)$$

$$\lambda \operatorname{erf}(y + \operatorname{erf}(x)) < \operatorname{erf}(x + \operatorname{erf}(y)) < \mu \operatorname{erf}(y + \operatorname{erf}(x)),$$

$$\lambda^* \operatorname{erf}(y \operatorname{erf}(x)) < \operatorname{erf}(x \operatorname{erf}(y)) \leq \mu^* \operatorname{erf}(y \operatorname{erf}(x))$$

hold for all $x_i \geq 0$ and $y \geq x > 0$ if and only if $\lambda_n \leq \alpha_n, \mu_n \geq \beta_n, \lambda \leq \operatorname{erf}(1) = 0.8427 \dots, \mu \geq 2/\sqrt{\pi} = 1.1283 \dots, \lambda^* \leq 0$ and $\mu^* \geq 1$, and inequality (1.1) holds for all $x_i \leq 0$ if and only if $\lambda_n \geq \alpha_n^*$ and $\mu_n \leq \beta_n^*$.

Neuman [24] proved that the double inequality

$$\frac{2x}{\sqrt{\pi}} e^{-\frac{x^2}{3}} \leq \operatorname{erf}(x) \leq \frac{2x}{\sqrt{\pi}} \frac{e^{-x^2} + 2}{3}$$

holds for all $x > 0$.

Let $\alpha \in (0, 1), \lambda(p), \mu = \mu(p), \eta(p)$ and $B(p, \alpha; x)$ be defined by

$$\lambda(p) = \frac{4[(7\pi - 20) - 5(\pi - 3)p]}{\pi(15p^2 - 40p + 28)}, \quad (1.2)$$

$$\mu(p) = \frac{4(5p - 7)}{5(3p - 4)}, \quad (1.3)$$

$$\eta(p) = \frac{16(5p - 7)}{(15p^2 - 40p + 28)(45p^2 - 60p - 4)}, \quad (1.4)$$

$$B(p, \alpha; x) = \sqrt{1 - \alpha e^{-px^2} - (1 - \alpha)e^{-\mu(p)x^2}}, \quad (1.5)$$

respectively.

Very recently, Yang, Chu and Zhang [37] provide necessary and sufficient conditions for the parameters r and s on the interval $(7/5, \infty)$ such that the double inequality

$$B(r, \eta(r); x) < \operatorname{erf}(x) < B(s, \eta(s); x)$$

holds for all $x > 0$.

From (1.2) we clearly see that $\lambda(p) \in (0, 1)$ for $p \in (7/5, (7\pi - 20)/(5\pi - 15))$. The main purpose of this paper is to present the best possible parameters p and q on the interval $(7/5, (7\pi - 20)/(5\pi - 15))$ such that the double inequality

$$B(p, \lambda(p); x) < \operatorname{erf}(x) < B(q, \lambda(q); x)$$

holds for all $x > 0$.

2. Lemmas

In order to prove our main results, we need to introduce an auxiliary function at first.

Let $-\infty \leq a < b \leq \infty$, f and g be differentiable on (a, b) , and $g' \neq 0$ on (a, b) . Then the function $H_{f,g}$ [40] is defined by

$$H_{f,g} \equiv \frac{f'}{g'} g - f. \quad (2.1)$$

LEMMA 2.1. (See [37, Lemma 2.1]) *Let $-\infty \leq a < b \leq \infty$, f and g be differentiable on (a, b) with $f(a^+) = g(a^+) = 0$, $g'(x) \neq 0$ and $g'(x)H_{f,g}(b^-) < (>)0$. If there exists $\lambda_0 \in (a, b)$ such that f'/g' is strictly increasing (decreasing) on (a, λ_0) and strictly decreasing (increasing) on (λ_0, b) , then there exists $\mu_0 \in (a, b)$ such that f/g is strictly increasing (decreasing) on (a, μ_0) and strictly decreasing (increasing) on (μ_0, b) .*

LEMMA 2.2. (See [39, Lemma 7]) *Let $a_i \geq 0$ for all $i = 0, 1, 2, \dots$ with $\sum_{i=0}^m a_i > 0$ and $\sum_{i=m+1}^{\infty} a_i > 0$, and*

$$P(t) = \sum_{i=0}^m a_i t^i - \sum_{i=m+1}^{\infty} a_i t^i$$

be a convergent power series on the interval $(0, \infty)$. Then there exists $t_0 \in (0, \infty)$ such that $P(t_0) = 0$, $P(t) > 0$ for $t \in (0, t_0)$ and $P(t) < 0$ for $t \in (t_0, \infty)$.

LEMMA 2.3. (See [37, Lemma 2.2 (2)]) *Let $u_n = (5p - 6)(5p - 8)n - (15p^2 - 40p + 28)$. Then $u_n < 0$ for all $n \geq 2$ if $p \in (7/5, 8/5]$.*

LEMMA 2.4. See [37, Theorem 3.2]) Let $\eta(p)$ and $B(p, \alpha; x)$ be defined by (1.4) and (1.5), respectively. Then the inequality

$$\operatorname{erf}(x) < B(p_0, \eta(p_0); x)$$

holds for $x > 0$, where $p_0 = (21\pi - 60 + \sqrt{3(147\pi^2 - 920\pi + 1440)})/[30(\pi - 3)] = 1.713\cdots$.

LEMMA 2.5. Let $x \in (0, \infty)$, $\lambda(p)$, $\mu(p)$, $B(p, \alpha; x)$ and $H_{f,g}(x)$ be respectively defined by (1.2), (1.3), (1.5) and (2.1), and $f_1(x)$, $g_1(x)$, $f_2(x)$ and $g_2(x)$ be respectively defined by

$$f_1(x) = B^2(p, \lambda(p); x) = 1 - \lambda(p)e^{-px^2} - (1 - \lambda(p))e^{-\mu(p)x^2}, \quad g_1(x) = \operatorname{erf}^2(x), \quad (2.2)$$

$$f_2(x) = \left[p\lambda(p)e^{(1-p)x^2} + \mu(p)(1 - \lambda(p))e^{(1-\mu(p))x^2} \right] x, \quad g_2(x) = \frac{2}{\sqrt{\pi}} \operatorname{erf}(x). \quad (2.3)$$

Then

$$H_{f_1, g_1}(\infty) = \lim_{x \rightarrow \infty} \left(\frac{f'_1(x)}{g'_1(x)} g_1(x) - f_1(x) \right) = \infty,$$

$$H_{f_2, g_2}(\infty) = \lim_{x \rightarrow \infty} \left(\frac{f'_2(x)}{g'_2(x)} g_2(x) - f_2(x) \right) = \infty$$

for $p \in (7/5, 8/5]$.

Proof. Let $t = (p - \mu(p))x^2$ and

$$\begin{aligned} h_1(t) &= p\lambda(p)(p - \mu(p)) - 2p\lambda(p)(p - 1)t \\ &\quad + \mu(p)(p - \mu(p))(1 - \lambda(p))e^t - 2\mu(p)(\mu(p) - 1)(1 - \lambda(p))te^t. \end{aligned} \quad (2.4)$$

Then (2.1)–(2.4) lead to

$$\begin{aligned} H_{f_1, g_1}(x) &= \frac{\sqrt{\pi}}{2} x \operatorname{erf}(x) \left[p\lambda(p)e^{(1-p)x^2} + \mu(p)(1 - \lambda(p))e^{(1-\mu(p))x^2} \right] \\ &\quad - \left[1 - \lambda(p)e^{-px^2} - (1 - \lambda(p))e^{-\mu(p)x^2} \right], \end{aligned} \quad (2.5)$$

$$H_{f_2, g_2}(x) = te^{\frac{1-\mu(p)}{p-\mu(p)}t} \left[\frac{\sqrt{\pi} e^{\frac{2-p}{p-\mu(p)}t}}{2(p - \mu(p))} \operatorname{erf}(x) \frac{h_1(t)}{te^{\frac{1-\mu(p)}{p-\mu(p)}t}} - \frac{p\lambda(p)e^{-t} + \mu(p)(1 - \lambda(p))}{\sqrt{(p - \mu(p))t}} \right]. \quad (2.6)$$

If $p \in (7/5, 8/5]$, then it is not difficult to verify that

$$p > \mu(p), \quad 0 < \lambda(p) < 1, \quad 0 < \mu(p) \leqslant 1, \quad (2.7)$$

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{h_1(t)}{te^{\frac{1-\mu(p)}{p-\mu(p)}t}} &= \lim_{t \rightarrow \infty} \left[\frac{p\lambda(p)(p-\mu(p))}{t} e^{\frac{\mu(p)-1}{p-\mu(p)}t} - 2p\lambda(p)(p-1)e^{\frac{\mu(p)-1}{p-\mu(p)}t} \right] \quad (2.8) \\ &\quad + \lim_{t \rightarrow \infty} \left[\frac{\mu(p)(1-\lambda(p))(p-\mu(p))}{t} e^{\frac{p-1}{p-\mu(p)}t} \right. \\ &\quad \left. + 2\mu(p)(1-\lambda(p))(1-\mu(p))e^{\frac{p-1}{p-\mu(p)}t} \right] \\ &= \infty. \end{aligned}$$

Therefore, Lemma 2.5 follows easily from (2.4)–(2.8). \square

LEMMA 2.6. *Let $x > 0$, and $\lambda(p)$ and $B(p, \alpha; x)$ be respectively defined by (1.2) and (1.5). Then the function $p \mapsto B(p, \lambda(p); x)$ is strictly increasing on $(7/5, (7\pi - 20)/(5\pi - 15))$*

Proof. Let $\mu(p)$ be defined by (1.3), $t = (p - \mu(p))x^2$ and $f(t)$ be defined by

$$f(t) = -[p - \mu(p)]\lambda'(p) + \lambda(p)t + [p - \mu(p)]\lambda'(p)e^t + \mu'(p)[1 - \lambda(p)]te^t.$$

Then elaborated computations lead to

$$2B(p, \lambda(p); x) \frac{\partial B(p, \lambda(p); x)}{\partial p} = \frac{e^{-px^2}}{p - \mu(p)} f(t), \quad (2.9)$$

$$f(0) = 0, \quad (2.10)$$

$$f'(t) = \lambda(p) + [p - \mu(p)]\lambda'(p)e^t + \mu'(p)[1 - \lambda(p)]e^t + \mu'(p)[1 - \lambda(p)]te^t,$$

$$f'(0) = 0, \quad (2.11)$$

$$f''(t) = \frac{4(\pi - 3)}{(3p - 4)\pi} e^t + \frac{4(\pi p - 4)}{(3p - 4)(15p^2 - 40p + 28)} te^t > 0 \quad (2.12)$$

for $t > 0$ and $p \in (7/5, (7\pi - 20)/(5\pi - 15))$.

Therefore, Lemma 2.6 follows easily from (2.9)–(2.12) and $p > \mu(p)$ for $p \in (7/5, (7\pi - 20)/(5\pi - 15))$. \square

3. Main results

THEOREM 3.1. *Let $\alpha \in (0, 1)$, $p \in (7/5, 8/5]$, and $\lambda(p)$ and $B(p, \alpha; x)$ be defined by (1.2) and (1.5), respectively. Then the inequality*

$$\text{erf}(x) > B(p, \alpha; x) \quad (3.1)$$

holds for all $x > 0$ if and only if $\alpha \leq \lambda(p)$.

Proof. If inequality (3.1) holds for all $x > 0$, then we clearly see that

$$\lim_{x \rightarrow 0^+} \frac{\operatorname{erf}(x) - B(p, \alpha; x)}{x} = \frac{2}{\sqrt{\pi}} - \sqrt{\alpha p + (1-\alpha)\mu(p)} \geq 0,$$

$$\alpha \leq \frac{4 - \pi\mu(p)}{\pi(p - \mu(p))} = \lambda(p).$$

Next, we prove that inequality (3.1) holds for all $x > 0$ if $\alpha \leq \lambda(p)$. We only need to prove that the inequality

$$\operatorname{erf}(x) > B(p, \lambda(p); x) \quad (3.2)$$

holds for all $x > 0$ due to the function $\alpha \mapsto B(p, \alpha; x)$ is strictly increasing on $(0, 1)$.

Let $t = (p - \mu(p))x^2$, $\mu(p)$, $\eta(p)$, u_n , $f_1(x)$, $g_1(x)$, $f_2(x)$, $g_2(x)$ and $h_1(t)$ be respectively defined by (1.3), (1.4), Lemma 2.3, (2.2), (2.3) and (2.4), and $h_2(t)$ be defined by

$$\begin{aligned} h_2(t) = & 2\mu(p)(1 - \lambda(p))(\mu(p) - 1)(\mu(p) - 2)te^t - \mu(p)(1 - \lambda(p))(3\mu(p) - 4) \\ & \times (p - \mu(p))e^t + 2p\lambda(p)(p - 1)(p - 2)t - p\lambda(p)(3p - 4)(p - \mu(p)), \end{aligned}$$

Then elaborated computations lead to

$$\frac{f'_1(x)}{g'_1(x)} = \frac{f_2(x)}{g_2(x)}, \quad (3.3)$$

$$\frac{f'_2(x)}{g'_2(x)} = \frac{\pi}{4(p - \mu(p))} e^{\frac{2-p}{p-\mu(p)}t} h_1(t),$$

$$\left(\frac{f'_2(x)}{g'_2(x)} \right)' = \frac{\pi}{4(p - \mu(p))} \frac{d}{dt} \left[e^{\frac{2-p}{p-\mu(p)}t} h_1(t) \right] \frac{dt}{dx} = \frac{\pi x}{2(p - \mu(p))} e^{\frac{2-p}{p-\mu(p)}t} h_2(t), \quad (3.4)$$

$$h_2(t) = 2\mu(p)(1 - \lambda(p))(\mu(p) - 1)(\mu(p) - 2) \sum_{n=1}^{\infty} \frac{t^n}{(n-1)!} \quad (3.5)$$

$$- \mu(p)(1 - \lambda(p))(3\mu(p) - 4)(p - \mu(p)) \sum_{n=0}^{\infty} \frac{t^n}{n!}$$

$$+ 2p\lambda(p)(p - 1)(p - 2)t - p\lambda(p)(3p - 4)(p - \mu(p))$$

$$= - \frac{(15p^2 - 40p + 28)^2(45p^2 - 60p - 4)}{125(3p - 4)^3} (\lambda(p) - \eta(p))$$

$$- \frac{2(p-1)(2-p)}{25(3p-4)^3} (\lambda(p) - \eta(p))t - \frac{4\mu(p)(1-\lambda(p))}{25(3p-4)^2} \sum_{n=2}^{\infty} \frac{u_n}{n!} t^n,$$

$$\lambda(p) - \eta(p) = \frac{20(3p-4)(p_0-p)(p-p_1)}{\pi(15p^2 - 40p + 28)(45p^2 - 60p - 4)}, \quad (3.6)$$

where

$$p_0 = \frac{21\pi - 60 + \sqrt{3(147\pi^2 - 920\pi + 1440)}}{30(\pi - 3)} = 1.713 \dots, \quad (3.7)$$

$$p_1 = \frac{21\pi - 60 - \sqrt{3(147\pi^2 - 920\pi + 1440)}}{30(\pi - 3)} = 1.099\dots \quad (3.8)$$

From (2.7) and (3.4)–(3.8) together with Lemmas 2.2 and 2.3 we clearly see that there exists $x_0 \in (0, \infty)$ such that $f'_2(x)/g'_2(x)$ is strictly decreasing on $(0, x_0)$ and strictly increasing on (x_0, ∞) . Then Lemmas 2.1 and 2.5 together with (3.3), $f_2(0) = g_2(0) = 0$ and $g'_2(x) > 0$ lead to the conclusion that there exists $x_1 \in (0, \infty)$ such that $f'_1(x)/g'_1(x)$ is strictly decreasing on $(0, x_1)$ and strictly increasing on (x_1, ∞) .

It follows from Lemma 2.1, $f_1(0) = g_1(0) = 0$ and $g'_1(x) > 0$ together with the piecewise monotonicity of $f'_1(x)/g'_1(x)$ that there exists $x_2 \in (0, \infty)$ such that $f_1(x)/g_1(x)$ is strictly decreasing on $(0, x_2)$ and strictly increasing on (x_2, ∞) .

Note that

$$\lim_{x \rightarrow 0^+} \frac{f_1(x)}{g_1(x)} = \lim_{x \rightarrow 0^+} \frac{f'_2(x)}{g'_2(x)} = \frac{\pi[p\lambda(p) + \mu(p)(1 - \lambda(p))]}{4} = 1, \quad \lim_{x \rightarrow \infty} \frac{f_1(x)}{g_1(x)} = 1. \quad (3.9)$$

Therefore, inequality (3.2) follows easily from (2.2) and (3.9) together with the piecewise monotonicity of $f_1(x)/g_1(x)$. \square

THEOREM 3.2. *Let $p_0 = (21\pi - 60 + \sqrt{3(147\pi^2 - 920\pi + 1440)})/[30(\pi - 3)] = 1.713\dots$, $\alpha \in (0, 1)$, $p \in [p_0, (7\pi - 20)/(5\pi - 15))$, and $\lambda(p)$ and $B(p, \alpha; x)$ be defined by (1.2) and (1.5), respectively. Then the inequality*

$$\operatorname{erf}(x) < B(p, \alpha; x) \quad (3.10)$$

holds for all $x > 0$ if and only if $\alpha \geq \lambda(p)$.

Proof. If inequality (3.10) holds for all $x > 0$, then we clearly see that

$$\lim_{x \rightarrow 0^+} \frac{\operatorname{erf}(x) - B(p, \alpha; x)}{x} = \frac{2}{\sqrt{\pi}} - \sqrt{\alpha p + (1 - \alpha)\mu(p)} \leq 0,$$

$$\alpha \geq \frac{4 - \pi\mu(p)}{\pi(p - \mu(p))} = \lambda(p).$$

Next, we prove that inequality (3.10) holds for all $x > 0$ if $\alpha \geq \lambda(p)$. We only need to prove that the inequality

$$\operatorname{erf}(x) < B(p, \lambda(p); x) \quad (3.11)$$

holds for all $x > 0$ due to the function $\alpha \mapsto B(p, \alpha; x)$ is strictly increasing on $(0, 1)$.

Inequality (3.11) follows easily from Lemma 2.4, Lemma 2.6, (3.6) and $p \geq p_0$. Indeed, we have

$$\operatorname{erf}(x) < B(p_0, \eta(p_0); x) = B(p_0, \lambda(p_0); x) \leq B(p, \lambda(p); x)$$

for all $x > 0$. \square

THEOREM 3.3. Let $p_0 = (21\pi - 60 + \sqrt{3(147\pi^2 - 920\pi + 1440)})/[30(\pi - 3)] = 1.713\cdots$, $\alpha \in (0, 1)$, $p \in (7/5, (7\pi - 20)/(5\pi - 15))$, and $\lambda(p)$, $\mu(p)$ and $B(p, \alpha; x)$ be respectively defined by (1.2), (1.3) and (1.5). Then the inequality

$$\operatorname{erf}(x) > B(p, \lambda(p); x) \quad (3.12)$$

holds for all $x > 0$ if and only if $p \in (7/5, 8/5]$, and the inequality

$$\operatorname{erf}(x) < B(p, \lambda(p); x) \quad (3.13)$$

holds for all $x > 0$ if and only if $p \in [p_0, (7\pi - 20)/(5\pi - 15))$.

Proof. From Theorems 3.1 and 3.2 we clearly see that inequality (3.12) holds for all $x > 0$ if $p \in (7/5, 8/5]$, and inequality (3.13) holds for all $x > 0$ if $p \in [p_0, (7\pi - 20)/(5\pi - 15))$.

Next, we prove that $p \leq 8/5$ if inequality (3.12) holds for all $x > 0$. Indeed,

$$\begin{aligned} & \lim_{x \rightarrow \infty} \frac{\operatorname{erf}^2(x) - B^2(p, \lambda(p); x)}{e^{-\mu(p)x^2}} \\ &= - \lim_{x \rightarrow \infty} \frac{\frac{4}{\sqrt{\pi}} e^{-x^2} \operatorname{erf}(x) - 2x \left(p\lambda(p)e^{-px^2} + \mu(p)(1 - \lambda(p))e^{-\mu(p)x^2} \right)}{2\mu(p)x e^{-\mu(p)x^2}} \\ &= \lim_{x \rightarrow \infty} \left(-\frac{2\operatorname{erf}(x)e^{(\mu(p)-1)x^2}}{\sqrt{\pi}\mu(p)} + \frac{p\lambda(p)}{\mu(p)} e^{-(p-\mu(p))x^2} + 1 - \lambda(p) \right) = -\infty \end{aligned}$$

due to $p > \mu(p)$ and $\mu(p) > 1$ if $p > 8/5$.

Finally, we prove that $p \geq p_0$ if inequality (3.13) holds for all $x > 0$. Let $x \rightarrow 0^+$, then making use of Taylor formula we get

$$\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}}x - \frac{2}{3\sqrt{\pi}}x^3 + o(x^3), \quad (3.14)$$

$$B(p, \lambda(p); x) = \sqrt{p\lambda(p) + \mu(p)(1 - \lambda(p))}x - \frac{p^2\lambda(p) + \mu^2(p)(1 - \lambda(p))}{4\sqrt{p\lambda(p) + \mu(p)(1 - \lambda(p))}}x^3 + o(x^3). \quad (3.15)$$

Note that

$$\sqrt{p\lambda(p) + \mu(p)(1 - \lambda(p))} = \frac{2}{\sqrt{\pi}}. \quad (3.16)$$

If inequality (3.13) holds for all $x > 0$, then (1.2), (1.3) and (3.14)–(3.16) lead to

$$\lim_{x \rightarrow 0^+} \frac{\operatorname{erf}(x) - B(p, \lambda(p); x)}{x^3} = -\frac{15(\pi - 3)p^2 - 3(7\pi - 20)p + 4}{30\sqrt{\pi}(3p - 4)} \leq 0. \quad (3.17)$$

Therefore, $p \geq p_0$ follows easily from (3.17) and $p > 7/5$. \square

Let $\lambda(p)$, $\mu(p)$ and $B(p, \alpha; x)$ be respectively defined by (1.2), (1.3) and (1.5). Then simple computations lead to

$$\lambda\left(\frac{8}{5}\right) = \frac{5(4-\pi)}{3\pi}, \quad \mu\left(\frac{8}{5}\right) = 1, \quad (3.18)$$

$$\lambda\left(\frac{3}{2}\right) = \frac{8(5-\pi)}{7\pi}, \quad \mu\left(\frac{3}{2}\right) = \frac{4}{5}, \quad (3.19)$$

$$\lambda\left(\frac{7}{5}\right) = \frac{20}{7\pi}, \quad \mu\left(\frac{7}{5}\right) = 0, \quad (3.20)$$

$$\lambda(2) = \frac{10-3\pi}{2\pi}, \quad \mu(2) = \frac{6}{5}, \quad (3.21)$$

$$\lambda\left(\frac{7\pi-20}{5\pi-15}\right) = 0, \quad \mu\left(\frac{7\pi-20}{5\pi-15}\right) = \frac{4}{\pi}. \quad (3.22)$$

From Lemma 2.6, Theorem 3.1, Theorem 3.2 and (3.18)–(3.22) together with $7/5 < 3/2 < 8/5$ and $(21\pi - 60 + \sqrt{3(147\pi^2 - 920\pi + 1440)})/[30(\pi - 3)] = 1.713\cdots < 2 < (7\pi - 20)/(5\pi - 15) = 2.181\cdots$, we get Corollary 3.4 immediately.

COROLLARY 3.4. Let $p_0 = (21\pi - 60 + \sqrt{3(147\pi^2 - 920\pi + 1440)})/[30(\pi - 3)]$, and $\lambda(p)$ and $\mu(p)$ be respectively defined by (1.2) and (1.3). Then the inequalities

$$\begin{aligned} \sqrt{\frac{20}{7\pi}(1-e^{-7x^2/5})} &< \sqrt{1 - \frac{8(5-\pi)}{7\pi}e^{-3x^2/2} - \frac{5(3\pi-8)}{7\pi}e^{-4x^2/5}} \\ &< \sqrt{1 - \frac{5(4-\pi)}{3\pi}e^{-8x^2/5} - \frac{4(2\pi-5)}{3\pi}e^{-x^2}} \\ &< \operatorname{erf}(x) < \sqrt{1 - \lambda(p_0)e^{-p_0x^2} - (1 - \lambda(p_0))e^{-\mu(p_0)x^2}} \\ &< \sqrt{1 - \frac{10-3\pi}{2\pi}e^{-2x^2} - \frac{5(\pi-2)}{2\pi}e^{-6x^2/5}} < \sqrt{1 - e^{-4x^2/\pi}} \end{aligned}$$

hold for all $x > 0$.

REMARK 3.5. Let $\lambda(p)$, $\eta(p)$ and $B(p, \alpha; x)$ be respectively defined by (1.2), (1.4) and (1.5), and $p_0 = (21\pi - 60 + \sqrt{3(147\pi^2 - 920\pi + 1440)})/[30(\pi - 3)] = 1.713\cdots$. Then from (3.6) we clearly see that $\lambda(p) > \eta(p)$ for $p \in (7/5, 8/5]$ and $\lambda(p) < \eta(p)$ for $p \in (p_0, (7\pi - 20)/(5\pi - 15))$. Therefore, Theorems 3.1 and 3.2 together with the monotonicity of the function $\alpha \rightarrow B(p, \alpha; x)$ lead to the conclusion that

$$\operatorname{erf}(x) > B(p, \lambda(p); x) > B(p, \eta(p); x)$$

for all $x > 0$ and $p \in (7/5, 8/5]$, and

$$\operatorname{erf}(x) < B(p, \lambda(p); x) < B(p, \eta(p); x)$$

for all $x > 0$ and $p \in (p_0, (7\pi - 20)/(5\pi - 15))$.

REFERENCES

- [1] K. M. ALUDAAT AND M. T. ALODAT, *A note on approximating the normal distribution function*, Appl. Math. Sci. **2**, 9–12 (2008), 425–429.
- [2] H. ALZER, *On some inequalities for the incomplete gamma function*, Math. Comp. **66**, 218 (1997), 771–778.
- [3] H. ALZER, *Functional inequalities for the error function*, Aequationes Math. **66**, 1–2 (2003), 119–127.
- [4] H. ALZER, *Functional inequalities for the error function II*, Aequationes Math. **78**, 1–2 (2009), 113–121.
- [5] H. ALZER, *Error function inequalities*, Adv. Comput. Math. **33**, 3 (2010), 349–379.
- [6] A. V. BOYD, *Inequalities for Mills' ratio*, Rep. Statist. Appl. Res. Un. Jap. Sci. Engrs. **6** 1959, 44–46.
- [7] J. T. CHU, *On bounds for the normal integral*, Biometrika **42** 1955, 263–265.
- [8] Y.-M. CHU, M. ADIL KHAN, T. ALI AND S. S. DRAGOMIR, *Inequalities for α -fractional differentiable functions*, J. Inequal. Appl. **2017** 2017, Article 93, 12 pages.
- [9] Y.-M. CHU, Y.-M. LI AND W.-F. XIA, *Best possible inequalities for the harmonic mean of error function*, J. Inequal. Appl. **2014** 2014, Article 525, 9 pages.
- [10] Y.-M. CHU, Y.-F. QIU AND M.-K. WANG, *Hölder mean inequalities for the complete elliptic integrals*, Integral Transforms Spec. Funct. **23**, 7 (2012), 521–527.
- [11] Y.-M. CHU AND M.-K. WANG, *Optimal Lehmer mean bounds for the Toader mean*, Results Math. **61**, 3–4 (2012), 223–229.
- [12] Y.-M. CHU, M.-K. WANG, S.-L. QIU AND Y.-P. JIANG, *Bounds for complete elliptic integrals of the second kind with applications*, Comput. Math. Appl. **63**, 7 (2012), 1177–1184.
- [13] Y.-M. CHU, G.-D. WANG AND X.-H. ZHANG, *Schur convexity and Hadamard's inequality*, Math. Inequal. Appl. **13**, 4 (2010), 725–731.
- [14] Y.-M. CHU, G.-D. WANG AND X.-H. ZHANG, *The schur multiplicative and harmonic convexities of the complete symmetric function*, Math. Nachr. **284**, 5–6 (2011), 653–663.
- [15] Y.-M. CHU, W.-F. XIA AND X.-H. ZHANG, *The Schur concavity, Shur multiplicative and harmonic convexities of the second dual form of the Hamy symmetric function with applications*, J. Multivariate Anal. **105** 2012, 412–421.
- [16] Y.-M. CHU, X.-H. ZHANG AND G.-D. WANG, *The Schur geometrical convexity of the extended mean values*, J. Convex Anal. **15**, 4 (2008), 725–731.
- [17] Y.-M. CHU AND T.-H. ZHAO, *Concavity of the error function with respect to Hölder means*, Math. Inequal. Appl. **19**, 2 (2016), 589–595.
- [18] W. J. CODY, *Rational Chebyshev approximations for the error function*, Math. Comp. **23** 1969, 631–637.
- [19] A. GASULL AND F. UTZET, *Approximating Mills ratio*, J. Math. Anal. Appl. **420**, 2 (2014), 1832–1853.
- [20] W. GAUTSCHI, *Some elementary inequalities relating to the gamma and incomplete gamma function*, J. Math. Phys. **38** 1959/60, 77–81.
- [21] R. G. HART, *A close approximation related to the error function*, Math. Comp. **20** 1966, 600–602.
- [22] A. LAFORGIA AND S. SISMONDI, *Monotonicity results and inequalities for the gamma and error functions*, J. Comput. Appl. Math. **23**, 1 (1988), 25–33.
- [23] Y.-M. LI, W.-F. XIA, Y.-M. CHU AND X.-H. ZHANG, *Optimal lower and upper bounds for the geometric convex combination of the error function*, J. Inequal. Appl. **2015** 2015, Article 382, 8 pages.
- [24] E. NEUMAN, *Inequalities and bounds for the incomplete gamma function*, Results Math. **63**, 3–4 (2013), 1209–1214.
- [25] G. PÓLYA, *Remarks on computing the probability integral in one and two dimensions*, Proceedings of the Berkeley Symposium on Mathematical Statistics and Probability, University of California Press, Berkeley and Los Angeles, 1949.
- [26] M.-K. WANG AND Y.-M. CHU, *Asymptotical bounds for complete elliptic integrals of the second kind*, J. Math. Anal. Appl. **402**, 1 (2013), 119–126.
- [27] M.-K. WANG AND Y.-M. CHU, *Refinements of transformation inequalities for zero-balanced hypergeometric functions*, Acta Math. Sci. **37B**, 3 (2017), 607–622.
- [28] M.-K. WANG, Y.-M. CHU AND Y.-P. JIANG, *Ramanujan's cubic transformation inequalities for zero-balanced hypergeometric functions*, Rocky Mountain J. Math. **46**, 2 (2016), 679–691.
- [29] M.-K. WANG, Y.-M. CHU AND S.-L. QIU, *Asymptotical formulas for Gaussian and generalized hypergeometric functions*, Appl. Math. Comput. **276** 2016, 44–60.
- [30] M.-K. WANG, Y.-M. CHU, S.-L. QIU AND Y.-P. JIANG, *Bounds for the perimeter of an ellipse*, J. Approx. Theory **164**, 7 (2012), 928–937.

- [31] M.-K. WANG, Y.-M. CHU, Y.-F. QIU AND S.-L. QIU, *An optimal power mean inequality for the complete elliptic integrals*, Appl. Math. Lett. **24**, 6 (2011), 887–890.
- [32] M.-K. WANG, Y.-M. LI AND Y.-M. CHU, *Inequalities and infinite product formula for Ramanujan generalized modular equation function*, Ramanujan J. doi:10.1007/s11139-017-9888-3.
- [33] M.-K. WANG, S.-L. QIU, Y.-M. CHU AND Y.-P. JIANG, *Generalized Hersch-Pfluger distortion function and complete elliptic integrals*, J. Math. Anal. Appl. **385**, 1 (2012), 221–229.
- [34] G.-D. WANG, X.-H. ZHANG AND Y.-M. CHU, *Inequalities for the generalized elliptic integrals and modular functions*, J. Math. Anal. Appl. **331**, 2 (2007), 1275–1283.
- [35] W.-F. XIA AND Y.-M. CHU, *Optimal inequalities for the convex combination of error function*, J. Math. Inequal. **9**, 1 (2015), 85–99.
- [36] Z.-H. YANG AND Y.-M. CHU, *On approximating Mills ratio*, J. Inequal. Appl. **2015** 2015, Article 273, 14 pages.
- [37] Z.-H. YANG AND Y.-M. CHU, *On approximating the error function*, J. Inequal. Appl. **2016** 2016, Article 311, 17 pages.
- [38] Z.-H. YANG AND Y.-M. CHU, *A monotonicity property involving the generalized elliptic integral of the first kind*, Math. Inequal. Appl. **20**, 3 (2017), 729–735.
- [39] Z.-H. YANG, Y.-M. CHU AND X.-J. TAO, *A double inequality for the trigamma function and its applications*, Abstr. Appl. Anal. **2014** 2014, Article ID 702718, 9 pages.
- [40] Z.-H. YANG, Y.-M. CHU AND M.-K. WANG, *Monotonicity criterion for the quotient of power series with applications*, J. Math. Anal. Appl. **428**, 1 (2015), 587–604.
- [41] Z.-H. YANG, W.-M. QIAN, Y.-M. CHU AND W. ZHANG, *Monotonicity rule for the quotient of two functions and its application*, J. Inequal. Appl. **2017** 2017, Article 106, 13 pages.
- [42] Z.-H. YANG, W.-M. QIAN, Y.-M. CHU AND W. ZHANG, *On rational bounds for the gamma function*, J. Inequal. Appl. **2017** 2017, Article 210, 17 pages.
- [43] Z.-H. YANG, W. ZHANG AND Y.-M. CHU, *Sharp Gautsch inequality for parameter $0 < p < 1$* , Math. Inequal. Appl. **20**, 4 (2017), 1107–1120.
- [44] X.-H. ZHANG, G.-D. WANG AND Y.-M. CHU, *Remarks on generalized elliptic integrals*, Proc. Roy. Edinburgh, Sect. A **139**, 2 (2009), 417–426.
- [45] X.-H. ZHANG, G.-D. WANG AND Y.-M. CHU, *Convexity with respect to Hölder mean involving zero-balanced hypergeometric functions*, J. Math. Anal. Appl. **353**, 1 (2009), 256–259.

(Received June 21, 2017)

Zhen-Hang Yang

Department of Mathematics
Huzhou University
Huzhou 313000, P. R. China
e-mail: yzhkm@163.com

Wei-Mao Qian

School of Distance Education
Huzhou Broadcast and TV University
Huzhou 313000, P. R. China
e-mail: qwm661977@126.com

Yu-Ming Chu

Department of Mathematics
Huzhou University
Huzhou 313000, P. R. China
e-mail: chuyuming2005@126.com

Wen Zhang

Friedman Brain Institute
Icahn School of Medicine at Mount Sinai
New York, NY 10029, USA
e-mail: zhang.wen81@gmail.com