

LANDEN INEQUALITIES FOR A CLASS OF HYPERGEOMETRIC FUNCTIONS WITH APPLICATIONS

MIAO-KUN WANG AND YU-MING CHU

(Communicated by I. Perić)

Abstract. In this paper, we study a class of Gaussian hypergeometric function ${}_2F_1(a, b; (a+b+1)/2; x)$ ($a, b > 0$), and find the maximal regions of ab plane in the first quadrant where the well-known Landen identities for the complete elliptic integrals of the first kind turn on respective inequalities valid for each $x \in (0, 1)$. Besides, the generalized Grötzsch ring function with two parameters $\mu_{a,b}(r)$ is introduced, and the analogs of duplication formula satisfied by Grötzsch ring function $\mu(r)$ for $\mu_{a,b}(r)$, in the form of inequalities, will be derived.

1. Introduction

For real numbers a, b , and c with $c \neq 0, -1, -2, \dots$, the Gaussian hypergeometric function is defined by [1, 8, 30, 40, 48, 49, 55, 72]

$$F(a, b; c; x) = {}_2F_1(a, b; c; x) = \sum_{n=0}^{\infty} \frac{(a, n)(b, n)}{(c, n)} \frac{x^n}{n!}$$

for $x \in (-1, 1)$, where

$$(a, n) = a(a+1)(a+2)\cdots(a+n-1)$$

for $n = 1, 2, \dots$, and $(a, 0) = 1$ for $a \neq 0$. The function $F(a, b; c; x)$ is called zero-balanced when $c = a + b$. The limiting values for $F(a, b; c; x)$ at 1 are as follows (see [8, Theorem 1.19 and 1.48])

$$\begin{cases} F(a, b; c; 1) = \Gamma(c)\Gamma(c-a-b)/[\Gamma(c-a)\Gamma(c-b)], & a+b < c, \\ B(a, b)F(a, b; c; z) + \log(1-z) = R(a, b) + O((1-z)\log(1-z)), & a+b = c, \\ F(a, b; c; z) = (1-z)^{c-a-b}F(c-a, c-b; c; z), & a+b > c. \end{cases} \quad (1.1)$$

Mathematics subject classification (2010): 33E05, 33C05.

Keywords and phrases: Hypergeometric function, complete elliptic integrals, generalized Grötzsch ring function, Landen inequalities, duplication formula.

This research was supported by the Natural Science Foundation of China (Grant Nos. 11626101, 11401191, 11601485, 11371125, 11701176), the Science and Technology Research Program of Zhejiang Educational Committee (Grant No. Y201635325).

Here $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$ ($\operatorname{Re}(x) > 0$) [60, 61, 68, 69, 70, 74, 75, 76] and $B(p, q) = [\Gamma(p)\Gamma(q)]/[\Gamma(p+q)]$ ($\operatorname{Re}(p) > 0$, $\operatorname{Re}(q) > 0$) are the classical gamma function and beta functions, respectively, and

$$R(a, b) = -\psi(a) - \psi(b) - 2\gamma, \quad R\left(\frac{1}{2}, \frac{1}{2}\right) = \log 16,$$

$\psi(z) = \Gamma'(z)/\Gamma(z)$ ($\operatorname{Re}(z) > 0$) [66, 67, 77] and γ is the Euler-Mascheroni constant defined by

$$\gamma = \lim_{n \rightarrow \infty} \left(\sum_{k=1}^n \frac{1}{k} - \log n \right) = 0.577 \dots$$

As special cases of hypergeometric functions, the complete elliptic integral of the first kind $\mathcal{K}(r)$ and the second kind $\mathcal{E}(r)$ can be expressed as

$$\begin{aligned} \mathcal{K}(r) &= \frac{\pi}{2} F\left(\frac{1}{2}, \frac{1}{2}; 1; r^2\right) = \int_0^{\pi/2} (1 - r^2 \sin^2 t)^{-1/2} dt, \quad r \in [0, 1], \quad \mathcal{K}(1^-) = +\infty, \\ \mathcal{E}(r) &= \frac{\pi}{2} F\left(-\frac{1}{2}, \frac{1}{2}; 1; r^2\right) = \int_0^{\pi/2} (1 - r^2 \sin^2 t)^{1/2} dt, \quad r \in [0, 1], \end{aligned}$$

respectively. Elliptic integrals have been studied over three centuries. Many important applications were found in physics, engineering, theory of mean values, number theory and other related fields [1, 2, 13, 14, 16, 25, 26, 27, 28, 32, 37, 38, 41, 43, 47, 58]. Recently, they also occur frequently in geometric function theory, particularly, these integrals enable us to investigate a lot of special functions that are indispensable for the study of quasiconformal mappings [3, 4, 5, 6, 11, 33]. For more properties and inequalities for $\mathcal{K}(r)$ and $\mathcal{E}(r)$ see e.g. [15, 17, 18, 19, 20, 21, 22, 23, 24, 31, 45, 50, 51, 52, 53, 57, 62, 64, 65, 73]. Recall that some of the most important properties of the elliptic integral $\mathcal{K}(r)$ are Landen identities proved in 1771 [59, p. 507]

$$\mathcal{K}\left(\frac{2\sqrt{r}}{1+r}\right) = (1+r)\mathcal{K}(r), \quad \mathcal{K}\left(\frac{1-r}{1+r}\right) = \frac{1+r}{2}\mathcal{K}(r'). \quad (1.2)$$

Here and in what follows $r' \equiv \sqrt{1-r^2}$. Equation (1.2) is equivalent to

$$\begin{aligned} F\left(\frac{1}{2}, \frac{1}{2}; 1; \frac{4r}{(1+r)^2}\right) &= (1+r)F\left(\frac{1}{2}, \frac{1}{2}; 1; r^2\right), \\ F\left(\frac{1}{2}, \frac{1}{2}; 1; \left(\frac{1-r}{1+r}\right)^2\right) &= \frac{1+r}{2}F\left(\frac{1}{2}, \frac{1}{2}; 1; 1-r^2\right). \end{aligned} \quad (1.3)$$

In 1992, a natural problem related to Landen identities was proposed by Anderson et al. in [7, p. 79]:

PROBLEM 1.1. Find an analog of Landen's transformation formulas in (1.2) for the zero-balanced hypergeometric function $F(a, b; a+b; x)(a, b > 0)$.

Later, Qiu and Vuorinen [35, Theorem 1.2] claimed that the function

$$s(r) = (1 + \sqrt{r})F(a, b; a + b; r) - F\left(a, b; a + b; \frac{4\sqrt{r}}{(1 + \sqrt{r})^2}\right)$$

is increasing in $r \in (0, 1)$ for $(a, b) \in \{a, b > 0 | a + b \leq 1\}$, and derived the Landen inequality

$$F\left(a, b; a + b; \frac{4r}{(1+r)^2}\right) \leq (1+r)F(a, b; a + b; r^2) \quad (1.4)$$

for each $r \in (0, 1)$ and $(a, b) \in \{a, b > 0 | a + b \leq 1\}$. Unfortunately, there is an error in the proof of the monotonicity of $s(r)$. Simić and Vuorinen [39, Lemma 3.2] gave a correct proof, and showed that $s(r)$ is also increasing on $(0, 1)$ for $(a, b) \in \{a, b > 0 | a + b > 1, ab \leq 1/4\}$, and decreasing for $(a, b) \in \{a, b > 0 | 1/a + 1/b \leq 4\}$. Besides, for each $r \in (0, 1)$, they found the maximal regions of ab plane in the first quadrant where equalities (1.2) become (1.4) or its reversed inequality.

Let \mathbf{B}^2 be the unit disk in the plane and $\mu(r)$ the modulus of the plane Grötzsch ring $\mathbf{B}^2 \setminus [0, r]$ for $r \in (0, 1)$. Then $\mu(r)$, called the Grötzsch ring function, has the following explicit expression in terms of complete elliptic integrals (see [29, p. 60])

$$\mu(r) = \frac{\pi}{2} \frac{\mathcal{K}(r')}{\mathcal{K}(r)}, \quad r \in (0, 1). \quad (1.5)$$

By (1.2) and (1.5), it is easy to see that

$$\mu(r) = 2\mu\left(\frac{2\sqrt{r}}{1+r}\right), \quad \mu(r) = \frac{1}{2}\mu\left(\frac{1-r'}{1+r'}\right). \quad (1.6)$$

The Grötzsch ring function plays a very important role in quasi-conformal theory and quasi-regular theory (see [8, 42]). Moreover, using $\mu(r)$, the classical modular equation of signature 2 and degree p ($p > 1$) in number theory also can be rewritten as (see [43])

$$\mu(s) = p\mu(r), \quad 0 < r < 1.$$

Because of its importance, numerous properties and inequalities of $\mu(r)$ have been obtained by many authors in the recent years (see [8, 10, 34]).

On the other hand, in the 1990's, Anderson, Vuorinen, Vamanamurthy and Qiu et al. began to study a natural generalization of $\mu(r)$ defined by

$$\mu_a(r) = \frac{\pi}{2\sin(\pi a)} \frac{F(a, 1-a; 1; 1-r^2)}{F(a, 1-a; 1; r^2)}, \quad r \in (0, 1)$$

in the theory of Ramanujan generalized modular equation, and a lot of beautiful properties and inequalities of $\mu(r)$ have been extended to the function $\mu_a(r)$ [12, 34, 36, 44, 46, 71]. For instance, Qiu and Vuorinen [34, 36] derived the analogs of the duplication formula and logarithmic inequalities satisfied by $\mu(r)$ for $\mu_a(r)$. The infinite-product representations for $\mu_{1/3}(r)$ and $\mu_{1/4}(r)$ which involves only r are established in [54, 56].

The objective of this paper is twofold. The first one is to study a class of hypergeometric function $F(a, b; (a+b+1)/2; x)$ with $a, b > 0$, and prove a generalization of Landen identities for it in the form of inequalities analogously. The other one is to introduce the generalized Grötzsch ring function with two parameters defined by

$$\mu_{a,b}(r) = \frac{B(a, b)}{2} \frac{F(a, b; (a+b+1)/2; 1-r^2)}{F(a, b; (a+b+1)/2; r^2)}, \quad r \in (0, 1), \quad (1.7)$$

and find the analogs of identities (1.6) for $\mu_{a,b}(r)$.

According to the definition of $\mu_{a,b}(r)$ in (1.7), it follows that $\mu_a(r) = \mu_{a,1-a}(r)$, $\mu(r) = \mu_{1/2}(r) = \mu_{1/2,1/2}(r)$, and the limiting values of $\mu_{a,b}(r)$ at 0 and 1 are

$$\begin{aligned} \mu_{a,b}(0^+) &= \lim_{r \rightarrow 0^+} \frac{B(a, b)}{2} F\left(a, b; \frac{a+b+1}{2}; 1-r^2\right) \\ &= \begin{cases} \frac{B(a, b)}{2} H(a, b), & a+b < 1; \\ +\infty, & a+b \geq 1, \end{cases} \end{aligned} \quad (1.8)$$

$$\mu_{a,b}(1^-) = \lim_{r \rightarrow 1^-} \frac{B(a, b)}{2F(a, b; \frac{a+b+1}{2}; r^2)} = \begin{cases} \frac{B(a, b)}{2H(a, b)}, & a+b < 1; \\ 0, & a+b \geq 1, \end{cases} \quad (1.9)$$

respectively. Here and in what follows

$$H(a, b) = \frac{B(\frac{a+b+1}{2}, \frac{1-a-b}{2})}{B(\frac{1+b-a}{2}, \frac{1+a-b}{2})}.$$

Moreover, due to [9, Lemmas 4.5, 4.6 and 4.8], $\mu_{a,b}(r)$ satisfies the following derivative formula

$$\frac{d\mu_{a,b}(r)}{dr} = -\frac{\Gamma(\frac{a+b+1}{2})^2}{\Gamma(a+b)} \frac{1}{r^{a+b} r^{a+b+1} F(a, b; (a+b+1)/2; r^2)^2}. \quad (1.10)$$

2. Main results

For convenience, we first list some regions in $\{(a, b) \in \mathbb{R}^2 | a > 0, b > 0\}$ (see Figure 1):

$$D_1 = \{(a, b) | a, b > 0, a+b \leq 1\},$$

$$D_2 = \left\{ (a, b) | a, b > 0, a+b > 1, ab + \frac{a+b}{2} - \frac{3}{4} < 0 \right\},$$

$$D_3 = \left\{ (a, b) | a, b > 0, ab + \frac{a+b}{2} - \frac{3}{4} \geq 0, ab - \frac{a+b+1}{8} < 0 \right\},$$

$$D_4 = \left\{ (a, b) | a, b > 0, ab - \frac{a+b+1}{8} \geq 0 \right\},$$




Figure 1: The regions D_1 , D_2 , D_3 , D_4 , D_{21} , D_{22} , D_{31} , and D_{32} , where $L : a+b=1$, $l_1 : ab+3(a+b)/2-7/4=0$, $l_2 : ab+(a+b)/2-3/4=0$, $l_3 : ab+7(a+b)/16-11/16=0$, $l_4 : ab-(a+b+1)/8=0$.

$$\begin{aligned} D_{21} &= \left\{ (a,b) | a,b > 0, a+b > 1, ab + \frac{3}{2}(a+b) - \frac{7}{4} < 0 \right\}, \\ D_{22} &= \left\{ (a,b) | a,b > 0, ab + \frac{3}{2}(a+b) - \frac{7}{4} \geq 0, ab + \frac{a+b}{2} - \frac{3}{4} < 0 \right\}, \\ D_{31} &= \left\{ (a,b) | a,b > 0, ab + \frac{a+b}{2} - \frac{3}{4} \geq 0, ab + \frac{7}{16}(a+b) - \frac{11}{16} < 0 \right\}, \\ D_{32} &= \left\{ (a,b) | a,b > 0, ab + \frac{7}{16}(a+b) - \frac{11}{16} \geq 0, ab - \frac{a+b+1}{8} < 0 \right\}. \end{aligned}$$

Clearly, $D_1 \cup D_2 \cup D_3 \cup D_4 = \{(a,b) \in \mathbb{R}^2 | a > 0, b > 0\}$, $D_{21} \subset D_2$, $D_{22} \subset D_2$, $D_{31} \subset D_3$ and $D_{32} \subset D_3$.

THEOREM 2.1. Let $a, b \in (0, \infty)$. Then the inequality

$$F\left(a, b; \frac{a+b+1}{2}; \frac{4r}{(1+r)^2}\right) \leq (1+r)F\left(a, b; \frac{a+b+1}{2}; r^2\right) \quad (2.1)$$

holds for all $r \in (0, 1)$ if and only if $(a, b) \in D_1$, and the reversed inequality

$$F\left(a, b; \frac{a+b+1}{2}; \frac{4r}{(1+r)^2}\right) \geq (1+r)F\left(a, b; \frac{a+b+1}{2}; r^2\right) \quad (2.2)$$

takes place for each $r \in (0, 1)$ if and only if $(a, b) \in D_4$, with equality in each instance if and only if $(a, b) = (1/2, 1/2)$.

In the remaining region $(a, b) \in D_2 \cup D_3$, neither of the above inequalities holds for each $r \in (0, 1)$.

THEOREM 2.2. For $a, b > 0$ with $(a, b) \neq (1/2, 1/2)$, define the function f on $(0, 1)$ by

$$f(r) = (1 + \sqrt{r})F\left(a, b; \frac{a+b+1}{2}; r\right) - F\left(a, b; \frac{a+b+1}{2}; \frac{4\sqrt{r}}{(1+\sqrt{r})^2}\right).$$

Then the following statements are true:

- (1) $f(r)$ is strictly increasing from $(0, 1)$ onto $(0, [R(a, b) - \log 16]/B(a, b))$ when $(a, b) \in L = \{(a, b)|a, b > 0, a+b = 1\}$;
- (2) $f(r)$ is strictly increasing from $(0, 1)$ onto $(0, H(a, b))$ when $(a, b) \in D_1 \setminus L$;
- (3) $f(r)$ is strictly decreasing from $(0, 1)$ onto $(-\infty, 0)$ when $(a, b) \in D_4$.

Moreover, for $(a, b) \in D_1 \setminus L$, the inequality

$$\begin{aligned} F\left(a, b; \frac{a+b+1}{2}; \frac{4r}{(1+r)^2}\right) &\leq (1+r)F\left(a, b; \frac{a+b+1}{2}; r^2\right) \\ &\leq F\left(a, b; \frac{a+b+1}{2}; \frac{4r}{(1+r)^2}\right) + H(a, b) \end{aligned} \quad (2.3)$$

holds for all $r \in (0, 1)$, and for $(a, b) \in L$, the following inequality is valid for each $r \in (0, 1)$,

$$\begin{aligned} F\left(a, b; \frac{a+b+1}{2}; \frac{4r}{(1+r)^2}\right) &\leq (1+r)F\left(a, b; \frac{a+b+1}{2}; r^2\right) \\ &\leq F\left(a, b; \frac{a+b+1}{2}; \frac{4r}{(1+r)^2}\right) + \frac{R(a, b) - \log 16}{B(a, b)}. \end{aligned} \quad (2.4)$$

THEOREM 2.3. For $(a, b) \in \{(a, b)|a, b > 0, ab \geq a+b - 10/9, a+b \geq 2\}$, define the function g on $(0, 1)$ by

$$g(r) = 2\mu_{a,b}\left(\frac{2\sqrt{r}}{1+r}\right) - \mu_{a,b}(r).$$

Then g is strictly increasing from $(0, 1)$ onto $(-\infty, 0)$. In particular, the inequality

$$2\mu_{a,b}\left(\frac{2\sqrt{r}}{1+r}\right) < \mu_{a,b}(r) \quad (2.5)$$

holds for each $r \in (0, 1)$ with $(a, b) \in \{(a, b)|a, b > 0, ab \geq a+b - 10/9, a+b \geq 2\}$.

REMARK 2.4. If $a+b=1$, then our Theorems 2.1 and 2.2 lead to Theorem 2.1 and Lemma 3.2 in [39], respectively.

3. Proofs of Theorems 2.1 and 2.2

Thought out this section, we let

$$F(x) = F\left(a, b; \frac{a+b+1}{2}; x\right), \quad G(x) = F\left(a+1, b+1; \frac{a+b+3}{2}; x\right) \quad (3.1)$$

with $a, b > 0$ and $(a, b) \neq (1/2, 1/2)$, and

$$F^*(x) = F\left(\frac{1}{2}, \frac{1}{2}; 1; x\right), \quad G^*(x) = F\left(\frac{3}{2}, \frac{3}{2}; 2; x\right). \quad (3.2)$$

LEMMA 3.1. ([63, Theorem 2.1]) Suppose that the power series $f(x) = \sum_{n=0}^{\infty} a_n x^n$ and $g(x) = \sum_{n=0}^{\infty} b_n x^n$ have the radius of convergence $r > 0$ with $b_n > 0$ for all $n \in \{0, 1, 2, \dots\}$. Let $h(x) = f(x)/g(x)$ and $H_{f,g} = (f'/g')g - f$, then the following statements are true:

- (1) If the non-constant sequence $\{a_n/b_n\}_{n=0}^{\infty}$ is increasing (decreasing), then $h(x)$ is strictly increasing (decreasing) on $(0, r)$;
- (2) If the non-constant sequence $\{a_n/b_n\}$ is increasing (decreasing) for $0 < n \leq n_0$ and decreasing (increasing) for $n > n_0$, then the function h is strictly increasing (decreasing) on $(0, r)$ if and only if $H_{f,g}(r^-) \geq (\leq) 0$. While if $H_{f,g}(r^-) < (>) 0$, then there exists $x_0 \in (0, r)$ such that $h(x)$ is strictly increasing (decreasing) on $(0, x_0)$ and strictly decreasing (increasing) on (x_0, r) .

LEMMA 3.2. (1) The function $\phi(x) = F(x)/F^*(x)$ is strictly decreasing on $(0, 1)$ if $(a, b) \in D_1$ and strictly increasing on $(0, 1)$ if $(a, b) \in D_4$. Moreover, if $(a, b) \in D_2 \cup D_3$, then there exists $x_0 \in (0, 1)$ such that $\phi(x)$ is strictly decreasing on $(0, x_0)$ and strictly increasing on $(x_0, 1)$;

(2) The function $\varphi(x) = G(x)/G^*(x)$ is strictly decreasing on $(0, 1)$ if $(a, b) \in D_1$ and strictly increasing on $(0, 1)$ if $(a, b) \in D_4 \cup D_{32}$. In the remaining region, namely $x \in D_2 \cup D_{31}$, $\varphi(x)$ is piecewise monotone on $(0, 1)$.

Proof. For part (1), let

$$A_n = \frac{(a, n)(b, n)}{\left(\frac{a+b+1}{2}, n\right) n!}, \quad A_n^* = \frac{(1/2, n)(1/2, n)}{(1, n)(n)!},$$

then

$$\phi(x) = \frac{F(x)}{F^*(x)} = \frac{\sum_{n=0}^{\infty} A_n x^n}{\sum_{n=0}^{\infty} A_n^* x^n}. \quad (3.3)$$

Simple computations show that the monotonicity of $\{A_n/A_n^*\}$ depends on the sign of

$$H_n = \left(\frac{a+b-1}{2} \right) n^2 + \left(ab + \frac{a+b}{2} - \frac{3}{4} \right) n + ab - \frac{a+b+1}{8}. \quad (3.4)$$

We divide the proof into three cases.

Case 1 $(a, b) \in D_1$. Then $H_n < 0$ for $n = 0, 1, 2, \dots$ follows from (3.4). Therefore, (3.3) and Lemma 3.1(1) lead to that $\phi(x)$ is strictly decreasing on $(0, 1)$.

Case 2 $(a, b) \in D_4$. Then (3.4) implies $H_n > 0$ for $n = 0, 1, 2, \dots$, and $\phi(x)$ is strictly increasing on $(0, 1)$ by (3.3) and Lemma 3.1(1).

Case 3 $(a, b) \in D_2 \cup D_3$. Then from (3.4) we conclude that the sequence $\{A_n/A_n^*\}$ is decreasing for $n \in \{1, 2, \dots, n_0\}$ and increasing for $n \geq n_0$, for some integer n_0 . Moreover, using (1.1) and the derivative formula of hypergeometric function

$$\frac{dF(a, b; c; x)}{dx} = \frac{ab}{c} F(a+1, b+1; c+1; x),$$

we get

$$\begin{aligned} H_{F,F^*}(x) &= \frac{8ab}{a+b+1} (1-x) \frac{F(a+1, b+1; (a+b+3)/2; x)}{F(1/2, 1/2; 2; x)} F\left(\frac{1}{2}, \frac{1}{2}; 1; x\right) \\ &\quad - F\left(a, b; \frac{a+b+1}{2}; x\right) \\ &= \frac{8ab}{a+b+1} (1-x) \frac{(1-x)^{-(a+b+1)/2} F\left(\frac{b-a+1}{2}, \frac{a-b+1}{2}; \frac{a+b+3}{2}; x\right)}{F(1/2, 1/2; 2; x)} F\left(\frac{1}{2}, \frac{1}{2}; 1; x\right) \\ &\quad - (1-x)^{\frac{1-a-b}{2}} F\left(\frac{b-a+1}{2}, \frac{a-b+1}{2}; \frac{a+b+1}{2}; x\right) \\ &= (1-x)^{\frac{1-a-b}{2}} \left[\frac{8ab}{a+b+1} \frac{F\left(\frac{b-a+1}{2}, \frac{a-b+1}{2}; \frac{a+b+3}{2}; x\right)}{F(1/2, 1/2; 2; x)} F\left(\frac{1}{2}, \frac{1}{2}; 1; x\right) \right. \\ &\quad \left. - F\left(\frac{b-a+1}{2}, \frac{a-b+1}{2}; \frac{a+b+1}{2}; x\right) \right], \\ &\lim_{x \rightarrow 1^-} H_{F,F^*}(x) = +\infty. \end{aligned} \quad (3.5)$$

Combining (3.5), (3.3) and Lemma 3.1(2) together with (1.1), we have that there exists $x_0 \in (0, 1)$ such that $\phi(x)$ is strictly decreasing on $(0, x_0)$ and strictly increasing on $(x_0, 1)$.

For part (2), let

$$B_n = \frac{(a+1, n)(b+1, n)}{\left(\frac{a+b+3}{2}, n\right) n!}, \quad B_n^* = \frac{(3/2, n)(3/2, n)}{(2, n)(n)!},$$

then

$$\varphi(x) = \frac{G(x)}{G^*(x)} = \frac{\sum_{n=0}^{\infty} B_n x^n}{\sum_{n=0}^{\infty} B_n^* x^n}. \quad (3.6)$$

Note that the monotonicity of $\{B_n/B_n^*\}$ depends on the sign of

$$H_n^* = \left(\frac{a+b-1}{2} \right) n^2 + \left[ab + \frac{3(a+b)}{2} - \frac{7}{4} \right] n + 2ab + \frac{7(a+b)}{8} - \frac{11}{8}. \quad (3.7)$$

Moreover, for $(a, b) \in D_2 \cup D_{31}^*$, we have

$$\begin{aligned} H_{G,G^*}(x) &= \frac{2(a+1)(b+1)}{a+b+3} \frac{F(a+2, b+2; \frac{a+b+5}{2}; x)}{9F(\frac{5}{2}, \frac{5}{2}; 3; x)/8} F\left(\frac{3}{2}, \frac{3}{2}; 2; x\right) \\ &\quad - F\left(a+1, b+1; \frac{a+b+3}{2}; x\right) \\ &= \frac{16(a+1)(b+1)}{9(a+b+3)} (1-x)^{-\frac{a+b+1}{2}} \frac{F\left(\frac{b-a+1}{2}, \frac{a-b+1}{2}; \frac{a+b+5}{2}; x\right)}{F(\frac{1}{2}, \frac{1}{2}; 3; x)} F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) \\ &\quad - (1-x)^{-\frac{a+b+1}{2}} F\left(\frac{b-a+1}{2}, \frac{a-b+1}{2}; \frac{a+b+3}{2}; x\right) \\ &= (1-x)^{-\frac{a+b+1}{2}} \left[\frac{16(a+1)(b+1)}{9(a+b+3)} \frac{F\left(\frac{b-a+1}{2}, \frac{a-b+1}{2}; \frac{a+b+5}{2}; x\right)}{F(\frac{1}{2}, \frac{1}{2}; 3; x)} F\left(\frac{1}{2}, \frac{1}{2}; 2; x\right) \right. \\ &\quad \left. - F\left(\frac{b-a+1}{2}, \frac{a-b+1}{2}; \frac{a+b+3}{2}; x\right) \right], \end{aligned}$$

$$\begin{aligned} &\lim_{x \rightarrow 1^-} (1-x)^{\frac{a+b+1}{2}} H_{G,G^*}(x) \\ &= \frac{16(a+1)(b+1)}{9(a+b+3)} \frac{\Gamma(\frac{a+b+5}{2}) \Gamma(\frac{a+b+3}{2})}{\Gamma(a+2)\Gamma(b+2)} \frac{\Gamma(2)\Gamma(1)}{\Gamma(\frac{3}{2})^2} \frac{\Gamma(\frac{5}{2})^2}{\Gamma(3)\Gamma(2)} - \frac{\Gamma(\frac{a+b+3}{2}) \Gamma(\frac{a+b+1}{2})}{\Gamma(a+1)\Gamma(b+1)} \\ &= \left(\frac{a+b-1}{2} \right) \frac{\Gamma(\frac{a+b+3}{2}) \Gamma(\frac{a+b+1}{2})}{\Gamma(a+1)\Gamma(b+1)} > 0, \end{aligned}$$

and thereby,

$$\lim_{x \rightarrow 1^-} H_{G,G^*}(x) = +\infty. \quad (3.8)$$

It follows from (3.6)-(3.8) together with the similar arguments in part (1) that the assertions of part (2) hold true. Here we omit the details. \square

Proof of Theorem 2.1. Since $y(r) = 4r/(1+r)^2 > r^2$ for $0 < r < 1$, Lemma 3.2(1) shows that $\phi(r^2) > \phi(y(r))$ for $(a, b) \in D_1$ and $\phi(r^2) < \phi(y(r))$ for $(a, b) \in D_4$. Therefore, it follows from (1.2) that

$$\frac{F(r^2)}{F^*(r^2)} > \frac{F(y(r))}{F^*(y(r))}, \quad F(y(r)) < \frac{F^*(y(r))}{F^*(r^2)} F(r^2) = (1+r)F(r^2)$$

for $(a, b) \in D_1$, which implies (2.1).

Inequality (2.2) can be derived analogously. The remaining conclusions easily follow from Lemma 3.2(1). \square

Proof of Theorem 2.2. Clearly $f(0^+) = 0$. From (1.1) we get

$$f(1^-) = \begin{cases} H(a,b), & a+b < 1, \\ \frac{R(a,b)-\log 16}{B(a,b)}, & a+b = 1, \\ -\infty, & a+b > 1. \end{cases} \quad (3.9)$$

Let $x = x(r) = 4\sqrt{r}/(1+\sqrt{r})^2$. Then

$$1-x = \left(\frac{1-\sqrt{r}}{1+\sqrt{r}} \right)^2, \quad \frac{dx}{dr} = \frac{2(1-\sqrt{r})}{\sqrt{r}(1+\sqrt{r})^3}. \quad (3.10)$$

Differentiating f yields

$$\begin{aligned} 2\sqrt{r}(1-\sqrt{r})f'(r) &= (1-\sqrt{r})F(r) + 2\sqrt{r}(1-r)\frac{2ab}{a+b+1}G(r) \\ &\quad - \frac{2ab}{a+b+1}\frac{4(1-x)}{1+\sqrt{r}}G(x). \end{aligned} \quad (3.11)$$

On the other hand, differentiating Landen identity (1.2) with replacing r by \sqrt{r} , we get

$$\frac{1}{4}\frac{2(1-\sqrt{r})}{\sqrt{r}(1+\sqrt{r})^3}G^*(x) = \frac{1}{2\sqrt{r}}F^*(r) + \frac{1}{4}(1+\sqrt{r})G^*(r),$$

namely,

$$\frac{1-x}{1+\sqrt{r}}\frac{G^*(x)}{G^*(r)} = \frac{(1-\sqrt{r})F^*(r)}{G^*(r)} + \frac{\sqrt{r}}{2}(1-r). \quad (3.12)$$

Since φ defined as in Lemma 3.2(2) is strictly decreasing in $(0, 1)$ on D_1 , we conclude that $\varphi(r) > \varphi(x)$ for all $r \in (0, 1)$, that is

$$G(x) < \frac{G^*(x)}{G^*(r)}G(r). \quad (3.13)$$

Equations (3.11) and (3.12) together with inequality (3.13) lead to

$$\begin{aligned} &2\sqrt{r}(1-\sqrt{r})f'(r) \\ &= (1-\sqrt{r})F(r) + 2\sqrt{r}(1-r)\frac{2ab}{a+b+1}G(r) - \frac{2ab}{a+b+1}\frac{4(1-x)}{1+\sqrt{r}}G(x) \\ &> (1-\sqrt{r})F(r) + 2\sqrt{r}(1-r)\frac{2ab}{a+b+1}G(r) - \frac{2ab}{a+b+1}\frac{4(1-x)}{1+\sqrt{r}}\frac{G^*(x)}{G^*(r)}G(r) \\ &= (1-\sqrt{r})F(r) + 2\sqrt{r}(1-r)\frac{2ab}{a+b+1}G(r) - \frac{8ab}{a+b+1} \left[\frac{(1-\sqrt{r})F^*(r)}{G^*(r)} + \frac{\sqrt{r}}{2}(1-r) \right] G(r) \\ &= (1-\sqrt{r}) \left[F(r) - \frac{8ab}{a+b+1} \frac{F^*(r)}{G^*(r)} G(r) \right] \\ &= (1-\sqrt{r}) \frac{4F(r)^2}{G^*(r)} \left(\frac{F^*(r)}{F(r)} \right)' . \end{aligned} \quad (3.14)$$

Lemma 3.2(1) shows that $r \rightarrow F^*(r)/F(r)$ is strictly increasing in $(0, 1)$ on $(a, b) \in D_1$. Thus by (3.14) one has $f'(r) > 0$ for all $r \in (0, 1)$ when $(a, b) \in D_1$, and f is strictly increasing in $(0, 1)$.

On the other hand, since φ defined as in Lemma 3.2(2) is strictly increasing in $(0, 1)$ on D_4 , we have $\varphi(x) > \varphi(r)$, namely

$$G(x) > \frac{G^*(x)}{G^*(r)} G(r).$$

With the similar argument, one has

$$\sqrt{r}f'(r) < \frac{2F(r)^2}{G^*(r)} \left(\frac{F^*(r)}{F(r)} \right)' < 0,$$

because $F(r)/F^*(r)$ is strictly increasing in $(0, 1)$ on D_4 by Lemma 3.2(1). Hence $f(r)$ is strictly decreasing in $(0, 1)$ on D_4 . It is now obvious that the remaining conclusions hold. \square

4. Proof of Theorem 2.3

LEMMA 4.1. Let $x' = \sqrt{1-x^2}$ for $0 < x < 1$. Then the function

$$f(x) = \frac{(xx')^{(a+b-1)/2} F(a, b; \frac{a+b+1}{2}; x^2)}{F(\frac{1}{2}, \frac{1}{2}; 1; x^2)} \quad (4.1)$$

is strictly increasing on $(0, 1)$ when $(a, b) \in \{(a, b) | a, b > 0, ab \geq a+b-10/9, a+b \geq 2\}$.

Proof. Differentiating f gives

$$\begin{aligned} f'(x) = & \frac{1}{F(\frac{1}{2}, \frac{1}{2}; 1; x^2)^2} \left\{ \left[\frac{a+b-1}{2} (xx')^{\frac{a+b-3}{2}} \left(\frac{1-2x^2}{x'} \right) F\left(a, b; \frac{a+b+1}{2}; x^2\right) \right. \right. \\ & + (xx')^{\frac{a+b-1}{2}} \frac{2ab}{a+b+1} F\left(a+1, b+1; \frac{a+b+3}{2}; x^2\right) \cdot 2x \Big] F\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right) \\ & \left. \left. - (xx')^{\frac{a+b-1}{2}} F\left(a, b; \frac{a+b+1}{2}; x^2\right) \cdot \frac{x}{2} F\left(\frac{3}{2}, \frac{3}{2}; 2; x^2\right) \right\}, \right. \end{aligned}$$

and thereby

$$\begin{aligned} f_1(x) \equiv & x'(xx')^{-\frac{a+b-3}{2}} F\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right)^2 f'(x) \\ = & \frac{a+b-1}{2} (1-2x^2) F\left(a, b; \frac{a+b+1}{2}; x^2\right) F\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right) \end{aligned}$$

$$\begin{aligned}
& + \frac{4ab}{a+b+1} x^2 x'^2 F\left(a+1, b+1; \frac{a+b+3}{2}; x^2\right) F\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right) \\
& - \frac{x^2 x'^2}{2} F\left(a, b; \frac{a+b+1}{2}; x^2\right) F\left(\frac{3}{2}, \frac{3}{2}; 2; x^2\right).
\end{aligned} \tag{4.2}$$

Since $x'^2 F\left(\frac{3}{2}, \frac{3}{2}; 2; x^2\right) = F\left(\frac{1}{2}, \frac{1}{2}; 2; x^2\right) \leq F\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right)$ for $x \in (0, 1)$, it follows from (4.2) that

$$\begin{aligned}
f_1(x) &\geq \frac{a+b-1}{2} (1-2x^2) F\left(a, b; \frac{a+b+1}{2}; x^2\right) F\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right) \\
& + \frac{4ab}{a+b+1} x^2 x'^2 F\left(a+1, b+1; \frac{a+b+3}{2}; x^2\right) F\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right) \\
& - \frac{x^2}{2} F\left(a, b; \frac{a+b+1}{2}; x^2\right) F\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right) \\
& = F\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right) \left[\frac{1}{2} (a+b-1 - (2a+2b-1)x^2) F\left(a, b; \frac{a+b+1}{2}; x^2\right) \right. \\
& \quad \left. + \frac{4ab}{a+b+1} x^2 (1-x^2) F\left(a+1, b+1; \frac{a+b+3}{2}; x^2\right) \right]. \tag{4.3}
\end{aligned}$$

Let $f_2(x) = f_1(x)/F\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right)$, then from (4.3) we get

$$\begin{aligned}
f_2(x) &\geq \frac{a+b-1}{2} \sum_{n=0}^{\infty} \frac{(a,n)(b,n)}{\left(\frac{a+b+1}{2}, n\right)} \frac{x^{2n}}{n!} - (a+b-\frac{1}{2}) \sum_{n=0}^{\infty} \frac{(a,n)(b,n)}{\left(\frac{a+b+1}{2}, n\right)} \frac{x^{2n+2}}{n!} \\
& + \frac{4ab}{a+b+1} \sum_{n=0}^{\infty} \frac{(a+1,n)(b+1,n)}{\left(\frac{a+b+3}{2}, n\right)} \frac{x^{2n+2}}{n!} \\
& - \frac{4ab}{a+b+1} \sum_{n=0}^{\infty} \frac{(a+1,n)(b+1,n)}{\left(\frac{a+b+3}{2}, n\right)} \frac{x^{2n+4}}{n!} \\
& = \frac{a+b-1}{2} + \left[\frac{ab(a+b-1)}{a+b+1} - (a+b-\frac{1}{2}) + \frac{4ab}{a+b+1} \right] x^2 \\
& + \frac{a+b-1}{2} \sum_{n=0}^{\infty} \frac{(a,n+2)(b,n+2)}{\left(\frac{a+b+1}{2}, n+2\right)} \frac{x^{2n+4}}{(n+2)!} \\
& - (a+b-\frac{1}{2}) \sum_{n=0}^{\infty} \frac{(a,n+1)(b,n+1)}{\left(\frac{a+b+1}{2}, n+1\right)} \frac{x^{2n+4}}{(n+1)!} \\
& + 2 \sum_{n=0}^{\infty} \frac{(a,n+2)(b,n+2)}{\left(\frac{a+b+1}{2}, n+2\right)} \frac{x^{2n+4}}{(n+1)!} - 2 \sum_{n=0}^{\infty} \frac{(a,n+1)(b,n+1)}{\left(\frac{a+b+1}{2}, n+1\right)} \frac{x^{2n+4}}{n!} \\
& = \frac{a+b-1}{2} \left(1 - \frac{x^2}{a+b+1}\right) + \frac{ab(a+b-1) - (a-b)^2}{a+b+1} x^2 \\
& + \sum_{n=0}^{\infty} \frac{(a,n+1)(b,n+1)}{\left(\frac{a+b+1}{2}, n+2\right)} \frac{C_n}{(n+2)!} x^{2n+4}, \tag{4.4}
\end{aligned}$$

where

$$\begin{aligned}
 C_n &= \frac{a+b-1}{2}(n+a+1)(n+b+1) - (n+2) \left(a+b - \frac{1}{2} \right) \left(n+1 + \frac{a+b+1}{2} \right) \\
 &\quad + 2(n+a+1)(n+b+1)(n+2) - 2(n+1)(n+2) \left(n+1 + \frac{a+b+1}{2} \right) \\
 &= \left(\frac{a+b-2}{2} \right) n^2 + \frac{1}{4}(a+b+8ab-9)n \\
 &\quad + \frac{1}{2} [ab(a+b+3) - (a-b)^2 - (a+b+2)]. \tag{4.5}
 \end{aligned}$$

If $(a, b) \in \{(a, b) | a, b > 0, ab \geq a+b-10/9, a+b \geq 2\}$, then by elementary computations one has

$$\begin{aligned}
 ab(a+b-1) - (a-b)^2 &\geq \left(a+b - \frac{10}{9} \right) (a+b-1) - (a-b)^2 \\
 &= 4ab - \frac{19(a+b)}{9} + \frac{10}{9} \geq \frac{17(a+b)}{9} - \frac{30}{9} \geq \frac{4}{9}, \\
 a+b+8ab-9 &\geq a+b+8 \left(a+b - \frac{10}{9} \right) - 9 = 9(a+b) - \frac{161}{9} \geq \frac{1}{9}, \\
 ab(a+b+3) - (a-b)^2 - (a+b+2) &\geq \left(a+b - \frac{10}{9} \right) (a+b+3) - (a-b)^2 - (a+b+2) \\
 &= 4ab + \frac{8(a+b)}{9} - \frac{16}{3} \geq 4 \left(a+b - \frac{10}{9} \right) + \frac{8(a+b)}{9} - \frac{16}{3} \\
 &= \frac{44(a+b)}{9} - \frac{88}{9} \geq 0.
 \end{aligned}$$

Thus it follows from (4.4) and (4.5) that $f_2(x) > 0$ for all $x \in (0, 1)$. Therefore by (4.2) and (4.3), f is strictly increasing on $(0, 1)$ when $(a, b) \in \{(a, b) | a, b > 0, ab \geq a+b-10/9, a+b \geq 2\}$. \square

REMARK 4.2. The main purpose of Lemma 4.1 is to find some $a, b > 0$ such that the function f defined as in (4.1) is strictly monotone on $(0, 1)$. In fact, for $a+b < 1$, we have that $f(0^+) = f(1^-) = +\infty$, so that f is not monotone on $(0, 1)$; While for $a+b = 1$, Lemma 3.2(1) shows that f is strictly decreasing on $(0, 1)$. In the remaining region $(a, b) \in \{a+b > 1\}$, note that $f(0^+) = 0$ and $f(1^-) = +\infty$, we find its subregion such that f is strictly increasing on $(0, 1)$ in Lemma 4.1.

COROLLARY 4.3. For $(a, b) \in \{(a, b) | a, b > 0, ab \geq a+b-10/9, a+b \geq 2\}$, let $x = x(r) = 2\sqrt{r}/(1+r)$, then the Landen-type inequality

$$(xx')^{(a+b-1)/2} F \left(a, b; \frac{a+b+1}{2}; x^2 \right) > (1+r)(rr')^{(a+b-1)/2} F \left(a, b; \frac{a+b+1}{2}; r^2 \right) \tag{4.6}$$

holds for all $r \in (0, 1)$.

Proof. Since $x = x(r) = 2\sqrt{r}/(1+r) > r$ for all $r \in (0, 1)$, Lemma 4.1 implies that $f(x) > f(r)$ for $0 < r < 1$, and consequently inequality (4.6) follows. \square

Proof of Theorem 2.3. Note that $g(1^-) = 0$ by (1.9), and

$$\begin{aligned} & \lim_{r \rightarrow 0^+} g(r) \\ &= \lim_{r \rightarrow 0^+} \frac{B(a, b)}{2} \left[2F \left(a, b; \frac{a+b+1}{2}; \left(\frac{1-r}{1+r} \right)^2 \right) - F \left(a, b; \frac{a+b+1}{2}; 1-r^2 \right) \right] \\ &= \lim_{r \rightarrow 0^+} \left\{ B(a, b) \left[1 - \left(\frac{1-r}{1+r} \right)^2 \right]^{(1-a-b)/2} F \left(\frac{-a+b+1}{2}, \frac{a-b+1}{2}; \frac{a+b+1}{2}; \left(\frac{1-r}{1+r} \right)^2 \right) \right. \\ &\quad \left. - \frac{B(a, b)}{2} r^{1-a-b} F \left(\frac{-a+b+1}{2}, \frac{a-b+1}{2}; \frac{a+b+1}{2}; 1-r^2 \right) \right\} \\ &= \frac{B(a, b)}{2} \frac{\Gamma(\frac{a+b+1}{2}) \Gamma(\frac{a+b-1}{2})}{\Gamma(a) \Gamma(b)} \lim_{r \rightarrow 0^+} \left[2 \cdot \left(\frac{2\sqrt{r}}{1+r} \right)^{1-a-b} - r^{1-a-b} \right] = -\infty. \end{aligned}$$

Let $x = 2\sqrt{r}/(1+r)$, then

$$x' = \sqrt{1-x^2} = \frac{1-r}{1+r}, \quad \frac{dx}{dr} = \frac{x'(1+x')^2}{2x}.$$

Differentiating g yields

$$\begin{aligned} g'(r) &= -2 \frac{\Gamma(\frac{a+b+1}{2})^2}{\Gamma(a+b)} \frac{1}{x^{a+b} x'^{a+b+1} F(a, b; \frac{a+b+1}{2}; x^2)^2} \cdot \frac{x'(1+x')^2}{2x} \\ &\quad + \frac{\Gamma(\frac{a+b+1}{2})^2}{\Gamma(a+b)} \cdot \frac{1}{r^{a+b} r'^{a+b+1} F(a, b; \frac{a+b+1}{2}; r^2)^2} \\ &= \frac{\Gamma(\frac{a+b+1}{2})^2}{\Gamma(a+b)} \frac{(1+x')^2}{(1+r)^2 x^{a+b+1} x'^{a+b} F(a, b; \frac{a+b+1}{2}; x^2)^2} \\ &\quad \times \left[\frac{(xx')^{a+b-1} F(a, b; \frac{a+b+1}{2}; x^2)^2}{(rr')^{a+b-1} F(a, b; \frac{a+b+1}{2}; r^2)^2} - (1+r)^2 \right]. \end{aligned} \tag{4.7}$$

Therefore, the monotonicity of g directly follows from (4.6) and (4.7). Inequality (2.5) is clear. \square

REFERENCES

- [1] M. ABRAMOWITZ AND I. A. STEGUN, *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*, Dover, New York, 1965.
- [2] M. ADIL KLAN, Y.-M. CHU, T. U. KHAN AND J. KHAN, *Some new inequalities of Hermite-Hadamard type of s -convex functions with applications*, Open Math. **15** (2017), 1414–1430.
- [3] H. ALZER AND K. RICHARDS, *On the modulus of the Grötzsch ring*, J. Math. Anal. Appl. **432**, 1 (2015), 134–141.
- [4] H. ALZER AND K. RICHARDS, *Inequalities for the ratio of complete elliptic integrals*, Proc. Amer. Math. Soc. **145**, 4 (2017), 1661–1670.
- [5] G. D. ANDERSON, S.-L. QIU AND M. K. VAMANAMURTHY, *Elliptic integral inequalities, with applications*, Constr. Approx. **14**, 2 (1998), 195–207.
- [6] G. D. ANDERSON, S.-L. QIU, M. K. VAMANAMURTHY AND M. VUORINEN, *Generalized elliptic integrals and modular equations*, Pacific J. Math. **192**, 1 (2000), 1–37.
- [7] G. D. ANDERSON, M. K. VAMANAMURTHY AND M. K. VUORINEN, *Hypergeometric functions and elliptic integrals*, in: Current Topics in Analytic Function Theory (H. M. Srivastava and S. Owa, eds.), 48–85, World Scientific Publ. Co., Singapore-London, 1992.
- [8] G. D. ANDERSON, M. K. VAMANAMURTHY AND M. VUORINEN, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.
- [9] G. D. ANDERSON, M. K. VAMANAMURTHY AND M. VUORINEN, *Topics in special functions, In: Papers on Analysis: A volume dedicated to Olli Martio on the occasion of his 60th birthday*, Report Univ. Jyväskylä **83** (2001), 5–26.
- [10] G. D. ANDERSON, M. K. VAMANAMURTHY AND M. VUORINEN, *Topics in special functions II*, Conform. Geom. Dyn. **11** (2007), 250–270.
- [11] R. W. BARNARD, K. PEARCE AND K. C. RICHARDS, *An inequality involving the generalized hypergeometric function and the arc length of an ellipse*, SIAM J. Math. Anal. **31**, 3 (2000), 693–699.
- [12] B. A. BHAYO AND M. VUORINEN, *On generalized complete elliptic integrals and modular functions*, Proc. Edinb. Math. Soc. (2) **55**, 3 (2012), 591–611.
- [13] J. M. BORWEIN AND P. B. BORWEIN, *Pi and AGM*, John Wiley & Sons, New York, 1987.
- [14] Y.-M. CHU, M. ADIL KLAN, S. S. DRAGOMIR, *Inequalities for α -fractional differentiable functions*, J. Inequal. Appl. **2017** (2017), Article 93, 12 pages.
- [15] Y.-M. CHU, S.-L. QIU AND M.-K. WANG, *Sharp inequalities involving the power mean and complete elliptic integral of the first kind*, Rocky Mountain J. Math. **43**, 3 (2013), 1489–1496.
- [16] Y.-M. CHU, Y.-F. QIU AND M.-K. WANG, *Hölder mean inequalities for the complete elliptic integrals*, Integral Transforms Spec. Funct. **23**, 7 (2012), 521–527.
- [17] Y.-M. CHU AND M.-K. WANG, *Optimal Lehmer mean bounds for the Toader mean*, Results Math. **61**, 3–4 (2012), 223–229.
- [18] Y.-M. CHU AND M.-K. WANG, *Inequalities between arithmetic-geometric, Gini, and Toader means*, Abstr. Appl. Anal. **2012** (2012), Article ID 830585, 11 pages.
- [19] Y.-M. CHU, M.-K. WANG, Y.-P. JIANG AND S.-L. QIU, *Concavity of the complete elliptic integrals of the second kind with respect to Hölder means*, J. Math. Anal. Appl. **395**, 2 (2012), 637–642.
- [20] Y.-M. CHU, M.-K. WANG AND X.-Y. MA, *Sharp bounds for Toader mean in terms of contraharmonic mean with applications*, J. Math. Inequal. **7**, 2 (2013), 161–166.
- [21] Y.-M. CHU, M.-K. WANG AND S.-L. QIU, *Optimal combinations bounds of root-square and arithmetic means for Toader mean*, Proc. Indian Acad. Sci. Math. Sci. **122**, 1 (2012), 41–51.
- [22] Y.-M. CHU, M.-K. WANG, S.-L. QIU AND Y.-P. JIANG, *Bounds for complete integrals of the second kind with applications*, Comput. Math. Appl. **63**, 7 (2012), 1177–1184.
- [23] Y.-M. CHU, M.-K. WANG, S.-L. QIU AND Y.-F. QIU, *Sharp generalized Seiffert mean bounds for Toader mean*, Abstr. Appl. Anal. **2011** (2011), Article ID 605259, 8 pages.
- [24] Y.-M. CHU, M.-K. WANG AND Y.-F. QIU, *On Alzer and Qiu's conjecture for complete elliptic integral and inverse hyperbolic tangent function*, Abstr. Appl. Anal. **2011** (2011), Article ID 697547, 7 pages.
- [25] Y.-M. CHU AND W.-F. XIA, *Two sharp inequalities for power mean, geometric mean, and harmonic mean*, J. Inequal. Appl. **2009** (2009), Article ID 741923, 6 pages.
- [26] Y.-M. CHU, X.-M. ZHANG AND G.-D. WANG, *The Schur geometrical convexity of the extended mean values*, J. Convex Anal. **15**, 4 (2008), 707–718.
- [27] A. GASULL, W. G. LI, J. LLIBRE AND Z.-F. ZHANG, *Chebyshev property of complete elliptic integrals and its applications to Abelian integrals*, Pacific. J. Math. **202**, 2 (2002), 341–361.
- [28] D. F. LAWDEN, *Elliptic Functions and Applications*, Springer-Verlag, New York, 1989.

- [29] O. LEHTO AND K. I. VIRTANEN, *Quasiconformal Mappings in the Plane* (2nd ed.), Springer-Verlag, New York-Heidelberg, 1973.
- [30] F. W. J. OLVER, D. W. LOZIER, R. F. BOISVERT AND C. W. CLARK (eds.), *NIST Handbook of Mathematical Functions*, Cambridge Univ. Press, Cambridge, 2010.
- [31] W.-M. QIAN AND Y.-M. CHU, *Sharp bounds for a special quasi-arithmetic mean in terms of arithmetic and geometric means with two parameters*, J. Inequal. Appl. **2017** (2017), Article 274, 10 pages.
- [32] W.-M. QIAN, X.-H. ZHANG AND Y.-M. CHU, *Sharp bounds for the Toader-Qi mean in terms of harmonic and geometric means*, J. Math. Inequal. **11**, 1 (2017), 121–127.
- [33] S.-L. QIU, M. K. VAMANAMURTHY AND M. VUORINEN, *Some inequalities for the Hersch-Pfluger distortion function*, J. Inequal. Appl. **4**, 2 (1999), 115–139.
- [34] S.-L. QIU AND M. VUORINEN, *Infinite products and the normalized quotients of hypergeometric functions*, SIAM J. Math. Anal. **30**, 5 (1999), 1057–1075.
- [35] S.-L. QIU AND M. VUORINEN, *Landen inequalities for hypergeometric functions*, Nagoya Math. J. **154** (1999), 31–56.
- [36] S.-L. QIU AND M. VUORINEN, *Duplication inequalities for the ratios of hypergeometric functions*, Forum Math. **12**, 1 (2000), 109–133.
- [37] J. SÁNDOR, *On certain inequalities for means*, J. Math. Anal. Appl. **189**, 2 (1995), 602–606.
- [38] D. SCHULTZ, *Cubic modular equations in two variables*, Adv. Math. **290** (2016), 329–363.
- [39] S. SIMIĆ AND M. VUORINEN, *Landen inequalities for zero-balanced hypergeometric functions*, Abstr. Appl. Anal. **2012** (2012), Article ID 932061, 11 pages.
- [40] Y.-Q. SONG, P.-G. ZHOU AND Y.-M. CHU, *Inequalities for the Gaussian hypergeometric function*, Sci. China Math. **57**, 11 (2014), 2369–2380.
- [41] M. K. VAMANAMURTHY AND M. VUORINEN, *Inequalities for means*, J. Math. Anal. Appl. **183**, 1 (1994), 155–166.
- [42] M. VUORINEN, *Conformal Geometry and Quasiregular Mappings*, Springer-Velag, Berlin, New York, 1988.
- [43] M. VUORINEN, *Singular values, Ramanujan modular equations, and Landen transformations*, Studia Math. **121**, 3 (1996), 221–230.
- [44] G.-D. WANG, X.-H. ZHANG AND Y.-M. CHU, *Inequalities for the generalized elliptic integrals and modular functions*, J. Math. Anal. Appl. **331**, 2 (2007), 1275–1283.
- [45] G.-D. WANG, X.-H. ZHANG AND Y.-M. CHU, *A power mean inequality involving the complete elliptic integrals*, Rocky Mountain J. Math. **44**, 5 (2014), 1661–1667.
- [46] G.-D. WANG, X.-H. ZHANG AND Y.-P. JIANG, *Concavity with respect to Hölder means involving the generalized Grötzsch function*, J. Math. Anal. Appl. **379**, 1 (2011), 200–204.
- [47] M.-K. WANG AND Y.-M. CHU, *Asymptotical bounds for complete elliptic integrals of the second kind*, J. Math. Anal. Appl. **402**, 1 (2013), 119–126.
- [48] M.-K. WANG AND Y.-M. CHU, *Refinements of transformation inequalities for zero-balanced hypergeometric functions*, Acta Math. Sci. **37B**, 3 (2017), 607–622.
- [49] M.-K. WANG, Y.-M. CHU AND Y.-P. JIANG, *Ramanujan's cubic transformation inequalities for zero-balanced hypergeometric functions*, Rocky Mountain J. Math. **46**, 2 (2016), 679–691.
- [50] M.-K. WANG, Y.-M. CHU AND S.-L. QIU, *Some monotonicity properties of generalized elliptic integrals with applications*, Math. Inequal. Appl. **16**, 3 (2013), 671–677.
- [51] M.-K. WANG, Y.-M. CHU AND S.-L. QIU, *Sharp bounds for generalized elliptic integrals of the first kind*, J. Math. Anal. Appl. **429**, 2 (2015), 744–757.
- [52] M.-K. WANG, Y.-M. CHU, S.-L. QIU AND Y.-P. JIANG, *Convexity of the complete elliptic integrals of the first kind with respect to Hölder means*, J. Math. Anal. Appl. **388**, 2 (2012), 1141–1146.
- [53] M.-K. WANG, Y.-M. CHU, Y.-F. QIU, S.-L. QIU, *An optimal power mean inequality for the complete elliptic integrals*, Appl. Math. Lett. **24**, 6 (2011), 887–890.
- [54] M.-K. WANG, Y.-M. CHU AND Y.-Q. SONG, *Ramanujan's cubic transformation and generalized modular equation*, Sci. China. Math. **58**, 11 (2015), 2387–2404.
- [55] M.-K. WANG, Y.-M. CHU AND Y.-Q. SONG, *Asymptotical formulas for Gaussian and generalized hypergeometric functions*, Appl. Math. Comput. **276** (2016), 44–60.
- [56] M.-K. WANG, Y. M. LI AND Y.-M. CHU, *Inequalities and infinite product formula for Ramanujan generalized modular equation function*, Ramanujan J. doi: 10.1007/s11139-0176-9888-3, 12 pages.
- [57] M.-K. WANG, S.-L. QIU, Y.-M. CHU AND Y.-P. JIANG, *Generalized Hersch-Pfluger distortion function and complete elliptic integrals*, J. Math. Anal. Appl. **385**, 1 (2012), 221–229.
- [58] M.-K. WANG, Y.-F. QIU AND Y.-M. CHU, *Sharp bounds for Seiffert means in terms of Lehmer means*, J. Math. Inequal. **4**, 4 (2010), 581–586.

- [59] E. T. WHITTAKER AND G. N. WATSON, *A Course of Modern Analysis*, Cambridge Univ. Press, New York, 1962.
- [60] ZH.-H. YANG AND Y.-M. CHU, *Asymptotic formulas for gamma function with applications*, Appl. Math. Comput. **270** (2015), 665–680.
- [61] ZH.-H. YANG AND Y.-M. CHU, *On approximating the modified Bessel function of the second kind*, J. Inequal. Appl. **2017** (2017), Article 41, 8 pages.
- [62] ZH.-H. YANG AND Y.-M. CHU, *A monotonicity property involving the generalized elliptic integral of the first kind*, Math. Inequal. Appl. **20**, 3 (2017), 729–735.
- [63] ZH.-H. YANG, Y.-M. CHU AND M.-K. WANG, *Monotonicity criterion for the quotient of power series with applications*, J. Math. Anal. Appl. **428**, 1 (2015), 587–604.
- [64] ZH.-H. YANG, Y.-M. CHU AND W. ZHANG, *Monotonicity of the ratio for the complete elliptic integral and Stolarsky mean*, J. Inequal. Appl. **2016** (2016), Article 176, 10 pages.
- [65] ZH.-H. YANG, Y.-M. CHU AND X.-H. ZHANG, *Sharp Stolarsky mean bounds for the complete elliptic integral of the second kind*, J. Nonlinear Sci. Appl. **10**, 3 (2017), 929–936.
- [66] ZH.-H. YANG, Y.-M. CHU AND X.-H. ZHANG, *Sharp bounds for psi function*, Appl. Math. Comput. **268** (2015), 1055–1063.
- [67] ZH.-H. YANG, Y.-M. CHU AND X.-H. ZHANG, *Necessary and sufficient conditions for functions involving the psi function to be completely monotonic*, J. Inequal. Appl. **2015** (2015), Article 157, 17 pages.
- [68] ZH.-H. YANG, W.-M. QIAN AND Y.-M. CHU, *On rational bounds for the gamma function*, J. Inequal. Appl. **2017** (2017), Article 210, 17 pages.
- [69] ZH.-H. YANG, W. ZHANG AND Y.-M. CHU, *Sharp Gautschi inequality for parameter $0 < p < 1$ with applications*, Math. Inequal. Appl. **20**, 4 (2017), 1107–1120.
- [70] X.-M. ZHANG AND Y.-M. CHU, *A double inequality for gamma function*, J. Inequal. Appl. **2009** (2009), Article ID 503782, 7 pages.
- [71] X.-H. ZHANG, G.-D. WANG AND Y.-M. CHU, *Some inequalities for the generalized Grötzsch function*, Proc. Edinb. Math. Soc. (2) **51**, 1 (2008), 265–272.
- [72] X.-H. ZHANG, G.-D. WANG AND Y.-M. CHU, *Convexity with respect to Hölder mean involving zero-balanced hypergeometric functions*, J. Math. Anal. Appl. **353**, 1 (2009), 256–259.
- [73] X.-H. ZHANG, G.-D. WANG AND Y.-M. CHU, *Remarks on generalized elliptic integrals*, Proc. Roy. Soc. Edinburgh Sect. A **139**, 2 (2009), 417–426.
- [74] T.-H. ZHAO AND Y.-M. CHU, *A class of logarithmically completely monotonic functions associated with a gamma function*, J. Inequal. Appl. **2010** (2010) Article ID 392431, 11 pages.
- [75] T.-H. ZHAO, Y.-M. CHU AND Y.-P. JIANG, *Monotonic and logarithmically convex properties of a function involving gamma functions*, J. Inequal. Appl. **2009** (2009), Article ID 728612, 13 pages.
- [76] T.-H. ZHAO, Y.-M. CHU AND H. WANG, *Logarithmically complete monotonicity properties relating to the gamma function*, Abstr. Appl. Anal. **2011** (2011), Article ID 896483, 13 pages.
- [77] T.-H. ZHAO, ZH.-H. YANG AND Y.-M. CHU, *Monotonicity properties of a function involving the psi function with applications*, J. Inequal. Appl. **2015** (2015), Article 193, 10 pages.

(Received September 20, 2017)

Miao-Kun Wang

Department of Mathematics

Huzhou University

Huzhou 313000, P. R. China

e-mail: wmk000@126.com; wangmiaokun@zjhu.edu.cn

Yu-Ming Chu

Department of Mathematics

Huzhou University

Huzhou 313000, P. R. China

e-mail: chuyuming2005@126.com