

## EXTENSIONS OF FEFFERMAN–STEIN MAXIMAL INEQUALITIES

WENHUA GAO AND GUOEN HU

*Abstract.* Let  $\beta_1, \dots, \beta_m \in [0, \infty)$  and  $\mathcal{M}_{L(\log L)^\beta}$  be the maximal operator defined by

$$\mathcal{M}_{L(\log L)^\beta}(f_1, \dots, f_m)(x) = \sup_{Q \ni x} \prod_{j=1}^m \|f_j\|_{L(\log L)^{\beta_j}, Q}.$$

In this paper, we establish the weighted bounds in terms of the  $A_p(\mathbb{R}^{mn})$  constant for  $\mathcal{M}_{L(\log L)^\beta}$  from  $L^{p_1}(I^{q_1}; \mathbb{R}^n, w_1) \times \dots \times L^{p_m}(I^{q_m}; \mathbb{R}^n, w_m)$  to  $L^p(I^q; \mathbb{R}^n, v_{\vec{w}})$ , where  $p_1, \dots, p_m, q_1, \dots, q_m \in (1, \infty)$ ,  $1/p = 1/p_1 + \dots + 1/p_m$ ,  $1/q = 1/q_1 + \dots + 1/q_m$  and  $\vec{w} = (w_1, \dots, w_m)$  a multiple  $A_p$  weights. A weak type endpoint inequality for vector-valued operator  $\mathcal{M}_{L(\log L)^\beta}$  is also given.

*Mathematics subject classification (2010):* 42B15.

*Keywords and phrases:* Maximal inequality, weighted bound, dyadic grid, Lerner's formula, sparse operator.

## REFERENCES

- [1] K. F. ANDERSON AND R. T. JOHN, *Weighted inequality for vector-valued maximal functions and singular integrals*, Studia Math. **69** (1980), 19–31.
- [2] T. A. BUI, J. CONDE-ALONSO, X. T. DUONG AND M. HORMOZI, *A note on weighted bounds for singular integral operators with nonsmooth kernels*, Studia Math. **236** (2017), 245–269.
- [3] D. CRUZ-URIBE, SFO, J. MARTELL AND C. PÉREZ, *Sharp weighted estimates for classical operators*, Adv. Math. **229** (2012), 408–441.
- [4] W. DAMIÁN, A. K. LERNER AND C. PÉREZ, *Sharp weighted bounds for multilinear maximal functions and Calderón-Zygmund operators*, J. Fourier Anal. Appl. **21** (2012), 751–765.
- [5] C. FEFFERMAN AND E. M. STEIN, *Some maximal inequalities*, Amer. J. Math. **93** (1971), 107–115.
- [6] L. GRAFAKOS, *Modern Fourier analysis*, GTM250, 2nd Edition, Springer, New York, 2008.
- [7] L. GRAFAKOS AND J. M. MARTELL, *Extrapolation of weighted norm inequalities for multivariable operators and applications*, J. Geom. Anal. **14** (2004), 19–46.
- [8] G. HU, *Weighted vector-valued estimates for a non-standard Calderón-Zygmund operator*, Nonlinear Anal. **165** (2017), 143–162.
- [9] T. HYTÖNEN, *The  $A_2$  theorem: Remarks and complements*, Contemp. Math. **612**, Amer. Math. Soc. Providence, RI, 2014, 91–106.
- [10] T. HYTÖNEN, M. T. LACEY AND C. PÉREZ, *Sharp weighted bounds for the  $q$ -variation of singular integrals*, Bull. Lond. Math. Soc. **45** (2013), 529–540.
- [11] T. HYTÖNEN AND C. PÉREZ, *Sharp weighted bounds involving  $A_\infty$* , Anal. PDE. **6** (2013), 777–818.
- [12] A. K. LERNER, *A pointwise estimate for the local sharp maximal function with applications to singular integrals*, Bull. London. Math. Soc. **42** (2010), 843–856.
- [13] A. LERNER, S. OMBROSI, C. PÉREZ, R. H. TORRES AND R. TRUJILLO-GONZÁLEZ, *New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory*, Adv. Math. **220** (2009), 1222–1264.
- [14] K. LI, K. MOEN AND W. SUN, *The sharp weighted bound for multilinear maximal functions and Calderón-Zygmund operators*, J. Four. Anal. Appl. **20** (2014), 751–765.
- [15] C. PÉREZ, G. PRADOLINI, R. H. TORRES AND R. TRUJILLO-GONZÁLEZ, *End-point estimates for iterated commutators of multilinear singular integrals*, Bull. London Math. Soc. **46** (2010), 26–42.

- [16] M. RAO AND Z. REN, *Theory of Orlicz spaces*, Monographs and Textbooks in Pure and Applied Mathematics, 146, Marcel Dekker Inc., New York, 1991.
- [17] E. M. STEIN, *Harmonic Analysis, Real Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Univ. Press, Princeton, NJ. 1993.