

DILATION-COMMUTING OPERATORS ON POWER-WEIGHTED ORLICZ CLASSES

RON KERMAN, RAMA RAWAT AND RAJESH K. SINGH

Abstract. Let Φ be a nondecreasing function from $\mathbb{R}_+ = (0, \infty)$ onto itself. Fix $\gamma \in \mathbb{R} = (-\infty, \infty)$ and let $L_{\Phi,t^\gamma}(\mathbb{R}_+)$ be the set of all Lebesgue-measurable functions f from \mathbb{R}_+ to \mathbb{R} for which

$$\int_{\mathbb{R}_+} \Phi(k|f(t)|) t^\gamma dt < \infty$$

for some $k > 0$. Define the gauge ρ_{Φ,t^γ} at $f \in L_{\Phi,t^\gamma}(\mathbb{R}_+)$ by

$$\rho_{\Phi,t^\gamma}(f) = \inf \left\{ \lambda > 0 : \int_{\mathbb{R}_+} \Phi\left(\frac{|f(t)|}{\lambda}\right) \frac{t^\gamma}{\lambda} dt \leq 1 \right\}.$$

Our principal goal in this paper is to find conditions on the nondecreasing functions Φ_1 and Φ_2 , $\gamma \in \mathbb{R}$ and an operator T so that the assertions

$$\rho_{\Phi_1,t^\gamma}(Tf) \leq C \rho_{\Phi_2,t^\gamma}(f) \quad (\text{G})$$

and

$$\int_{\mathbb{R}_+} \Phi_1(|(Tf)(t)|) t^\gamma dt \leq K \int_{\mathbb{R}_+} \Phi_2(|f(s)|) s^\gamma ds, \quad (\text{M})$$

concerning $f \in S(\mathbb{R}_+)$, the class of simple functions supported in \mathbb{R}_+ , are equivalent and to then find necessary and sufficient conditions in order that (M) holds.

In addition, we investigate the connection between (G) and the assertion that

$$T : \hat{L}_{\Phi_2,t^\gamma}(\mathbb{R}_+) \rightarrow L_{\Phi_1,t^\gamma}(\mathbb{R}_+),$$

where $\hat{L}_{\Phi_2,t^\gamma}(\mathbb{R}_+)$ is the closure of $S(\mathbb{R}_+)$ in $L_{\Phi_2,t^\gamma}(\mathbb{R}_+)$.

Mathematics subject classification (2010): Primary 42B25, 26D15, Secondary 28A25.

Keywords and phrases: Dilation-commuting operators, Orlicz spaces, norm inequalities, modular inequalities, Hardy operator, maximal function, Hilbert transform.

REFERENCES

- [1] S. BLOOM AND R. KERMAN, *Weighted L_Φ integral inequalities for operators of Hardy type*, Studia Math. **110**, 1 (1994), 35–52.
- [2] S. BLOOM AND R. KERMAN, *Weighted Orlicz space integral inequalities for the Hardy-Littlewood maximal operator*, Studia Math. **110**, 2 (1994), 149–167.
- [3] C. BENNETT AND R. SHARPLEY, *Interpolation of operators*, Pure and Applied Mathematics, vol. **129**, Academic Press Inc., Boston, MA, 1988.
- [4] A. CIANCHI, *Hardy inequalities in Orlicz spaces*, Trans. Amer. Math. Soc. **351**, 6 (1999), 2459–2478.
- [5] R. COIFMAN AND C. FEFFERMAN, *Weighted norm inequalities for maximal functions and singular integrals*, Studia Math. **51**, 3 (1974), 241–250.
- [6] M. FABIAN, P. HABAŁA, P. HÁJEK, V. MONTESINOS AND V. ZIZLER, *Banach space theory*, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, Springer, 2011.
- [7] R. KERMAN AND L. PICK, *Explicit formulas for optimal rearrangement-invariant norms in Sobolev imbedding inequalities*, Studia Math. **206**, 2 (2011), 97–119.

- [8] R. KERMAN AND A. TORCHINSKY, *Integral inequalities with weights for the Hardy maximal function*, Studia Math. **71**, 3 (1982), 277–284.
- [9] L. MALIGRANDA, *Orlicz spaces and interpolation*, Sem. Math. 5, Dep. Mat., Univ. Estadual de Campinas, Campinas SP, Brazil (1989).
- [10] W. MATUSZEWSKA AND W. ORLICZ, *On certain properties of φ -functions*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **8** (1960), 439–443.
- [11] S. MAZUR AND W. ORLICZ, *On some classes of linear spaces*, Studia Math. **17** (1958), 97–119.
- [12] W. ORLICZ, *A note on modular spaces. I*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **9** (1961), 157–162.
- [13] A. TORCHINSKY, *Real-variable methods in harmonic analysis*, Pure and Applied Mathematics, vol. **123**, Academic Press, Inc., Orlando, FL, 1986.