

MONOTONICITY AND INEQUALITIES INVOLVING ZERO-BALANCED HYPERGEOMETRIC FUNCTION

MIAO-KUN WANG, YU-MING CHU* AND WEN ZHANG

(Communicated by J. Pečarić)

Abstract. In the article, we present a monotonicity property involving the zero-balanced hypergeometric function $F(a, b; a+b; x)$ for all $a, b > 0$, and establish several sharp inequalities for $F(a, b; a+b; x)$ in the first quadrant of ab -plane, which are the generalizations of the previously results.

1. Introduction

For real numbers a, b , and c with $c \neq 0, -1, -2, \dots$, the Gaussian hypergeometric function $F(a, b; c; x)$ [49, 51, 52, 53, 64, 80, 88] is defined by

$$F(a, b; c; x) = {}_2F_1(a, b; c; x) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{x^n}{n!}$$

for $x \in (-1, 1)$, where $(a)_n$ is the Pochhammer symbol given by

$$(a)_n = a(a+1)\cdots(a+n-1) = \frac{\Gamma(n+a)}{\Gamma(a)}$$

for $n = 1, 2, \dots$, and $(a)_0 = 1$ for $a \neq 0$, $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$ is the classical Euler Gamma function [3, 15, 36, 38, 40, 69, 72, 79, 81, 83, 84, 85, 87, 90, 91, 92]. The function $F(a, b; c; x)$ is said to be zero-balanced if $c = a+b$. The asymptotic properties for $F(a, b; c; x)$ as $x \rightarrow 1$ are as follows (see [9, Theorems 1.19 and 1.48])

$$F(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}, \quad a+b < c, \quad (1.1)$$

$$F(a, b; c; x) = (1-x)^{c-a-b} F(c-a, c-b; c; x), \quad a+b > c, \quad (1.2)$$

Mathematics subject classification (2010): 33C05, 33E05.

Keywords and phrases: Gaussian hypergeometric function, generalized elliptic integral, monotonicity, inequality.

This work was supported by the Natural Science Foundation of China (Grant Nos. 11701176, 11626101, 11601485) and the Science and Technology Research Program of Zhejiang Educational Committee (Grant no. Y201635325).

* Corresponding author.

and when $a + b = c$,

$$B(a, b)F(a, b; c; x) + \log(1 - x) = R(a, b) + O((1 - x)\log(1 - x)), \quad (1.3)$$

where $B(z, w) = \Gamma(z)\Gamma(w)/[\Gamma(z+w)]$ ($\Re(z) > 0$, $\Re(w) > 0$) is the classical Beta function, and

$$R(a, b) = -\psi(a) - \psi(b) - 2\gamma,$$

$\psi(z) = \Gamma'(z)/\Gamma(z)$ ($\Re(z) > 0$) and γ is the Euler-Mascheroni constant. Equation (1.3) was established by Ramanujan [12, pp. 33–34].

It is well known that the Gaussian hypergeometric function $F(a, b; c; x)$ has many important applications in other branches of mathematics [14, 27, 28, 30, 31, 32, 33, 34, 42, 66], and a lot of special functions and elementary functions are the particular cases or limiting cases [2, 4, 29, 48, 63, 70]. For example, for $r \in [0, 1]$ and $a \in (0, 1/2]$, the complete elliptic integrals $\mathcal{K}(r)$ [1, 6, 7, 16, 17, 18, 20, 23, 25, 35, 41, 58, 59, 62, 68, 82] and $\mathcal{E}(r)$ [19, 21, 22, 24, 26, 50, 54, 57, 65, 75, 76, 77, 78] of the first and second kinds, and their generalizations $\mathcal{K}_a(r)$ and $\mathcal{E}_a(r)$ [8, 13, 55, 67, 89] can be expressed by $F(a, b; c; x)$ as follows:

$$\begin{aligned} \mathcal{K}(r) &= \frac{\pi}{2} F\left(\frac{1}{2}, \frac{1}{2}; 1; r^2\right), \quad \mathcal{K}(0) = \frac{\pi}{2}, \quad \mathcal{K}(1) = +\infty, \\ \mathcal{K}_a(r) &= \frac{\pi}{2} F(a, 1-a; 1; r^2), \quad \mathcal{K}_a(0) = \frac{\pi}{2}, \quad \mathcal{K}_a(1) = +\infty, \\ \mathcal{E}(r) &= \frac{\pi}{2} F\left(-\frac{1}{2}, \frac{1}{2}; 1; r^2\right), \quad \mathcal{E}(0) = \frac{\pi}{2}, \quad \mathcal{E}(1) = 1, \\ \mathcal{E}_a(r) &= \frac{\pi}{2} F(a-1, 1-a; 1; r^2), \quad \mathcal{E}_a(0) = \frac{\pi}{2}, \quad \mathcal{E}_a(1) = \frac{\sin(\pi a)}{2(1-a)}. \end{aligned}$$

In the past few years, $F(a, b; c; x)$ has been extensively studied by many authors in geometric function theory and modular equations. Numerous remarkable properties and inequalities for this function have been obtained. In [11, 37, 39, 86] the authors studied Legendre's relation of hypergeometric function and related \mathcal{M} -function, the Landen inequalities for zero-balanced hypergeometric function can be found in the literature [43, 47, 74], and the quotient of two hypergeometric functions as the generalization of the modulus of the plane Grötzsch ring in conformal geometry was introduced and investigated in [44, 45, 61]. For the above, or more properties see the Anderson-Vamanamurthy-Vuorinen book “Conformal Invariants, Inequalities, and Quasiconformal Mappings” [9] or a survey “Topics in special functions” [10].

Since the hypergeometric series $F(a, b; c; x)$ converges for all $x \in (-1, 1)$, the asymptotic properties and inequalities at $x = 1$ for $F(a, b; c; x)$ have been the subject of intensive research. Especially when $c = a + b$, equation (1.3) shows that the zero-balanced hypergeometric function $F(a, b; a+b; x)$, as well as its special cases \mathcal{K} and \mathcal{K}_a , has a logarithmic singularity at $x = 1$, namely,

$$F(a, b; a+b; x) \sim -\frac{1}{B(a, b)} \log(1-x), \quad x \rightarrow 1. \quad (1.4)$$

Thus it is very interesting to establish some asymptotic formulas or sharp inequalities for $F(a, b; a+b; x)$ as $x \rightarrow 1$.

Qiu and Vuorinen [43] proved the monotonicity properties for the functions $x \rightarrow xF(a, b; a+b; x)/\log[1/(1-x)]$, $x \rightarrow B(a, b)F(a, b; a+b; x) + (1/x)\log(1-x)$ and $x \rightarrow [B(a, b)F(a, b; a+b; x) + \log(1-x) - R(a, b)]/([(1-x)/x]\log[1/(1-x)])$. As applications, some sharp inequalities for $F(a, b; a+b; x)$ were derived. Recently, Wang, Chu and Song [60] refined Qiu and Vuorinen's results, in which the authors gave a complete answer to the monotonicity properties of the above functions for arbitrary $(a, b) \in \{(a, b) | a > 0, b > 0\}$.

For the complete elliptic integral $\mathcal{K}(r)$, Alzer [5] proved that the double inequality

$$1 + \alpha(1 - r^2) < \frac{\mathcal{K}(r)}{\log\left(\frac{4}{\sqrt{1-r^2}}\right)} < 1 + \beta(1 - r^2) \quad (1.5)$$

holds for all $r \in (0, 1)$ if and only if $\alpha \leq \pi/(4\log 2) - 1$ and $\beta \geq 1/4$. In 2015, Wang, Chu and Qiu [56] generalized inequality (1.5) and obtained

$$1 + \alpha(1 - r^2) < \frac{\mathcal{K}_a(r)}{\sin(\pi a) \log\left[\frac{e^{R(a,1-a)/2}}{\sqrt{1-r^2}}\right]} < 1 + \beta(1 - r^2) \quad (1.6)$$

for all $a \in (0, 1/2]$ and $r \in (0, 1)$ if and only if $\alpha \leq \pi/[R(a, 1-a) \sin \pi a] - 1$ and $\beta \geq a(1-a)$.

Very recently, making use of the following two-side inequality established in [56]

$$\frac{R(a, 1-a)^2}{(1+a-a^2)R(a, 1-a)-1} < \frac{\pi}{\sin(\pi a)} < (1+a-a^2)R(a, 1-a), \quad a \in (0, 1/2], \quad (1.7)$$

the authors [73] proved that the function

$$\begin{aligned} Y(r) &= \frac{2\mathcal{K}_a(r)}{\sin(\pi a)(1-r^2)\log[e^{R(a,1-a)/(1-r^2)}]} - \frac{1}{1-r^2} \\ &= \frac{B(a, 1-a)F(a, 1-a; 1; r^2) - \log[e^{R(a,1-a)/(1-r^2)}]}{(1-r^2)\log[e^{R(a,1-a)/(1-r^2)}]} \end{aligned} \quad (1.8)$$

is strictly increasing from $(0, 1)$ onto $(\pi/[R(a, 1-a) \sin \pi a] - 1, a(1-a))$ for all $a \in (0, 1/2]$, and consequently inequality (1.6) can be also derived. Actually, in order to search for beautiful inequalities for the zero-balanced hypergeometric function, or $\mathcal{K}(r)$ and $\mathcal{K}_a(r)$, Qiu and Vuorinen [43] raised the following open problem about a generalization of $Y(r)$.

PROBLEM 1.1. Let $a, b \in (0, 1)$ with $a+b < 1$ and define F^* on $(0, 1)$ by

$$F^*(x) = \frac{B(a, b)F(a, b; a+b; x) - \log[e^{R(a,b)/(1-x)}]}{(1-x)\log[e^{R(a,b)/(1-x)}]}. \quad (1.9)$$

Is it true that the function F^* has a Maclaurin expansion $\sum_{n=0}^{\infty} d_n x^n$ with non-negative coefficients d_n .

Problem 1.1 is very difficult, and until now, it is still open. The main purpose of this paper is to prove the monotonicity property of $F^*(x)$ for arbitrary $(a, b) \in \{(a, b) | a > 0, b > 0\}$. This result lead to some sharp inequalities for $F(a, b; a + b; x)$, which extend inequality (1.6).

2. Main results

Throughout this paper, for $a, b > 0$, we denote

$$M_1(a, b) = \frac{1 - abB(a, b)/(a + b)}{1 - ab(a + 1)(b + 1)B(a, b)/[(a + b)(a + b + 1)]} \quad (2.1)$$

and

$$M_2(a, b) = R(a, b) - 1 - \frac{R(a, b)}{B(a, b) - R(a, b)} \left[1 - B(a, b) \frac{ab}{a + b} \right]. \quad (2.2)$$




Figure 1: The regions D_{11} , D_{12} , D_{21} , D_{22} , D_{12}^* , D_{12}^{**} , D_{21}^* and D_{21}^{**} , where $C_1 : 2ab - a - b = 0$, $C_2 : R(a, b) = 0$, $C_3 : R(a, b) = 1$, $C_4 : M_2(a, b) = 0$, $C_5 : M_1(a, b) = R(a, b) - 1$.

Let

$$D_1 = \{(a, b) | a, b > 0, R(a, b) \geq 1\},$$

$$D_2 = \{(a, b) | a, b > 0, R(a, b) < 1\},$$

$$\begin{aligned}
D_{11} &= \{(a,b) | a,b > 0, R(a,b) \geq 1, R(a,b) - 1 \geq M_1(a,b)\}, \\
D_{12} &= \{(a,b) | a,b > 0, R(a,b) \geq 1, R(a,b) - 1 < M_1(a,b)\}, \\
D_{21} &= \{(a,b) | a,b > 0, R(a,b) < 1, 2ab - a - b < 0\}, \\
D_{22} &= \{(a,b) | a,b > 0, R(a,b) < 1, 2ab - a - b \geq 0\}, \\
D_{12}^* &= \{(a,b) | a,b > 0, R(a,b) \geq 1, R(a,b) - 1 < M_1(a,b), M_2(a,b) \geq 0\}, \\
D_{12}^{**} &= \{(a,b) | a,b > 0, R(a,b) \geq 1, R(a,b) - 1 < M_1(a,b), M_2(a,b) < 0\}, \\
D_{21}^* &= \{(a,b) | a,b > 0, 0 \leq R(a,b) < 1, 2ab - a - b < 0\}
\end{aligned}$$

and

$$D_{21}^{**} = \{(a,b) | a,b > 0, R(a,b) < 0, 2ab - a - b < 0\}.$$

Then $D_{12}^* \cup D_{12}^{**} = D_{12}$, $D_{21}^* \cup D_{21}^{**} = D_{21}$, $D_{11} \cup D_{12} = D_1$, $D_{21} \cup D_{22} = D_2$ and $D_1 \cup D_2 = \{a,b | a,b > 0\}$ (see Figure 1).

REMARK 2.1. According to Lemma 3.4 in Section 3, inequalities $1 - abB(a,b)/(a+b) > 0$ and $1 - ab(a+1)(b+1)B(a,b)/(a+b)/(a+b+1) > 0$ hold for all $a,b > 0$. Hence $M_1(a,b)$ in (2.1) is positive for each $(a,b) \in \{(a,b) | a > 0, b > 0\}$. On the other hand, Theorem 1.52(2) in [9] shows that the function $x \rightarrow B(a,b)F(a,b;a+b;x) + \log(1-x)(a,b > 0)$ is strictly decreasing from $(0,1)$ onto $(R(a,b), B(a,b))$. Thus $B(a,b) > R(a,b)$ for all $a,b > 0$, and thereby $M_2(a,b)$ is well-defined.

THEOREM 2.2. *Let*

$$F(x) = \frac{(1-x)\log[e^{R(a,b)}/(1-x)]}{B(a,b)F(a,b;a+b;x) - \log[e^{R(a,b)}/(1-x)]}, \quad x \in (0,1).$$

Then the following statements hold

- (1) *If $(a,b) \in D_{11} \cup D_{12}^*$, then the function $F(x)$ is strictly decreasing from $(0,1)$ onto $(1/(ab), R(a,b)/[B(a,b) - R(a,b)])$;*
- (2) *If $(a,b) \in D_{22}$, then the function $F(x)$ is strictly increasing from $(0,1)$ onto $(R(a,b)/[B(a,b) - R(a,b)], 1/(ab))$;*
- (3) *If $(a,b) \in D_{12}^{**} \cup D_{21}$, then there exists $x_0 \in (0,1)$ such that $F(x)$ is strictly increasing on $(0,x_0)$, and strictly decreasing on $(x_0,1)$.*

Using Theorem 2.2, we can derive the monotonicity property of F^* immediately.

COROLLARY 2.3. *Let*

$$F^*(x) = \frac{1}{F(x)} = \frac{B(a,b)F(a,b;a+b;x) - \log[e^{R(a,b)}/(1-x)]}{(1-x)\log[e^{R(a,b)}/(1-x)]}.$$

Then the following statements hold

- (1) *$(a,b) \in \{(a,b) | R(a,b) \geq 0\}$. Then F^* is strictly increasing from $(0,1)$ onto $(B(a,b)/R(a,b) - 1, ab)$ if $(a,b) \in D_{11} \cup D_{12}^*$, and if $(a,b) \in D_{12}^{**} \cup D_{21}^*$, then there*

exists $x_0 \in (0, 1)$ such that F^* is strictly decreasing on $(0, x_0)$, and strictly increasing on $(x_0, 1)$. Moreover, for $(a, b) \in D_{11} \cup D_{12}^*$, the double inequality

$$1 + \frac{B(a, b) - R(a, b)}{R(a, b)}(1-x) < \frac{B(a, b)F(a, b, ; a+b; x)}{\log [e^{R(a,b)} / (1-x)]} < 1 + ab(1-x) \quad (2.3)$$

holds for all $x \in (0, 1)$ with the best possible constants $[B(a, b) - R(a, b)]/R(a, b)$ and ab , and for $(a, b) \in D_{12}^{**} \cup D_{21}^*$, inequality

$$\frac{B(a, b)F(a, b, ; a+b; x)}{\log [e^{R(a,b)} / (1-x)]} < 1 + \max \left\{ ab, \frac{B(a, b) - R(a, b)}{R(a, b)} \right\} (1-x) \quad (2.4)$$

holds for all $x \in (0, 1)$.

(2) $(a, b) \in \{(a, b) | R(a, b) < 0\}$. Denote $x_0^* = x_0^*(a, b)$ by the solution of equation $R(a, b) = \log(1-x)$, one has

(i) If $(a, b) \in D_{21}^{**}$, then F^* is strictly decreasing from $(0, x_0^*)$ onto $(-\infty, [B(a, b) - R(a, b)]/R(a, b))$, and there exist $x_0^{**} \in (x_0^*, 1)$ such that F^* is strictly decreasing on (x_0^*, x_0^{**}) and strictly increasing on $(x_0^{**}, 1)$;

(ii) If $(a, b) \in D_{22}$, then F^* is strictly decreasing from $(0, x_0^*)$ onto $(-\infty, [B(a, b) - R(a, b)]/R(a, b))$, and strictly decreasing from $(x_0^*, 1)$ onto $(ab, +\infty)$.

REMARK 2.4. If $b = 1 - a > 0$, then $B(a, 1-a) = \pi / \sin(\pi a)$, and

$$M_2(a, b) = \frac{B(a, 1-a)[(1+a-a^2)R(a, 1-a) - 1] - R(a, 1-a)^2}{B(a, 1-a) - R(a, 1-a)}.$$

It follows (1.7) that $\{(a, b) | a, b > 0, a+b=1\} \subset D_{11} \cup D_{12}^*$, so that $Y(r)$ in (1.8), which is equal to $F^*(r^2)$, is strictly increasing on $(0, 1)$ by Corollary 2.3. Besides, Substituting r^2 for x in inequality (2.3), we obtain inequality (1.6).

3. Lemmas

In order to prove Theorem 2.2 we need several lemmas, which we present in this section.

LEMMA 3.1. ([9, Theorem 1.25]) For $-\infty < a < b < \infty$, let $f, g : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$, and be differentiable on (a, b) , let $g'(x) \neq 0$ on (a, b) . If $f'(x)/g'(x)$ is increasing (decreasing) on (a, b) , then so are

$$\frac{f(x) - f(a)}{g(x) - g(a)} \text{ and } \frac{f(x) - f(b)}{g(x) - g(b)}.$$

If $f'(x)/g'(x)$ is strictly monotone, then the monotonicity in the conclusion is also strict.

LEMMA 3.2. ([74, Theorem 2.1]) Suppose that the power series $f(x) = \sum_{n=0}^{\infty} a_n x^n$ and $g(x) = \sum_{n=0}^{\infty} b_n x^n$ have the radius of convergence $r > 0$ with $b_n > 0$ for all $n \in \mathbb{N}$

$\{0, 1, 2, \dots\}$. Let $h(x) = f(x)/g(x)$ and $H_{f,g} = (f'/g')g - f$, then the following statements] are true:

- (1) If the non-constant sequence $\{a_n/b_n\}_{n=0}^{\infty}$ is increasing (decreasing), then $h(x)$ is strictly increasing (decreasing) on $(0, r)$;
- (2) If the non-constant sequence $\{a_n/b_n\}$ is increasing (decreasing) for $0 < n \leq n_0$ and decreasing (increasing) for $n > n_0$, then the function h is strictly increasing (decreasing) on $(0, r)$ if and only if $H_{f,g}(r^-) \geq (\leq) 0$. While if $H_{f,g}(r^-) < (>) 0$, then there exists $x_0 \in (0, r)$ such that $h(x)$ is strictly increasing (decreasing) on $(0, x_0)$ and strictly decreasing (increasing) on (x_0, r) .

LEMMA 3.3. ([71, Theorem 9]) For $-\infty \leq a < b \leq \infty$, let f and g be differentiable functions on (a, b) with $g' \neq 0$ on (a, b) , $\text{sgn}(\cdot)$ be signum function, and $H_{f,g} = (f'/g')g - f$. Suppose that (i) $g' \neq 0$ on (a, b) ; (ii) $f(b^-) = g(b^-) = 0$; (iii) there exists $c \in (a, b)$ such that f'/g' is increasing (decreasing) on (a, c) and decreasing (increasing) on (c, b) . Then

- (1) when $\text{sgn}(g')\text{sgn}H_{f,g}(a^+) \leq (\geq) 0$, f/g is decreasing (increasing) on (a, b) ;
- (2) when $\text{sgn}(g')\text{sgn}H_{f,g}(a^+) > (<) 0$, there is a unique number $x_b \in (a, b)$ such that f/g is increasing (decreasing) on (a, x_b) and decreasing (increasing) on (x_b, b) .

LEMMA 3.4. ([9, Lemma 1.50 (1)]) For $a, b > 0$, the sequence

$$f(n) = \frac{(a)_n(b)_n}{(a+b)_n(n-1)!}$$

is strictly increasing to the limit $1/B(a, b)$.

LEMMA 3.5. For $a, b > 0$ with $1/a + 1/b \leq 2$, then

$$R(a, b) \leq 0,$$

with equality if and only if $a = b = 1$.

Proof. Since $R(a, b) = -\Psi(a) - \Psi(b) - 2\gamma$ is strictly decreasing in a and b , we have

$$\begin{aligned} R(a, b) &\leq R\left(a, \frac{a}{2a-1}\right) = -\Psi(a) - \Psi\left(\frac{a}{2a-1}\right) - 2\gamma \\ &= \frac{1}{a} + \frac{1}{\frac{a}{2a-1}} - \sum_{k=1}^{\infty} \left(\frac{2}{k} - \frac{1}{k+a} - \frac{1}{k+\frac{a}{2a-1}} \right) \\ &= 2 - \sum_{k=1}^{\infty} \left[\frac{\frac{2a^2}{2a-1}(k+1)}{k\left(k^2 + \frac{2a^2}{2a-1}k + \frac{a^2}{2a-1}\right)} \right] \end{aligned} \tag{3.1}$$

by employing $\Psi(x) = -\gamma - 1/x - \sum_{k=1}^{\infty} [1/k - 1/(k+x)]$. It is easy to check that $2a^2/(2a-1) \geq 2$ for $a > 1/2$, and $x \rightarrow x/(k^2 + xk + x/2)$ ($k \in \mathbb{N}^+$) is strictly increasing on $[2, \infty)$. Thus from (3.1) one has

$$\begin{aligned} R(a, b) &\leq 2 - \sum_{k=1}^{\infty} \left[\frac{\frac{2a^2}{2a-1}(k+1)}{k \left(k^2 + \frac{2a^2}{2a-1}k + \frac{a^2}{2a-1} \right)} \right] \\ &\leq 2 - \sum_{k=1}^{\infty} \frac{2(k+1)}{k(k^2 + 2k + 1)} = 2 - \sum_{k=1}^{\infty} \frac{2}{k(k+1)} = 0. \end{aligned} \quad (3.2)$$

Both inequalities (3.1) and (3.2) become equalities if and only if $a = b = 1$. This completes the proof of Lemma 3.5. \square

LEMMA 3.6. Let $\{A_n\}_{n=1}^{\infty}$ and $\{B_n\}_{n=1}^{\infty}$ be defined by

$$A_1 = R(a, b) - 1, A_n = \frac{1}{n-1} (n \geq 2), B_n = 1 - \frac{(a)_n(b)_n}{(a+b)_n(n-1)!} B(a, b) \quad (3.3)$$

with $a, b > 0$. Then the sequence $\{A_n/B_n\}$ is strictly decreasing for $n \geq 2$.

Proof. By Lemma 3.4, $B_n > 0$ for all $a, b > 0$ and $n \geq 1$. Let

$$C_n = \frac{B_n}{A_n} = n-1 - B(a, b) \frac{(a)_n(b)_n}{(a+b)_n(n-2)!}, \quad n \geq 2. \quad (3.4)$$

Then

$$\begin{aligned} C_{n+1} - C_n &= n - B(a, b) \frac{(a)_{n+1}(b)_{n+1}}{(a+b)_{n+1}(n-1)!} - (n-1) + B(a, b) \frac{(a)_n(b)_n}{(a+b)_n(n-2)!} \\ &= 1 - B(a, b) \frac{(a)_n(b)_n}{(a+b)_{n+1}(n-1)!} [(n+a)(n+b) - (n+a+b)(n-1)] \\ &= 1 - B(a, b) \frac{(a)_n(b)_n}{(a+b)_n(n-1)!} \frac{n+ab+a+b}{n+a+b}. \end{aligned} \quad (3.5)$$

Let

$$D_n = \frac{(a)_n(b)_n}{(a+b)_n(n-1)!} \frac{n+ab+a+b}{n+a+b}. \quad (3.6)$$

Then

$$\frac{D_{n+1}}{D_n} - 1 = \frac{ab(1+a+b+ab)}{n(n+1+a+b)(n+ab+a+b)} > 0. \quad (3.7)$$

It follows from Lemma 3.4 that

$$\lim_{n \rightarrow \infty} D_n = \frac{1}{B(a, b)}. \quad (3.8)$$

Hence, by (3.6)–(3.8), $D_n < 1/B(a, b)$ for $a, b > 0$, so that $\{C_n\}$ is strictly increasing for $n \geq 2$ from (3.5).

Therefore, Lemma 3.6 follows from (3.4) and the monotonicity of the sequence $\{C_n\}$. \square

LEMMA 3.7. For $a, b > 0$, let

$$f_1(x) = R(a, b) - 1 + \log\left(\frac{1}{1-x}\right) \quad (3.9)$$

and

$$\begin{aligned} g_1(x) &= -B(a, b) \frac{ab}{a+b} F(a+1, b+1; a+b+1; x) + \frac{1}{1-x} \\ &= \frac{1}{1-x} \left[1 - B(a, b) \frac{ab}{a+b} F(a, b; a+b+1; x) \right]. \end{aligned} \quad (3.10)$$

Then

$$\lim_{x \rightarrow 1^-} H_{f_1, g_1}(x) = \lim_{x \rightarrow 1^-} \left[\frac{f'_1(x)}{g'_1(x)} g_1(x) - f_1(x) \right] = \frac{2ab - a - b}{ab}. \quad (3.11)$$

Proof. Differentiating f_1 and g_1 gives

$$f'_1(x) = \frac{1}{1-x}, \quad (3.12)$$

and

$$\begin{aligned} g'_1(x) &= \frac{1}{(1-x)^2} \left[1 - \frac{abB(a, b)}{a+b} F(a, b; a+b+1; x) \right] \\ &\quad - \frac{1}{1-x} \frac{a^2 b^2 B(a, b)}{(a+b)(a+b+1)} F(a+1, b+1; a+b+2; x), \end{aligned} \quad (3.13)$$

where we use the derivative formula of hypergeometric function

$$\frac{dF(a, b; c; x)}{dx} = \frac{ab}{c} F(a+1, b+1; c+1; x).$$

It follows from (3.9), (3.10), (3.12) and (3.13) that

$$\begin{aligned} H_{f_1, g_1}(x) &= \frac{f'_1(x)}{g'_1(x)} g_1(x) - f_1(x) \\ &= \frac{1 - \frac{abB(a, b)}{a+b} F(a, b; a+b+1; x)}{h(x)} - \left[R(a, b) - 1 + \log\left(\frac{1}{1-x}\right) \right] \\ &= \frac{(1-x)^{-1} \left[1 - \frac{abB(a, b)}{a+b} F(a, b; a+b+1; x) - h(x) (R(a, b) - 1 + \log(\frac{1}{1-x})) \right]}{(1-x)^{-1} h(x)} \end{aligned} \quad (3.14)$$

where

$$\begin{aligned} h(x) &= 1 - \frac{abB(a, b)}{a+b} F(a, b; a+b+1; x) \\ &\quad - (1-x) \frac{a^2 b^2 B(a, b)}{(a+b)(a+b+1)} F(a+1, b+1; a+b+2; x). \end{aligned} \quad (3.15)$$

Since Gaussian hypergeometric function $F(a, b; c; x)$ has the following asymptotic expansions (see [46, (2.10)])

$$F(a+1, b+1; a+b+2; x) = \frac{1}{B(a+1, b+1)} \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n}{(n!)^2} u_n (1-x)^n,$$

$$\begin{aligned} F(a, b; a+b+1; z) &= \frac{\Gamma(a+b+1)}{\Gamma(a+1)\Gamma(b+1)} \\ &\quad + \frac{\Gamma(a+b+1)}{\Gamma(a)\Gamma(b)} (x-1) \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n}{(n!)(n+1)!} v_n (1-x)^n, \end{aligned}$$

where

$$u_n = 2\Psi(n+1) - \Psi(n+a+1) - \Psi(n+b+1) - \log(1-x) \quad (3.16)$$

and

$$v_n = \Psi(n+1) + \Psi(n+2) - \Psi(n+a+1) - \Psi(n+b+1) - \log(1-x), \quad (3.17)$$

one has

$$\begin{aligned} &1 - \frac{abB(a, b)}{a+b} F(a, b; a+b+1; x) \\ &= 1 - \frac{abB(a, b)}{a+b} \frac{\Gamma(a+b+1)}{\Gamma(a+1)\Gamma(b+1)} - \frac{abB(a, b)}{a+b} \frac{\Gamma(a+b+1)}{\Gamma(a)\Gamma(b)} (x-1) \\ &\quad \times \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n}{(n!)(n+1)!} v_n (1-x)^n \\ &= ab(1-x) \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n}{(n!)(n+1)!} v_n (1-x)^n, \end{aligned}$$

$$\begin{aligned} &(1-x)^{-1} \left[1 - \frac{abB(a, b)}{a+b} F(a, b; a+b+1; x) - h(x) \left(R(a, b) - 1 + \log\left(\frac{1}{1-x}\right) \right) \right] \\ &= (1-x)^{-1} \left[ab(1-x) \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n}{(n!)(n+1)!} v_n (1-x)^n - (R(a, b) - 1 - \log(1-x)) \right. \\ &\quad \times \left(ab(1-x) \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n}{(n!)(n+1)!} v_n (1-x)^n \right. \\ &\quad \left. \left. - (1-x) \frac{a^2 b^2 B(a, b)}{(a+b)(a+b+1)} \frac{1}{B(a+1, b+1)} \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n}{(n!)^2} u_n (1-x)^n \right) \right] \\ &= ab \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n}{(n!)(n+1)!} v_n (1-x)^n - (R(a, b) - 1 - \log(1-x)) \\ &\quad \times \left(ab \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n}{(n!)(n+1)!} v_n (1-x)^n - ab \sum_{n=0}^{\infty} \frac{(a+1)_n (b+1)_n}{(n!)^2} u_n (1-x)^n \right) \end{aligned}$$

$$= ab \sum_{n=0}^{\infty} \frac{(a+1)_n(b+1)_n}{(n!)(n+1)!} (1-x)^n \{ [v_n - (n+1)u_n] [1 - R(a,b) + \log(1-x)] + v_n \}, \quad (3.18)$$

$$\begin{aligned} (1-x)^{-1}h(x) &= \frac{1}{1-x} \left(ab(1-x) \sum_{n=0}^{\infty} \frac{(a+1)_n(b+1)_n}{(n!)(n+1)!} v_n (1-x)^n \right) \\ &\quad - \frac{a^2 b^2 B(a,b)}{(a+b)(a+b+1)} F(a+1, b+1; a+b+2; x) \\ &= ab \sum_{n=0}^{\infty} \frac{(a+1)_n(b+1)_n}{(n!)(n+1)!} v_n (1-x)^n - ab \sum_{n=0}^{\infty} \frac{(a+1)_n(b+1)_n}{(n!)^2} u_n (1-x)^n \\ &= ab \sum_{n=0}^{\infty} \frac{(a+1)_n(b+1)_n}{(n!)(n+1)!} (1-x)^n [v_n - (n+1)u_n]. \end{aligned} \quad (3.19)$$

It follows from (3.14)–(3.19) that

$$\begin{aligned} &\lim_{x \rightarrow 1^-} H_{f_1, g_1}(x) \\ &= \lim_{x \rightarrow 1^-} \frac{(v_0 - u_0)[1 - R(a,b) + \log(1-x)] + v_0 + o((1-x)\log(1-x))}{v_0 - u_0 + o((1-x)\log(1-x))} \\ &= \lim_{x \rightarrow 1^-} \frac{1 - R(a,b) + \log(1-x) + \Psi(1) + \Psi(2) - \Psi(a+1) - \Psi(b+1) - \log(1-x)}{1 + o((1-x)\log(1-x))} \\ &= 1 - R(a,b) + 2\Psi(1) + 1 - \Psi(a) - \Psi(b) - \frac{1}{a} - \frac{1}{b} = \frac{2ab - a - b}{ab}. \quad \square \end{aligned}$$

4. Proof of Theorem 2.2

Obviously

$$\lim_{x \rightarrow 0^+} F(x) = \frac{R(a,b)}{B(a,b) - R(a,b)}, \quad (4.1)$$

and making use of L'Hôpital's rule we get

$$\begin{aligned} \lim_{x \rightarrow 1^-} F(x) &= \lim_{x \rightarrow 1^-} \frac{1 - R(a,b) + \log(1-x)}{B(a,b) \frac{ab}{a+b} F(a+1, b+1; a+b+1; x) - \frac{1}{1-x}} \\ &= \lim_{x \rightarrow 1^-} \frac{(1-x)[1 - R(a,b)] + (1-x)\log(1-x)}{B(a,b) \frac{ab}{a+b} F(a, b; a+b+1; x) - 1} \\ &= \lim_{x \rightarrow 1^-} \frac{R(a,b) - 2 - \log(1-x)}{B(a,b) \frac{a^2 b^2}{(a+b)(a+b+1)} F(a+1, b+1; a+b+2; x)} \\ &= \lim_{x \rightarrow 1^-} \frac{1}{(1-x) \frac{a^2 b^2 B(a,b)}{(a+b)(a+b+1)} \frac{(a+1)(b+1)}{(a+b+2)} F(a+2, b+2; a+b+3; x)} \end{aligned}$$

$$\begin{aligned}
&= \lim_{x \rightarrow 1^-} \frac{1}{\frac{d^2 b^2 B(a,b)}{(a+b)(a+b+1)} \frac{(a+1)(b+1)}{(a+b+2)} F(a+1, b+1; a+b+3; x)} \\
&= \frac{1}{B(a,b) \frac{a^2 b^2 (a+1)(b+1)}{(a+b)(a+b+1)(a+b+2)} \frac{\Gamma(a+b+3)\Gamma(1)}{\Gamma(a+2)\Gamma(b+2)}} = \frac{1}{ab}.
\end{aligned} \tag{4.2}$$

If we let

$$f(x) = (1-x)[R(a,b) - \log(1-x)] \tag{4.3}$$

and

$$g(x) = B(a,b)F(a,b;a+b;x) - \log\left(\frac{e^{R(a,b)}}{1-x}\right), \tag{4.4}$$

then $F(x) = f(x)/g(x)$, $f(1^-) = g(1^-) = 0$,

$$f'(x) = -R(a,b) + \log(1-x) + 1 = -(R(a,b) - 1) - \sum_{n=2}^{\infty} \frac{x^{n-1}}{n-1} = -\sum_{n=1}^{\infty} A_n x^{n-1},$$

$$\begin{aligned}
g'(x) &= B(a,b) \frac{ab}{a+b} F(a+1, b+1; a+b+1; x) - \frac{1}{1-x} \\
&= B(a,b) \sum_{n=1}^{\infty} \frac{(a)_n (b)_n}{(a+b)_n (n-1)!} x^{n-1} - \frac{1}{1-x} = -\sum_{n=1}^{\infty} B_n x^{n-1},
\end{aligned} \tag{4.5}$$

$$\frac{f'(x)}{g'(x)} = \frac{f_1(x)}{g_1(x)} = \frac{\sum_{n=1}^{\infty} A_n x^{n-1}}{\sum_{n=1}^{\infty} B_n x^{n-1}} \tag{4.6}$$

and thereby

$$\begin{aligned}
\lim_{x \rightarrow 0^+} H_{f,g}(x) &= \lim_{x \rightarrow 0^+} \left[\frac{f'(x)}{g'(x)} g(x) - f(x) \right] \\
&= \frac{R(a,b) - 1}{1 - \frac{ab}{a+b} B(a,b)} [B(a,b) - R(a,b)] - R(a,b) \\
&= \frac{B(a,b) - R(a,b)}{1 - \frac{ab}{a+b} B(a,b)} M_2(a,b).
\end{aligned} \tag{4.7}$$

Here

$$A_1 = R(a,b) - 1, \quad A_n = \frac{1}{n-1} (n \geq 2), \quad B_n = 1 - \frac{(a)_n (b)_n}{(a+b)_n (n-1)!} B(a,b), \tag{4.8}$$

and $f_1(x)$ and $g_1(x)$ are defined by (3.9) and (3.10) in Lemma 3.7.

Next, we divide the proof into five cases.

Case 1: $(a,b) \in D_{11}$. Then $A_1/B_1 \geq A_2/B_2$, and from Lemma 3.6 we know that the non-constant sequence $\{A_n/B_n\}$ is strictly decreasing. Equation (4.6) and Lemma 3.2(1) imply that the function $f_1(x)/g_1(x)$ is strictly decreasing on $(0, 1)$, and so is $F(x)$ by applying Lemma 3.1.

Case 2: $(a, b) \in D_{12}^*$. Then $A_1/B_1 < A_2/B_2$, and the non-constant sequence $\{A_n/B_n\}$ is strictly increasing for $1 \leq n \leq 2$, and strictly decreasing for $n \geq 2$. By Lemma 3.5, we conclude that $H_{f_1,g_1}(1^-) = (2ab - a - b)/(ab) < 0$. So that equation (4.6) and Lemma 3.2(2) lead to the conclusion that there exists $\xi \in (0, 1)$ such that $f'(x)/g'(x)$ is strictly increasing on $(0, \xi)$, and strictly decreasing on $(\xi, 1)$.

Equations (2.1), (4.5) and (4.7) show that $\operatorname{sgn}(g') < 0$ and $H_{f,g}(0^+) \geq 0$. Thus by application of Lemma 3.3(1) one has that $F(x)$ is strictly decreasing on $(0, 1)$.

Case 3: $(a, b) \in D_{12}^{**}$. Then $A_1/B_1 < A_2/B_2$, with the similar argument in Case 2, we conclude that there exists $\eta \in (0, 1)$ such that $f'(x)/g'(x)$ is strictly increasing on $(0, \eta)$, and strictly decreasing on $(\eta, 1)$. Since $\operatorname{sgn}(g') < 0$ and $H_{f,g}(0^+) > 0$, by Lemma 3.3(2) one has that there exists $x_0 \in (0, 1)$ such that $F(x)$ is strictly increasing on $(0, x_0)$, and strictly decreasing on $(x_0, 1)$.

Case 4: $(a, b) \in D_{22}$. Then $A_1/B_1 < 0 < A_2/B_2$, and the non-constant sequence $\{A_n/B_n\}$ is strictly increasing for $1 \leq n \leq 2$, and strictly decreasing for $n \geq 2$. Since $H_{f_1,g_1}(1^-) = (2ab - a - b)/(ab) \geq 0$, Lemma 3.2(2) and (4.6) lead to the conclusion that $f'(x)/g'(x)$ is strictly increasing on $(0, 1)$, and so is $F(x)$ by applying Lemma 3.1.

Case 5: $(a, b) \in D_{21}$. Then $A_1/B_1 < 0 < A_2/B_2$, and the non-constant sequence $\{A_n/B_n\}$ is strictly increasing for $1 \leq n \leq 2$, and strictly decreasing for $n \geq 2$. Lemma 3.2(2) and $H_{f_1,g_1}(1^-) = (2ab - a - b)/(ab) < 0$ show that there exist $\delta \in (0, 1)$ such that $f'(x)/g'(x)$ is strictly increasing on $(0, \delta)$, and strictly decreasing on $(\delta, 1)$.

Next we claim that $H_{f,g}(0^+) < 0$ for $(a, b) \in D_{21}$. In fact, by Remark 2.1, $B(a, b) > R(a, b)$ and $1 > abB(a, b)/(a + b)$ for all $a, b > 0$, and it is clear to see that $M_2(a, b) < 0$ for $(a, b) \in D_{21}^*$, and for $(a, b) \in D_{21}^{**}$,

$$M_2(a, b) = \frac{B(a, b)R(a, b) - B(a, b) - R(a, b)^2 + abB(a, b)R(a, b)/(a + b)}{B(a, b) - R(a, b)} < 0.$$

Finally, inequalities $H_{f,g}(0^+) < 0$ for $(a, b) \in D_{21}$ and $\operatorname{sgn}(g') < 0$ for $x \in (0, 1)$ together with Lemma 3.3(2) and the piecewise monotonicity of $f'(x)/g'(x)$ yield that there exists x_1 such that $F(x)$ is strictly increasing on $(0, x_1)$ and strictly decreasing on $(x_1, 1)$.

REFERENCES

- [1] M. ABRAMOWITZ, I. A. STEGUN, *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*, Dover, New York, 1965.
- [2] M. ADIL KHAN, Y.-M. CHU, A. KASHURI, R. LIKO, *Conformable fractional integrals versions of Hermite-Hadamard inequalities and their generalizations*, J. Funct. Spaces **2018** (2018), Article ID 6928130, 9 pages.
- [3] M. ADIL KHAN, Y.-M. CHU, A. KASHURI, R. LIKO, *Hermite-Hadamard type fractional integral inequalities for $MT_{(r,g,m,\phi)}$ -preinvex functions*, J. Comput. Anal. Appl. **28**, (8) (2019), 1487–1503.
- [4] M. ADIL KHAN, Y.-M. CHU, T. U. KHAN, J. KHAN, *Some new inequalities of Hermite-Hadamard type for s -convex functions with applications*, Open Math. **15** (2017), 1414–1430.
- [5] H. ALZER, *Sharp inequalities for the complete elliptic integral of the first kind*, Math. Proc. Cambridge Philos. Soc. **24**, 2 (1998), 309–314.
- [6] H. ALZER, S.-L. QIU, *Monotonicity theorems and inequalities for the complete elliptic integrals*, J. Comput. Appl. Math. **172**, 2 (2004), 289–312.

- [7] G. D. ANDERSON, S.-L. QIU, M. K. VAMANAMURTHY, *Elliptic integral inequalities, with applications*, Constr. Approx. **14**, 2 (1998), 195–207.
- [8] G. D. ANDERSON, S.-L. QIU, M. K. VAMANAMURTHY, M. VUORINEN, *Generalized elliptic integrals and modular equations*, Pacific J. Math. **192**, 1 (2000), 1–37.
- [9] G. D. ANDERSON, M. K. VAMANAMURTHY, M. VUORINEN, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.
- [10] G. D. ANDERSON, M. K. VAMANAMURTHY, M. VUORINEN, *Topics in special functions*, In: Papers on Analysis: A volume dedicated to Olli Martio on the occasion of his 60th birthday, Report Univ. Jyväskylä, **83** (2001), 5–26.
- [11] R. BALASUBRAMANIAN, S. NAIK, S. PONNUSAMY, M. VUORINEN, *Elliott's identity and hypergeometric functions*, J. Math. Anal. Appl. **271**, 1 (2002), 232–256.
- [12] B. C. BERNDT, *Ramanujan's Notebooks II*, Springer-Verlag, New York, 1989.
- [13] B. A. BHAYO, M. VUORINEN, *On generalized complete elliptic integrals and modular functions*, Proc. Edinb. Math. Soc. (2) **55**, 3 (2012), 591–611.
- [14] Y.-M. CHU, Y.-H. HAO, X.-G. LIU, *Global weak solutions to a general liquid crystals system*, Discrete Contin. Dyn. Syst. **33**, 7 (2013), 2681–2710.
- [15] Y.-M. CHU, Y.-M. LI, W.-F. XIA, X.-H. ZHANG, *Best possible inequalities for the harmonic mean of error function*, J. Inequal. Appl. **2014** (2014), Article 525, 9 pages.
- [16] Y.-M. CHU, Y.-F. QIU, M.-K. WANG, *Hölder mean inequalities for the complete elliptic integrals*, Integral Transforms Spec. Funct. **23**, 7 (2012), 521–527.
- [17] Y.-M. CHU, Y.-F. QIU, M.-K. WANG, Y.-F. QIU, *On Alzer and Qiu's conjecture for complete elliptic integral and inverse hyperbolic tangent function*, Abstr. Appl. Anal. **2011** (2011), Article ID 697547, 7 pages.
- [18] Y.-M. CHU, M.-K. WANG, *Optimal inequalities between harmonic, geometric, logarithmic, and arithmetic-geometric means*, J. Appl. Math. **2011** (2011), Article ID 618929, 9 pages.
- [19] Y.-M. CHU, M.-K. WANG, *Optimal Lehmer mean bounds for the Toader mean*, Results Math. **61**, 3–4 (2012), 223–229.
- [20] Y.-M. CHU, M.-K. WANG, *Inequalities between arithmetic-geometric, Gini, and Toader means*, Abstr. Appl. Anal. **2012** (2012), Article ID 830585, 11 pages.
- [21] Y.-M. CHU, M.-K. WANG, Y.-P. JIANG, S.-L. QIU, *Concavity of the complete elliptic integrals of the second kind with respect to Hölder means*, J. Math. Anal. Appl. **395**, 2 (2012), 637–642.
- [22] Y.-M. CHU, M.-K. WANG, X.-Y. MA, *Sharp bounds for Toader mean in terms of contraharmonic mean with applications*, J. Math. Inequal. **7**, 2 (2013), 161–166.
- [23] Y.-M. CHU, M.-K. WANG AND S.-L. QIU, *Optimal combinations bounds of root-square and arithmetic means for Toader mean*, Proc. Indian Acad. Sci. Math. Sci. **122**, (1) (2012), 41–51.
- [24] Y.-M. CHU, M.-K. WANG, S.-L. QIU, Y.-P. JIANG, *Bounds for complete elliptic integrals of the second kind with applications*, Comput. Math. Appl. **63**, 7 (2012), 1177–1184.
- [25] Y.-M. CHU, M.-K. WANG, Y.-F. QIU, X.-Y. MA, *Sharp two parameter bounds for the logarithmic mean and the arithmetic-geometric mean of Gauss*, J. Math. Inequal. **7**, 3 (2013), 349–355.
- [26] Y.-M. CHU, M.-K. WANG, S.-L. QIU, Y.-F. QIU, *Sharp generalized Seiffert mean bounds for Toader mean*, Abstr. Appl. Anal. **2011** (2011), Article ID 605259, 8 pages.
- [27] Y.-M. CHU, G.-D. WANG, X.-H. ZHANG, *Schur convexity and Hadamard's inequality*, Math. Inequal. Appl. **13**, 4 (2010), 725–731.
- [28] Y.-M. CHU, G.-D. WANG, X.-H. ZHANG, *The Schur multiplicative and harmonic convexities of the complete symmetric function*, Math. Nachr. **284**, 5–6 (2011), 653–663.
- [29] Y.-M. CHU, W.-F. XIA, *Two sharp inequalities for power mean, geometric mean, and harmonic mean*, J. Inequal. Appl. **2009** (2009), Article ID 741923, 6 pages.
- [30] Y.-M. CHU, W.-F. XIA, *Solution of an open problem for Schur convexity or concavity of the Gini mean values*, Sci. China **52A**, 10 (2009), 2099–2106.
- [31] Y.-M. CHU, W.-F. XIA, X.-H. ZHANG, *The Schur concavity, Schur multiplicative and harmonic convexities of the second dual form of the Hamy symmetric function with applications*, J. Multivariate Anal. **105** (2012), 412–421.
- [32] Y.-M. CHU, W.-F. XIA, T.-H. ZHAO, *Schur convexity for a class of symmetric functions*, Sci. China Math. **53**, 2 (2010), 465–474.
- [33] Y.-M. CHU, X.-M. ZHANG, *Necessary and sufficient conditions such that extended mean values are Schur-convex or Schur-concave*, J. Math. Kyoto Univ. **48**, 1 (2008), 229–238.

- [34] Y.-M. CHU, X.-M. ZHANG, G.-D. WANG, *The Schur geometrical convexity of the extended mean values*, J. Convex Anal. **15**, 4 (2008), 707–718.
- [35] Y.-M. CHU, T.-H. ZHAO, *Convexity and concavity of the complete elliptic integrals with respect to Lehmer mean*, J. Inequal. Appl. **2015** (2015), Article 396, 6 pages.
- [36] Y.-M. CHU, T.-H. ZHAO, *Concavity of the error function with respect to Hölder means*, Math. Inequal. Appl. **19**, 2 (2016), 589–595.
- [37] V. HEIKKALA, H. LINDÉN, M. K. VAMANAMURTHY, M. VUORINEN, *Generalized elliptic integrals and the Legendre \mathcal{M} -function*, J. Math. Anal. Appl. **338**, 1 (2008), 223–243.
- [38] T.-R. HUANG, B.-W. HAN, X.-Y. MA, Y.-M. CHU, *Optimal bounds for the generalized Euler-Mascheroni constant*, J. Inequal. Appl. **2018** (2018), Article 118, 9 pages.
- [39] E. A. KARATSUBA, M. VUORINEN, *On hypergeometric functions and generalizations of Legendre's relation*, J. Math. Anal. Appl. **260**, 2 (2001), 623–640.
- [40] Y.-M. LI, W.-F. XIA, Y.-M. CHU, X.-H. ZHANG, *Optimal lower and upper bounds for the geometric convex combination of the error function*, J. Inequal. Appl. **2015** (2015), Article 382, 8 pages.
- [41] W.-M. QIAN, Y.-M. CHU, *Sharp bounds for a special quasi-arithmetic mean in terms of arithmetic and geometric means with two parameters*, J. Inequal. Appl. **2017** (2017), Article 274, 10 pages.
- [42] W.-M. QIAN, X.-H. ZHANG, Y.-M. CHU, *Sharp bounds for the Toader-Qi mean in terms of harmonic and geometric means*, J. Math. Inequal. **11**, 1 (2017), 121–127.
- [43] S.-L. QIU, M. VUORINEN, *Landen inequalities for hypergeometric functions*, Nagoya Math. J. **154** (1999), 31–56.
- [44] S.-L. QIU, M. VUORINEN, *Infinite products and the normalized quotients of hypergeometric functions*, SIAM J. Math. Anal. **30**, 5 (1999), 1057–1075.
- [45] S.-L. QIU, M. VUORINEN, *Duplication inequalities for the ratios of hypergeometric functions*, Forum Math. **12**, 1 (2000), 109–133.
- [46] S.-L. QIU, M. VUORINEN, *Special functions in geometric function theory*, In: Handbook of Complex Analysis: geometric function theory, 2, 621–659, Elsevier Sci. B. V., Amsterdam, 2005.
- [47] S. SIMIĆ, M. VUORINEN, *Landen inequalities for zero-balanced hypergeometric functions*, Abstr. Appl. Anal. **2012** (2012), Article ID 932061, 11 pages.
- [48] Y.-Q. SONG, M. ADIL KHAN, S. ZAHEER ULLAH, Y.-M. CHU, *Integral inequalities involving strongly convex functions*, J. Funct. Spaces **2018** (2018), Article ID 6595921, 8 pages.
- [49] Y.-Q. SONG, P.-G. ZHOU, Y.-M. CHU, *Inequalities for the Gaussian hypergeometric function*, Sci. China Math. **57**, 11 (2014), 2369–2380.
- [50] M.-K. WANG AND Y.-M. CHU, *Asymptotical bounds for complete elliptic integrals of the second kind*, J. Math. Anal. Appl. **402**, 1 (2013), 119–126.
- [51] M.-K. WANG AND Y.-M. CHU, *Refinements of transformation inequalities for zero-balanced hypergeometric functions*, Acta Math. Sci. **37B**, 3 (2017), 607–622.
- [52] M.-K. WANG, Y.-M. CHU, *Landen inequalities for a class of hypergeometric functions with applications*, Math. Inequal. Appl. **21**, 2 (2018), 521–537.
- [53] M.-K. WANG, Y.-M. CHU AND Y.-P. JIANG, *Ramanujan's cubic transformation inequalities for zero-balanced hypergeometric functions*, Rocky Mountain J. Math. **46**, 2 (2016), 679–691.
- [54] M.-K. WANG, Y.-M. CHU, Y.-P. JIANG, S.-L. QIU, *Bounds of the perimeter of an ellipse using arithmetic, geometric and harmonic means*, Math. Inequal. Appl. **17**, 1 (2014), 101–111.
- [55] M.-K. WANG, Y.-M. CHU AND S.-L. QIU, *Some monotonicity properties of generalized elliptic integrals with applications*, Math. Inequal. Appl. **16**, 3 (2013), 671–677.
- [56] M.-K. WANG, Y.-M. CHU AND S.-L. QIU, *Sharp bounds for generalized elliptic integrals of the first kind*, J. Math. Anal. Appl. **429**, 2 (2015), 744–757.
- [57] M.-K. WANG, Y.-M. CHU, S.-L. QIU, Y.-P. JIANG, *Bounds for the perimeter of an ellipse*, J. Approx. Theory **164**, 7 (2012), 928–937.
- [58] M.-K. WANG, Y.-M. CHU, S.-L. QIU AND Y.-P. JIANG, *Convexity of the complete elliptic integrals of the first kind with respect to Hölder means*, J. Math. Anal. Appl. **388**, 2 (2012), 1141–1146.
- [59] M.-K. WANG, Y.-M. CHU, Y.-F. QIU, S.-L. QIU, *An optimal power mean inequalities for the complete elliptic integrals*, Appl. Math. Lett. **24**, 6 (2011), 887–890.
- [60] M.-K. WANG, Y.-M. CHU, Y.-Q. SONG, *Asymptotical formulas for Gaussian and generalized hypergeometric functions*, Appl. Math. Comput. **276** (2016), 44–60.
- [61] M.-K. WANG, Y.-M. LI, Y.-M. CHU, *Inequalities and infinite product formula for Ramanujan generalized modular equation function*, Ramanujan J. **46**, 1 (2018), 189–200.

- [62] H. WANG, W.-M. QIAN, Y.-M. CHU, *Optimal bounds for Gaussian arithmetic-geometric mean with applications to complete elliptic integral*, J. Funct. Spaces **2016** (2016), Article ID 3698463, 6 pages.
- [63] M.-K. WANG, Y.-F. QIU, Y.-M. CHU, *Sharp bounds for Seiffert means in terms of Lehmer means*, J. Math. Inequal. **4**, 4 (2010), 581–586.
- [64] M.-K. WANG, S.-L. QIU, Y.-M. CHU, *Infinite series formula for Hübner upper bound function with applications to Hersch-Pfluger distortion functions*, Math. Inequal. Appl. **21**, 3 (2018), 629–648.
- [65] M.-K. WANG, S.-L. QIU, Y.-M. CHU, Y.-P. JIANG, *Generalized Hersch-Pfluger distortion function and complete elliptic integrals*, J. Math. Anal. Appl. **385** (2012), 221–229.
- [66] M.-K. WANG, Z.-K. WANG, Y.-M. CHU, *An optimal double inequality between geometric and identity means*, Appl. Math. Lett. **25** (2012), 471–475.
- [67] G.-D. WANG, X.-H. ZHANG, Y.-M. CHU, *Inequalities for the generalized elliptic integrals and modular functions*, J. Math. Anal. Appl. **331**, 2 (2007), 1275–1283.
- [68] G.-D. WANG, X.-H. ZHANG, Y.-M. CHU, *A power mean inequality involving the complete elliptic integrals*, Rocky Mountain J. Math. **44**, 5 (2014), 1661–1667.
- [69] W.-F. XIA, Y.-M. CHU, *Optimal inequalities for the convex combination of error function*, J. Math. Inequal. **9**, 1 (2015), 85–99.
- [70] H.-Z. XU, Y.-M. CHU, W.-M. QIAN, *Sharp bounds for the Sándor-Yang means in terms of arithmetic and contra-harmonic means*, J. Inequal. Appl. **2018** (2018), Article 127, 13 pages.
- [71] ZH.-H YANG, *A new way to prove L'Hospital monotone rules with applications*, arXiv:1409.6408v1 [math. CA], available online at <http://arxiv.org/pdf/1409.6408v1.pdf>.
- [72] ZH.-H. YANG, Y.-M. CHU, *Asymptotic formulas for gamma function with applications*, Appl. Math. Comput. **270** (2015), 665–680.
- [73] ZH.-H. YANG, Y.-M. CHU, *A monotonicity property involving the generalized elliptic integral of the first kind*, Math. Inequal. Appl. **20**, 3 (2017), 729–735.
- [74] ZH.-H. YANG, Y.-M. CHU, M.-K. WANG, *Monotonicity criterion for the quotient of power series with applications*, J. Math. Anal. Appl. **428**, 1 (2015), 587–604.
- [75] ZH.-H. YANG, Y.-M. CHU, W. ZHANG, *Accurate approximations for the complete elliptic integral of the second kind*, J. Math. Anal. Appl. **438**, 2 (2016), 875–888.
- [76] ZH.-H. YANG, Y.-M. CHU, W. ZHANG, *Monotonicity of the ratio for the complete elliptic integral and Stolarsky mean*, J. Inequal. Appl. **2016** (2016), Article 176, 10 pages.
- [77] ZH.-H. YANG, Y.-M. CHU, W. ZHANG, *Sharp Stolarsky mean bounds for the complete elliptic integral of the second kind*, J. Nonlinear Sci. Appl. **10**, 3 (2017), 929–936.
- [78] ZH.-H. YANG, W.-M. QIAN, Y.-M. CHU, W. ZHANG, *Monotonicity rule for the quotient of two functions and its application*, J. Inequal. Appl. **2017** (2017), Article 106, 13 pages.
- [79] ZH.-H. YANG, W.-M. QIAN, Y.-M. CHU, W. ZHANG, *On rational bounds for the gamma function*, J. Inequal. Appl., **2017** (2017), Article 210, 17 pages.
- [80] ZH.-H. YANG, W.-M. QIAN, Y.-M. CHU, W. ZHANG, *On approximating the arithmetic-geometric mean and complete elliptic integral of the first kind*, J. Math. Anal. Appl. **462**, 2 (2018), 1714–1726.
- [81] ZH.-H. YANG, W.-M. QIAN, Y.-M. CHU, W. ZHANG, *On approximating the error function*, Math. Inequal. Appl. **21**, 2 (2018), 469–479.
- [82] ZH.-H. YANG, Y.-Q. SONG, Y.-M. CHU, *Sharp bounds for the arithmetic-geometric mean*, J. Inequal. Appl. **2014** (2014), Article 192, 13 pages.
- [83] ZH.-H. YANG, W. ZHANG, Y.-M. CHU, *Monotonicity and inequalities involving the incomplete gamma function*, J. Inequal. Appl. **2016** (2016), Article 221, 10 pages.
- [84] ZH.-H. YANG, W. ZHANG, Y.-M. CHU, *Monotonicity of the incomplete gamma function with applications*, J. Inequal. Appl. **2016** (2016), Article 251, 10 pages.
- [85] ZH.-H. YANG, W. ZHANG, Y.-M. CHU, *Sharp Gautschi inequality for parameter $0 < p < 1$ with applications*, Math. Inequal. Appl. **20**, 4 (2017), 1107–1120.
- [86] X.-H. ZHANG, *Solution to a conjecture on the Legendre M-function with an application to the generalized modulus*, J. Math. Anal. Appl. **431**, 2 (2015), 1190–1196.
- [87] X.-M. ZHANG, Y.-M. CHU, *A double inequality for gamma function*, J. Inequal. Appl. **2009** (2009), Article ID 503782, 7 pages.
- [88] X.-H. ZHANG, G.-D. WANG, Y.-M. CHU, *Convexity with respect to Hölder mean involving zero-balanced hypergeometric functions*, J. Math. Anal. Appl. **353**, 1 (2009), 256–259.
- [89] X.-H. ZHANG, G.-D. WANG, Y.-M. CHU, *Remarks on generalized elliptic integrals*, Proc. Roy. Soc. Edinburgh **139A**, 2 (2009), 417–426.

- [90] T.-H. ZHAO, Y.-M. CHU, *A class of logarithmically completely monotonic functions associated with gamma function*, J. Inequal. Appl. **2010** (2010), Article ID 392431, 11 pages.
- [91] T.-H. ZHAO, Y.-M. CHU, Y.-P. JIANG, *Monotonic and logarithmically convex properties of a function involving gamma functions*, J. Inequal. Appl. **2009** (2009), Article ID 728612, 13 pages.
- [92] T.-H. ZHAO, Y.-M. CHU, H. WANG, *Logarithmically complete monotonicity properties relating to the gamma function*, Abstr. Appl. Anal. **2011** (2011), Article ID 896483, 13 pages.

(Received March 4, 2018)

Miao-Kun Wang
 College of Science
 Hunan City University
 Yiyang 413000, P. R. China
e-mail: wmk000@126.com, wangmiaokun@zjhu.edu.cn

Yu-Ming Chu
 Department of Mathematics
 Huzhou University
 Huzhou 313000, P. R. China
e-mail: chuyuming2005@126.com

Wen Zhang
 Friedman Brain Institute
 Icahn School of Medicine at Mount Sinai
 New York NY 10029, USA
e-mail: zhang.wen81@gmail.com