

A CRITERION FOR THE CONVERSE OF THE HERMITE–HADAMARD INEQUALITY ON SIMPLICES

ELEUTHERIUS SYMEONIDIS

Abstract. It is already known that if a function satisfies the left or the right Hermite-Hadamard inequality for all simplices in its domain of definition, then it is convex, provided that the density of the measure is continuous and does not vanish identically on any segment. Here we show that this condition can be relaxed.

Mathematics subject classification (2010): 26B25.

Keywords and phrases: Convexity, Hermite-Hadamard inequality, simplex.

REFERENCES

- [1] J. HADAMARD, *Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann*, Journ. de Math., (4) **IX** (1893), 171–215.
- [2] CH. HERMITE, *Sur deux limites d'une intégrale définie*, Mathesis, **3** (1883), 82.
- [3] F. - C. MITROI, E. SYMEONIDIS, *The converse of the Hermite-Hadamard inequality on simplices*, Expo. Math., **30** (2012), 389–396.
- [4] C. P. NICULESCU, L. - E. PERSSON, *Convex Functions and Their Applications. A Contemporary Approach. Second Edition*, CMS Books in Mathematics (Canadian Mathematical Society), Springer, 2018.
- [5] R. R. PHELPS, *Lectures on Choquet's Theorem*, D. van Nostrand Company Inc., Princeton, 1966.