

SOME GENERALIZATIONS AND COMPLEMENTS OF DETERMINANTAL INEQUALITIES

HASSANE ABBAS* AND MOHAMMAD M. GHABRIES

(Communicated by J.-C. Bourin)

Abstract. K. Audenaert in [1] formulated a determinantal inequality arising from diffusion tensor imaging. Very recently M. Lin proved in [6] a complement and proposed a conjecture. In this short note, we generalize his conjecture and we prove it in a wild case, when the matrix is singular. We also present a refinement of the complement found by Lin and finally we present a series of determinantal inequalities followed by a conjecture.

1. Introduction

Audenaert formulated in his work [1] the following inequality for all $A, B \geq 0$ of same size $n \geq 1$,

$$\det(A^2 + |BA|) \leq \det(A^2 + AB) \quad (1)$$

He proved it in order to get the following determinantal inequality arising from diffusion tensor imaging

$$\det(A + U^*B) \leq \det(A + B)$$

where A and B are two n -square positive semi-definite matrices and U is a specified n -square unitary matrix arising from the polar decomposition of the matrix BA . Throughout this paper, let M_n be the space of $n \times n$ complex matrices. I_n denotes the identity matrix in M_n . The modulus of a complex matrix X is the unique positive semi-definite square root of the X^*X denoted by $|X| = (X^*X)^{1/2}$. For $X, Y \in M_n$ Hermitian matrices we say $X \geq Y$ if $X - Y$ is positive semi-definite matrix. The spectrum of X is the multiset of the eigenvalues of X denoted by $Sp(X)$, we can simply rearrange the eigenvalues of X in decreasing order if they are all real, that is

$$\lambda_1(X) \geq \lambda_2(X) \geq \dots \geq \lambda_n(X).$$

For every $X \in M_n$ Hermitian we have $\lambda(X) = (\lambda_1(X), \lambda_2(X), \dots, \lambda_n(X))^t$ is a real vector of order n . The spectral norm of a $X \in M_n$ is defined by $\|X\|_{op} = \rho^{1/2}(X^*X)$ where

$$\rho(X) = \sup_{\lambda \in Sp(X)} |\lambda(X)|.$$

Mathematics subject classification (2010): 15A45, 15A60.

Keywords and phrases: Determinantal inequality, Hermitian matrix, positive definite matrix, log-majorization, eigenvalues.

* Corresponding author.

Let us recall some definitions of majorizations: for a vector $x \in \mathbb{R}^n$, the vector obtained after rearranging the components of x in decreasing order is denoted by $x = (x_1^\downarrow, x_2^\downarrow, \dots, x_n^\downarrow)^t$, we say $x \in \mathbb{R}^n$ is weakly log majorized by $y \in \mathbb{R}^n$ denoted by $x \prec_{w, \log} y$ if

$$\prod_{i=1}^k (x_i^\downarrow) \leq \prod_{i=1}^k (y_i^\downarrow) \quad k = 1, 2, \dots, n \tag{2}$$

and x is log majorized by y ($x \prec_{\log} y$) if (2) is true and equality holds for $k = n$.

M. Lin proved in [6] a complement and a generalization for (1),

$$\det(A^2 + |AB|) \geq \det(A^2 + AB) \tag{3}$$

and

$$\det(A^2 + |BA|^p) \leq \det(A^2 + A^p B^p) \quad 0 \leq p \leq 2 \tag{4}$$

And he also introduced the following conjecture.

CONJECTURE 1.1. Let A, B be two positive semi-definite matrices. Then

$$\det(A^2 + |AB|^p) \geq \det(A^2 + A^p B^p), \quad 0 \leq p \leq 2$$

In this paper, we will show determinantal inequalities that are inspired by (1), (3) and (4).

2. Main Results

The main result in this paper are the following:

1. Let A, B be two positive semi-definite matrices. Then

$$\det(A^{kp} + |BA|^p) \leq \det(A^{kp} + A^p B^p) \quad k \geq 1, \quad 0 \leq p \leq 2 \tag{5}$$

2. Let A, B be two positive semi-definite matrices. Then, for all $0 \leq p \leq 2$,

- $\det(A^p + |BA|^p) \leq \det(A^p + A^p B^p)$.
- $\det(I_n + |BA|^p) \geq \det(I_n + A^p B^p)$.

3. Let A, B be two positive semi-definite matrices. Then, for all $p \geq 2$,

- $\det(A^p + |BA|^p) \geq \det(A^p + A^p B^p)$.
- $\det(I_n + |BA|^p) \leq \det(I_n + A^p B^p)$.

4. Let A, B be two n -square hermitian matrices. Then

$$\det(A^4 + |AB|^2) \geq \det(A^4 + A^2 B^2).$$

5. Let A, B be two positive semi-definite matrices. Then for every $k \geq 1$

$$\det(A^k + |AB|) \geq \det(A^k + AB).$$

6. Let A, B be two positive semi-definite matrices. Then,

$$\det(A^2 + |BA|^2) = \det(A^2 + A^2B^2) \leq \det(A^2 + (AB)^2) \leq \det(A^2 + |AB|^2).$$

We remark that for two positive semi-definite matrices A and B with A singular, the following general result holds

$$\det(A^k + |BA|^p) = \det(A^k + A^pB^p) \quad \text{and} \quad \det(A^k + |AB|^p) \geq \det(A^k + A^pB^p)$$

for all $k > 0, p \geq 0$. Which gives a partial answer of the positivity of Lin’s conjecture.

We can find a generalization for (4), to do this we need the following lemmas where the proof of the first one is in [6] and the proof of the second is a corollary of Furuta’s inequality and it can be found in [3, p. 128] and the third lemma proved in the interesting reference [5].

LEMMA 1. *If $\lambda(A), \lambda(B) \in \mathbb{R}_+^n$ such that $\lambda(A) \prec_{w, \log} \lambda(B)$ then*

$$\det(I_n + A) \leq \det(I_n + B).$$

LEMMA 2. *Let A, B be two positive semi-definite matrices such that $A \geq B$. Then, for all $p \geq 1, r \geq 0$,*

$$A^{(p+2r)/p} \geq (A^rB^pA^r)^{1/p}.$$

LEMMA 3. *Let X and Y be two positive semi-definite matrices. Then for every unitarily invariant norm, we have*

$$|||X^tY^tX^t||| \leq |||(XYX)^t||| \quad 0 \leq t \leq 1 \tag{6}$$

and

$$|||X^tY^tX^t||| \geq |||(XYX)^t||| \quad t \geq 1 \tag{7}$$

The following theorem is one of our main result.

THEOREM 1. *Let A, B be two positive semi-definite matrices. Then for all $0 \leq p \leq 2$ and for all $k \geq 1$,*

$$\det(A^{kp} + |BA|^p) \leq \det(A^{kp} + A^pB^p) \tag{8}$$

Proof. It is enough to prove the result for $k > 1$ as the case $k = 1$ is followed by limit argument. Assume that A is invertible, for A singular the inequality is true. First we need to prove that, for all $0 \leq t \leq 1$ and $k > 1$, we have

$$\lambda(A^{kt/2}(A^{-1/2}BA^{-1/2})^t A^{kt/2}) \prec_{\log} \lambda(A^{(k-1)t}B^t). \tag{9}$$

To achieve (9), it is enough to show that

$$A^{\frac{(k-1)t}{2}}B^tA^{\frac{(k-1)t}{2}} \leq I_n \Rightarrow A^{kt/2}(A^{-1/2}BA^{-1/2})^t A^{kt/2} \leq I_n.$$

Assume that $A^{\frac{(k-1)t}{2}}B^tA^{\frac{(k-1)t}{2}} \leq I_n$, so $0 \leq B^t \leq A^{-(k-1)t}$ and by applying Lemma 2, we get

$$A^{-(k-1)t(p+2r)/p} \geq \left(A^{-(k-1)tr}B^{tp}A^{-(k-1)tr} \right)^{1/p}.$$

Now, by replacing p with $\frac{1}{t} \geq 1$ and r with $\frac{1}{2(k-1)t} > 0$ we obtain

$$A^{-(k-1)t^2\left(\frac{1}{t} + \frac{1}{(k-1)t}\right)} \geq \left(A^{-1/2}BA^{-1/2} \right)^t$$

which implies

$$A^{-kt} \geq \left(A^{-1/2}BA^{-1/2} \right)^t.$$

Therefore $A^{kt/2}(A^{-1/2}BA^{-1/2})^t A^{kt/2} \leq I_n$.

Let $a = \lambda_1\left(A^{\frac{(k-1)t}{2}}B^tA^{\frac{(k-1)t}{2}}\right)$. If $a = 0$, then it is obvious that (9) is true. If $a > 0$, we observe that

$$A^{\frac{(k-1)t}{2}}B^tA^{\frac{(k-1)t}{2}} \leq a I_n \text{ and } \left(\frac{1}{a^{1/kt}}A \right)^{\frac{(k-1)t}{2}} \left(\frac{1}{a^{1/kt}}B \right)^t \left(\frac{1}{a^{1/kt}}A \right)^{\frac{(k-1)t}{2}} \leq I_n.$$

This yields

$$\left(\frac{1}{a^{1/kt}}A \right)^{kt/2} \left[\left(\frac{1}{a^{1/kt}}A \right)^{-1/2} \left(\frac{1}{a^{1/kt}}B \right) \left(\frac{1}{a^{1/kt}}A \right)^{-1/2} \right]^t \left(\frac{1}{a^{1/kt}}A \right)^{kt/2} \leq I_n.$$

Thus

$$A^{kt/2}(A^{-1/2}BA^{-1/2})^t A^{kt/2} \leq a I_n.$$

And hence

$$\lambda_1\left(A^{kt/2}(A^{-1/2}BA^{-1/2})^t A^{kt/2}\right) \leq \lambda_1\left(A^{\frac{(k-1)t}{2}}B^tA^{\frac{(k-1)t}{2}}\right) \tag{10}$$

Now, using the antisymmetric tensor product, we have

$$\wedge^s(A^{kt/2}(A^{-1/2}BA^{-1/2})^t A^{kt/2}) = (\wedge^s A)^{kt/2} \left((\wedge^s A)^{-1/2}(\wedge^s B)(\wedge^s A)^{-1/2} \right)^t (\wedge^s A)^{kt/2}$$

and

$$\wedge^s \left(A^{\frac{(k-1)t}{2}} B^t A^{\frac{(k-1)t}{2}} \right) = (\wedge^s A)^{\frac{(k-1)t}{2}} (\wedge^s B)^t (\wedge^s A)^{\frac{(k-1)t}{2}} \text{ for } 1 \leq s \leq n.$$

Replacing A and B with $\wedge^s A$ and $\wedge^s B$ respectively in (10) yields

$$\lambda_1 \left(\wedge^s (A^{kt/2} (A^{-1/2} B A^{-1/2})^t A^{kt/2}) \right) \leq \lambda_1 \left(\wedge^s (A^{\frac{(k-1)t}{2}} B^t A^{\frac{(k-1)t}{2}}) \right).$$

And so, for all $1 \leq s \leq n - 1$, we have

$$\prod_{i=1}^s \lambda_i \left(A^{kt/2} (A^{-1/2} B A^{-1/2})^t A^{kt/2} \right) \leq \prod_{i=1}^s \lambda_i \left(A^{\frac{(k-1)t}{2}} B^t A^{\frac{(k-1)t}{2}} \right).$$

And as in general $\det(A^{kt/2} (A^{-1/2} B A^{-1/2})^t A^{kt/2}) = \det(A^{\frac{(k-1)t}{2}} B^t A^{\frac{(k-1)t}{2}})$, we obtain

$$\prod_{i=1}^n \lambda_i \left(A^{kt/2} (A^{-1/2} B A^{-1/2})^t A^{kt/2} \right) = \prod_{i=1}^n \lambda_i \left(A^{\frac{(k-1)t}{2}} B^t A^{\frac{(k-1)t}{2}} \right).$$

and by consequently,

$$\lambda(A^{kt/2} (A^{-1/2} B A^{-1/2})^t A^{kt/2}) \prec_{\log} \lambda(A^{(k-1)t} B^t).$$

By applying Lemma 1, we get

$$\det(I_n + A^{kt/2} (A^{-1/2} B A^{-1/2})^t A^{kt/2}) \leq \det(I_n + A^{(k-1)t} B^t).$$

Now taking $A = A^{-2}$, $B = B^2$ and $t = p/2$ yields

$$\det(I_n + A^{-kp/2} (A B^2 A)^{p/2} A^{-kp/2}) \leq \det(I_n + A^{-kp+p} B^p) \tag{11}$$

Pre-post multiplying both sides of (11) by $\det(A^{kp/2})$ leads to the result for $k > 1$ and $0 < p \leq 2$. Finally, it is easy to see that (8) is true for $p = 0$. \square

The next theorem shows some reverse inequalities.

THEOREM 2. *Let A and B be two positive semi-definite matrices. Then for $0 \leq p \leq 2$,*

$$\det(I_n + |BA|^p) \geq \det(I_n + A^p B^p) \tag{12}$$

Proof. Take $Y = B^2$, $X = A$ and the spectral norm in (6) gives

$$\lambda_1(B^t A^{2t} B^t) \leq \lambda_1((BA^2 B)^t) \quad 0 \leq t \leq 1 \tag{13}$$

Note that $\wedge^s(A^t B^{2t} A^t) = (\wedge^s A)^t (\wedge^s B)^{2t} (\wedge^s A)^t$ and $\wedge^s((AB^2 A)^t) = (\wedge^s A (\wedge^s B)^2 \wedge^s A)^t$.

Replacing A and B with $\wedge^s A$ and $\wedge^s B$ respectively in (13) yields

$$\lambda_1(\wedge^s(A^t B^{2t} A^t)) \leq \lambda_1(\wedge^s((AB^2A)^t)).$$

And as $\det((AB^2A)^t) = \det(A^t B^{2t} A^t)$, then for $0 \leq t \leq 1$

$$\lambda(A^t B^{2t} A^t) \prec_{\log} \lambda((AB^2A)^t).$$

Assume that A is positive definite matrix. For $t = p/2$ and by Lemma 1 we get

$$\det(I_n + A^{p/2} B^p A^{p/2}) \leq \det(I_n + |BA|^p)$$

Therefore we get the desired for A positive semi-definite matrix by continuity argument. \square

With a similar proof using (7), we can get the following.

THEOREM 3. *Let A, B be two positive semi-definite matrices. Then for all $p \geq 2$,*

- $\det(A^p + |BA|^p) \geq \det(A^p + A^p B^p)$.
- $\det(I_n + |BA|^p) \leq \det(I_n + A^p B^p)$.

We can find a more general complement for (8) when $k = 2$ and $p = 2$ as the following theorem shows.

THEOREM 4. *Let A, B be two n -square hermitian matrices. Then*

$$\det(A^4 + |AB|^2) \geq \det(A^4 + A^2 B^2).$$

Proof. Again, assume that A is an invertible matrix, the case of A singular is true by continuity argument. It is well known, in [7, p. 352], that if a matrix $M = \begin{pmatrix} X & Y \\ Y^* & Z \end{pmatrix} \in \mathbb{M}_{n+n}(\mathbb{C})$ is positive semi-definite, then

$$|\lambda(Y)| \prec_{w,\log} \lambda^{\frac{1}{2}}(X) \circ \lambda^{\frac{1}{2}}(Z).$$

where $X \circ Y$ represents the Hadamard product of the two matrices X and Y .

Replacing $X = BA^{-2}B$, $Y = B^2A^{-2}$ and $Z = A^{-2}BA^2BA^{-2}$, and by using Schur's complement we get $M = \begin{pmatrix} BA^{-2}B & B^2A^{-2} \\ A^{-2}B^2 & A^{-2}BA^2BA^{-2} \end{pmatrix} \geq 0$. Thus

$$\| \|B^2A^{-2}\| \|_{op}^2 \leq \| \|BA^{-2}B\| \|_{op} \cdot \| \|A^{-2}BA^2BA^{-2}\| \|_{op}.$$

If $\| \|B^2A^{-2}\| \|_{op} = 0$, then the desired determinantal inequality is true.

Noticing that $|||B^2A^{-2}|||_{op} = |||BA^{-2}B|||_{op}$, and dividing both sides by $|||B^2A^{-2}|||_{op} > 0$ gives

$$|||A^{-1}B^2A^{-1}|||_{op} = |||B^2A^{-2}|||_{op} \leq |||A^{-2}BA^2BA^{-2}|||_{op}$$

which is

$$\lambda_1(A^{-1}B^2A^{-1}) \leq \lambda_1(A^{-2}BA^2BA^{-2}). \tag{14}$$

Observe that $\wedge^s(A^{-1}B^2A^{-1}) = (\wedge^s A)^{-1}(\wedge^s B)^2(\wedge^s A)^{-1}$,

$$\wedge^s(A^{-2}BA^2BA^{-2}) = (\wedge^s A)^{-2}(\wedge^s B)(\wedge^s A)^2(\wedge^s B)(\wedge^s A)^{-2}$$

and replacing A with $\wedge^s A$ and B with $\wedge^s B$ in (14) yields

$$\lambda_1(\wedge^s(A^{-1}B^2A^{-1})) \leq \lambda_1(\wedge^s(A^{-2}BA^2BA^{-2})), \quad 1 \leq s \leq n-1.$$

Also as $\det(A^{-1}B^2A^{-1}) = \det(A^{-2}BA^2BA^{-2})$ we obtain

$$\lambda(A^{-1}B^2A^{-1}) \prec_{\log} \lambda(A^{-2}BA^2BA^{-2}).$$

Using Lemma 1 gives

$$\det(I_n + A^{-1}B^2A^{-1}) \leq \det(I_n + A^{-2}BA^2BA^{-2}).$$

Pre-post multiplying by $\det(A^2)$ both sides yields

$$\det(A^4 + A^2B^2) \leq \det(A^4 + BA^2B) = \det(A^4 + |AB|^2). \quad \square$$

We may ask whether the following conjecture is true

CONJECTURE 2.1. Let A and B be two positive semi-definite matrices. Then

$$\det(A^k + A^2B^2) \leq \det(A^k + (AB)^2) \text{ for all } k \geq 1 \tag{15}$$

If (15) is true we get

$$\det(A^{k'} + A^2B^2) \leq \det(A^{k'} + |AB|^2) \text{ for all } k' \geq 1 \tag{16}$$

Also, if (16) is true then (15) is true.

The inequality (15) is true for $k = 1, 3$

- When $k = 1$ we have

$$\begin{aligned} \det(A + (AB)^2) &= \det(A) \cdot \det(I_n + BAB) \\ &= \det(A) \cdot \det(I_n + AB^2) \\ &= \det(A + A^2B^2) \end{aligned}$$

- When $k = 3$ we have

$$\begin{aligned}\det(A^3 + (AB)^2) &= \det(A) \cdot \det((A^{1/2})^4 + |A^{1/2}B|^2) \\ &\geq \det(A) \cdot \det((A^{1/2})^4 + (A^{1/2})^2 B^2) \quad (\text{using Theorem 4}) \\ &= \det(A^3 + A^2 B^2)\end{aligned}$$

The inequality (15) is not valid for $k < 1$ as the following example shows.

EXAMPLE 1. For $A = \begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}$, $B = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ and $k = 1/2$, we have

$$\det(A^{1/2} + A^2 B^2) = 60 > \det(A^{1/2} + (AB)^2) = 54.$$

Acknowledgement. The authors sincerely thank Prof. J.C.Bourin for his valuable suggestions and many useful comments which have helped to improve the paper and clarify the details. Also many thanks go to the Editor and Editor-in-Chief for giving us a chance to revise. The authors acknowledge financial support from the Lebanese University research grants program.

REFERENCES

- [1] K. M. R. AUDENAERT, *A determinantal inequality for the geometric mean with an application in diffusion tensor*, arXiv 1502.06902v2, 2015.
- [2] R. BHATIA, *Matrix Analysis*, Springer-Verlag, New York, 1997.
- [3] R. BHATIA, *Positive Definite Matrices*, Princeton University Press, New Jersey, 2007.
- [4] J. C. BOURIN, E.-Y. LEE AND M. LIN, *Positive matrices partitioned into a small number of Hermitian blocks*, Linear Algebra Appl., 483, 2013, 2591–2598.
- [5] J. C. BOURIN, E.-Y. LEE, *Matrix inequalities from a two variables functional*, Internat. J. Math., 27, 9, 2016, 19 pp.
- [6] M. LIN, *On a determinantal inequality arising from diffusion tensor imaging*, Commun. Contemp. Math, 19, 1650044, 2017, 6 pp.
- [7] F. ZHANG, *Matrix Theory: Basic Results and Techniques*, Springer, New York, 2nd ed., 2011.

(Received October 29, 2018)

Hassane Abbas
Department of Mathematics, Faculty of science
KALMA Laboratory, Lebanese University
Hadat-Beirut, Lebanon
e-mail: habbas@ul.edu.lb

Mohammad M. Ghabries
Department of Mathematics, Faculty of science
KALMA Laboratory, Lebanese University
Hadat-Beirut, Lebanon
e-mail: mahdi.ghabries@gmail.com