

OPTIMAL BOUNDS OF THE ARITHMETIC MEAN IN TERMS OF NEW SEIFFERT-LIKE MEANS

MONIKA NOWICKA AND ALFRED WITKOWSKI

Abstract. We provide the optimal bounds for the arithmetic mean in terms of the arithmetic, harmonic, quadratic, inverse quadratic and geometric combination of means generated by sine, tangent, hyperbolic sine and hyperbolic tangent functions.

Mathematics subject classification (2010): 26D15.

Keywords and phrases: Sine mean, tangent mean, Seiffert function, hyperbolic sine mean, hyperbolic tangent mean.

REFERENCES

- [1] G. D. ANDERSON, M. K. VAMANAMURTHY AND M. K. VOURINEN, *Conformal invariants, inequalities, and quasiconformal maps*, Canadian Mathematical Society Series of Monographs and Advanced Texts. John Wiley & Sons, Inc., New York, 1997.
- [2] I. LAZAREVIĆ, *Certain inequalities with hyperbolic functions*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. No. 159–170, (1966) 41–48 (in Serbo-Croatian).
- [3] D. S. MITRINović AND D. D. ADAMoviĆ, *Sur une inégalité élémentaire où interviennent des fonctions trigonométriques*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. No. 143–155, (1965) 23–34 (in French).
- [4] I. PINELIS, *L'Hospital type results for monotonicity, with applications*, JIPAM. J. Inequal. Pure Appl. Math. **3** 1 (2002), Article 5.
- [5] A. WITKOWSKI, *On Seiffert-like means*, J. Math. Inequal. 2015; **9** 4 (2015), 1071–1092. doi: 10.7153/jmi-09-83.