

ASYMPTOTIC EXPANSION AND BOUNDS FOR COMPLETE ELLIPTIC INTEGRALS

MIAO-KUN WANG, YU-MING CHU*, YONG-MIN LI AND WEN ZHANG

(Communicated by J. Pečarić)

Abstract. In the article, we present several new bounds for the the complete elliptic integrals $\mathcal{K}(r) = \int_0^{\pi/2} (1-r^2 \sin^2 \theta)^{-1/2} d\theta$ and $\mathcal{E}(r) = \int_0^{\pi/2} (1-r^2 \sin^2 \theta)^{1/2} d\theta$, and find an asymptotic expansion for $\mathcal{K}(r)$ as $r \rightarrow 1$, which are the refinements and improvements of the previously well-known results.

1. Introduction

For $r \in [0, 1]$, Lengedre's complete elliptic integrals of the first and second kind [18, 20, 21, 22, 37, 60, 72, 73, 74, 82, 83, 87] are defined by

$$\mathcal{K} = \mathcal{K}(r) = \int_0^{\pi/2} (1-r^2 \sin^2 \theta)^{-1/2} d\theta, \quad \mathcal{K}' = \mathcal{K}'(r) = \mathcal{K}(r'),$$

$$\mathcal{K}(0) = \pi/2, \quad \mathcal{K}(1) = \infty,$$

and

$$\mathcal{E} = \mathcal{E}(r) = \int_0^{\pi/2} (1-r^2 \sin^2 \theta)^{1/2} d\theta, \quad \mathcal{E}' = \mathcal{E}'(r) = \mathcal{E}(r'),$$

$$\mathcal{E}(0) = \pi/2, \quad \mathcal{E}(1) = 1,$$

respectively. Here and in what follows we set $r' = \sqrt{1-r^2}$. These integrals are special cases of Gaussian hypergeometric function [43, 55, 56, 57, 58, 62, 63, 66, 79, 91, 92, 94, 95]

$$F(a, b; c; x) = {}_2F_1(a, b; c; x) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{x^n}{n!} \quad (-1 < x < 1),$$

Mathematics subject classification (2010): 33E05, 26E60.

Keywords and phrases: Asymptotic expansion, complete elliptic integrals, arithmetic-geometric mean, infinite series, bound.

* Corresponding author.

where $(a)_0 = 1$, $(a)_n = \prod_{k=0}^{n-1} (a+k) = \Gamma(n+a)/\Gamma(a)$ for $n \geq 1$ and $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$ is the Euler gamma function [29, 30, 46, 49, 50, 65, 78, 84, 85, 86, 89, 93]. Indeed, we have

$$\mathcal{K}(r) = \frac{\pi}{2} F\left(\frac{1}{2}, \frac{1}{2}; 1; r^2\right) \quad \text{and} \quad \mathcal{E}(r) = \frac{\pi}{2} F\left(-\frac{1}{2}, \frac{1}{2}; 1; r^2\right).$$

It is well known that the complete elliptic integrals have many important applications in physics, engineering, geometric function theory, theory of mean values, number theory and other related fields [1, 4, 7, 10, 31, 34, 35, 48, 52, 75]. Particularly, in the past two decades, a great deal of mathematical effort in the complete elliptic integrals has been devoted to the study of distortion estimates in quasiconformal mappings [24, 25, 26, 27, 28, 32, 69, 70]. For more properties and recent development of \mathcal{K} and \mathcal{E} see the book [13], and the research articles [23, 36, 61, 64, 68, 81, 88].

1.1. The complete elliptic integrals and their related means

An important method to study the complete elliptic integrals is to consider their convexity and monotonicity properties, and their related bivariate means [2, 3, 5, 6, 8, 9, 16, 33, 39, 47, 76, 77, 90]. In 1875, Lagrange firstly defined the arithmetic-geometric mean $AG(a, b)$ [19, 38, 42, 45, 71, 80] of two positive real numbers a and b as the common limit of the following sequences $\{a_n\}$ and $\{b_n\}$:

$$\begin{aligned} a_0 &= a, & b_0 &= b, \\ a_{n+1} &= A(a_n, b_n) = \frac{a_n + b_n}{2}, & b_{n+1} &= G(a_n, b_n) = \sqrt{a_n b_n}. \end{aligned}$$

Later, Gauss proved that

$$AG(a, b) = \frac{\pi/2}{\int_0^{\pi/2} \frac{d\theta}{\sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta}}} = \begin{cases} \pi a / \left[2 \mathcal{K} \left(\sqrt{1 - (b/a)^2} \right) \right], & a \geq b, \\ \pi b / \left[2 \mathcal{K} \left(\sqrt{1 - (a/b)^2} \right) \right], & a < b. \end{cases}$$

From then on, $AG(a, b)$ and related iterations have been used for efficient numerical evaluation of the complete elliptic integral $\mathcal{K}(r)$.

In 1998, a class of quasi-arithmetic mean related to the arithmetic-geometric mean was introduced by Toader in [53], which is defined by

$$M_{p,n}(a, b) = p^{-1} \left(\frac{1}{\pi} \int_0^\pi p(r_n(\theta)) d\theta \right) = p^{-1} \left(\frac{2}{\pi} \int_0^{\pi/2} p(r_n(\theta)) d\theta \right),$$

where $r_n(\theta) = (a^n \cos^2 \theta + b^n \sin^2 \theta)^{1/n}$ for $n \neq 0$, and $r_0(\theta) = a^{\cos^2 \theta} b^{\sin^2 \theta}$, and p is a strictly monotonic function. In the particular case $p = x^{-1}$ and $n = 2$, $M_{p,n}(a, b)$ reduces to $AG(a, b)$. Moreover, taking $p = \sqrt{x}$ and $n = 1$, a new symmetric mean $E(a, b)$ [17, 40, 41, 67, 96] involving the complete elliptic integral of the second kind

was obtained

$$E(a, b) = \begin{cases} 4a \left[\mathcal{E} \left(\sqrt{1 - b/a} \right) \right]^2 / \pi^2, & a \geq b, \\ 4b \left[\mathcal{E} \left(\sqrt{1 - a/b} \right) \right]^2 / \pi^2, & a < b, \end{cases}$$

which will be studied and derived several asymptotic bounds for \mathcal{E} in this paper.

1.2. Some well known bounds for \mathcal{K} and \mathcal{E}

Ramanujan's work on the asymptotic behavior of the hypergeometric functions shows that $\mathcal{K}(r)$ satisfies (see [13, 1.48])

$$\mathcal{K}(r) + \log r' = \log 4 + O((1 - r^2) \log(1 - r^2)), \quad r \rightarrow 1. \quad (1.1)$$

Anderson, Vamanamurthy and Vuorinen [12] approximated $\mathcal{K}(r)$ by the inverse hyperbolic tangent function arth , obtaining the inequalities

$$\frac{\pi}{2} \left(\frac{\operatorname{arth} r}{r} \right)^{1/2} < \mathcal{K}(r) < \frac{\pi}{2} \frac{\operatorname{arth} r}{r} \quad (1.2)$$

for all $r \in (0, 1)$. In 1995, Sándor [51] refined the elegant inequalities for $AG(a, b)$ in terms of arithmetic and the logarithmic means

$$\frac{a - b}{\log a - \log b} = L(a, b) < AG(a, b) < A(a, b) = \frac{a + b}{2},$$

or equivalently

$$\frac{\pi}{2} \frac{1}{A(1, r')} < \mathcal{K}(r) < \frac{\pi}{2} \frac{1}{L(1, r')}$$

and proved that for $r \in (0, 1)$

$$\frac{\pi}{2} \left(\frac{2/\pi}{A(1, r')} + \frac{1 - 2/\pi}{L(1, r')} \right) < \mathcal{K}(r) < \frac{\pi}{2} \left(\frac{12/(5\pi)}{A(1, r')} + \frac{1 - 12/(5\pi)}{L(1, r')} \right). \quad (1.3)$$

In 2004, Alzer and Qiu [11] refined (1.2) and (1.3) as

$$\frac{\pi}{2} \left(\frac{\operatorname{arth} r}{r} \right)^{3/4} < \mathcal{K}(r) < \frac{\pi}{2} \frac{\operatorname{arth} r}{r} \quad (1.4)$$

with the best possible exponents $3/4$ and 1 , and

$$\frac{\pi}{2} \left(\frac{\alpha}{A(1, r')} + \frac{1 - \alpha}{L(1, r')} \right) < \mathcal{K}(r) < \frac{\pi}{2} \left(\frac{\beta}{A(1, r')} + \frac{1 - \beta}{L(1, r')} \right) \quad (1.5)$$

with the optimal parameters $\alpha = 2/\pi$ and $\beta = 3/4$.

In 1997, Vuorinen [54] conjectured that the inequality

$$\mathcal{E}(r) > \frac{\pi}{2} \left(\frac{1 + r'^{3/2}}{2} \right)^{2/3} \quad (1.6)$$

holds for all $r \in (0, 1)$. Later, Barnard et al. [14, 15] verified the conjecture. And in [11] the authors also provided an optimal upper bound for $\mathcal{E}(r)$ in terms of power mean of 1 and r' , namely, for $0 < r < 1$,

$$\mathcal{E}(r) < \frac{\pi}{2} \left(\frac{1 + r'^{\log 2 / \log(\pi/2)}}{2} \right)^{\log(\pi/2) / \log 2}. \tag{1.7}$$

In this paper, we shall give an asymptotic expansion for $\mathcal{K}(r)$ as $r \rightarrow 1$ in Section 2, which is a refinement of (1.1). In Section 3, several sharp symmetrical bounds for $AG(a, b)$ and $E(a, b)$ are found. These results lead to several new bounds for $\mathcal{K}(r)$ and $\mathcal{E}(r)$ in Section 4. Finally, in Section 5 we shall compare our results with (1.4)-(1.7).

In order to prove our main results we need some formulas [13, Appendix E, pp. 474-475] and a technical lemma, which we present in this section.

$$\frac{d\mathcal{K}}{dr} = \frac{\mathcal{E} - r'^2 \mathcal{K}}{rr'^2}, \quad \frac{d\mathcal{E}}{dr} = \frac{\mathcal{E} - \mathcal{K}}{r}, \quad \frac{d(\mathcal{E} - r'^2 \mathcal{K})}{dr} = r\mathcal{K},$$

$$\mathcal{K} \left(\frac{2\sqrt{r}}{1+r} \right) = (1+r)\mathcal{K}(r), \quad \mathcal{E} \left(\frac{2\sqrt{r}}{1+r} \right) = \frac{2\mathcal{E} - r'^2 \mathcal{K}}{1+r}.$$

LEMMA 1.1. [13, Theorem 1.25]. For $-\infty < a < b < \infty$, let $f, g : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$, and be differentiable on (a, b) , let $g'(x) \neq 0$ on (a, b) . If $f'(x)/g'(x)$ is increasing (decreasing) on (a, b) , then so are

$$\frac{f(x) - f(a)}{g(x) - g(a)} \quad \text{and} \quad \frac{f(x) - f(b)}{g(x) - g(b)}.$$

If $f'(x)/g'(x)$ is strictly monotone, then the monotonicity in the conclusion is also strict.

2. An asymptotic expansion for $\mathcal{K}(r)$ as $r \rightarrow 1$

In this section, we find an infinite series for the function $r \rightarrow \mathcal{K}(r) + \log r'$ defined on $(0, 1]$. This result yields an asymptotic expansion for $\mathcal{K}(r)$ as $r \rightarrow 1$, which is a refinement of (1.1).

THEOREM 2.1. Let $r \in (0, 1]$, $r_0 = r$, $r_1 = 2\sqrt{r}/(1+r) = \varphi_2(r_0)$, $r_2 = 2\sqrt{r_1}/(1+r_1) = \varphi_2(r_1) = \varphi_4(r_0), \dots, r_n = 2\sqrt{r_{n-1}}/(1+r_{n-1}) = \varphi_2(r_{n-1}) = \varphi_{2^n}(r_0)$. Then

$$\mathcal{K}(r) + \log r' = \frac{1}{2} \sum_{k=0}^{\infty} \frac{(3+r_k) \log(1+r_k) - (1-r_k) \log(1-r_k)}{\prod_{i=0}^k (1+r_i)}.$$

Proof. By simple computation we get

$$\mathcal{K}(r_1) + \log r'_1 = \mathcal{K} \left(\frac{2\sqrt{r}}{1+r} \right) + \log \left(\frac{1-r}{1+r} \right) = (1+r)\mathcal{K}(r) + \log \left(\frac{1-r}{1+r} \right)$$

$$\begin{aligned}
&= (1+r)[\mathcal{K}(r) + \log r'] - \frac{1+r}{2} \log(1-r^2) + \log\left(\frac{1-r}{1+r}\right) \\
&= (1+r)[\mathcal{K}(r) + \log r'] - \frac{(3+r)\log(1+r) - (1-r)\log(1-r)}{2},
\end{aligned}$$

that is,

$$\frac{\mathcal{K}(r_1) + \log r'_1}{1+r_0} = \mathcal{K}(r_0) + \log r'_0 - \frac{(3+r_0)\log(1+r_0) - (1-r_0)\log(1-r_0)}{2(1+r_0)}. \quad (2.1)$$

Putting r_{n-1} into (2.1) instead of r_0 , one has

$$\begin{aligned}
&\frac{\mathcal{K}(r_n) + \log r'_n}{1+r_{n-1}} \\
&= \mathcal{K}(r_{n-1}) + \log r'_{n-1} - \frac{(3+r_{n-1})\log(1+r_{n-1}) - (1-r_{n-1})\log(1-r_{n-1})}{2(1+r_{n-1})}. \quad (2.2)
\end{aligned}$$

It follows from (2.1) and (2.2) that

$$\begin{aligned}
&\mathcal{K}(r) + \log r' - \frac{(3+r_0)\log(1+r_0) - (1-r_0)\log(1-r_0)}{2(1+r_0)} \\
&\quad - \frac{(3+r_1)\log(1+r_1) - (1-r_1)\log(1-r_1)}{2(1+r_0)(1+r_1)} \\
&= \frac{\mathcal{K}(r_2) + \log r'_2}{(1+r_0)(1+r_1)},
\end{aligned}$$

and

$$\begin{aligned}
&\mathcal{K}(r) + \log r' - \frac{(3+r_0)\log(1+r_0) - (1-r_0)\log(1-r_0)}{2(1+r_0)} \\
&\quad - \frac{(3+r_1)\log(1+r_1) - (1-r_1)\log(1-r_1)}{2(1+r_0)(1+r_1)} \\
&\quad - \frac{(3+r_2)\log(1+r_2) - (1-r_2)\log(1-r_2)}{2(1+r_0)(1+r_1)(1+r_2)} \\
&= \frac{\mathcal{K}(r_3) + \log r'_3}{(1+r_0)(1+r_1)(1+r_2)}.
\end{aligned}$$

Generally, by recursion we can obtain

$$\mathcal{K}(r) + \log r' - \frac{1}{2} \sum_{k=0}^n \frac{(3+r_k)\log(1+r_k) - (1-r_k)\log(1-r_k)}{\prod_{i=0}^k (1+r_i)} = \frac{\mathcal{K}(r_{n+1}) + \log r'_{n+1}}{\prod_{i=0}^n (1+r_i)}.$$

Since the function $r \rightarrow \mathcal{K}(r) + \log r'$ is strictly decreasing from $(0, 1]$ onto $[\log 4, \pi/2)$,

$$0 < \mathcal{K}(r) + \log r' - \frac{1}{2} \sum_{k=0}^n \frac{(3+r_k)\log(1+r_k) - (1-r_k)\log(1-r_k)}{\prod_{i=0}^k (1+r_i)} < \frac{\pi/2}{\prod_{i=0}^n (1+r_i)}.$$

Letting $n \rightarrow \infty$, then $\prod_{i=0}^{\infty} (1 + r_i) = +\infty$ for $r \in (0, 1]$, and thereby

$$\mathcal{K}(r) + \log r' = \frac{1}{2} \sum_{k=0}^{\infty} \frac{(3 + r_k) \log(1 + r_k) - (1 - r_k) \log(1 - r_k)}{\prod_{i=0}^k (1 + r_i)}.$$

COROLLARY 2.2. *Let r_n be defined as in Theorem 2.1. Then the complete elliptic integrals of the first kind $\mathcal{K}(r)$ has the following asymptotic formula*

$$\mathcal{K}(r) + \log r' = \log 4 - \sum_{k=0}^{\infty} \frac{(1 - r_k) \log(1 - r_k)}{2^{k+2}}, \quad r \rightarrow 1.$$

3. Bounds for $AG(a, b)$ and $E(a, b)$

In this section, we shall prove several sharp symmetrical bounds for $AG(a, b)$ and $E(a, b)$ in terms of some classical mean values, such as arithmetic mean $A(a, b)$, geometric mean $G(a, b)$, harmonic mean $H(a, b)$ and logarithmic mean $L(a, b)$. We first establish three monotonicity lemmas involving the complete elliptic integrals $\mathcal{K}(r)$ and $\mathcal{E}(r)$.

LEMMA 3.1. (1) *The function $r \mapsto (\mathcal{E} - r'^2 \mathcal{K})/r^2$ is strictly increasing from $(0, 1)$ onto $(\pi/4, 1)$;*

(2) *the function $r \mapsto 2\mathcal{E} - r'^2 \mathcal{K}$ is strictly increasing from $(0, 1)$ onto $(\pi/2, 2)$;*

(3) *the function $r \mapsto [\mathcal{K} - \mathcal{E} - (\mathcal{E} - r'^2 \mathcal{K})]/r^4$ is strictly increasing from $(0, 1)$ onto $(\pi/16, \infty)$;*

(4) *the function $r \mapsto r'^{3/4}(\mathcal{K} - \mathcal{E})/r^2$ is strictly decreasing from $(0, 1)$ onto $(\pi/4, \infty)$;*

(5) *the function $r \mapsto \mathcal{E} + r^2 \mathcal{K}$ is strictly increasing from $(0, 1)$ onto $(\pi/2, \infty)$;*

(6) *the function $r \mapsto [4(2\mathcal{E} - r'^2 \mathcal{K})^2/\pi^2 - (1 - r^2)]/r^2$ is strictly increasing from $(0, 1)$ onto $(3/2, 16/\pi^2)$.*

Proof. Parts (1)-(4) can be found in [13, Theorem 3.21(1), Exercise 3.43(13)], [59, Lemma 2.2(4)] and [11, Theorem 15].

For part (5), let $g(r) = \mathcal{E} + r^2 \mathcal{K}$, then by differentiation we have

$$g'(r) = \frac{\mathcal{E} - \mathcal{K}}{r} + 2r\mathcal{K} + r^2 \frac{\mathcal{E} - r'^2 \mathcal{K}}{rr'^2} = \frac{\mathcal{E} - r'^2 \mathcal{K} + r^2 r'^2 \mathcal{K}}{rr'^2} > 0$$

for all $r \in (0, 1)$. Thus $g(r)$ is strictly increasing on $(0, 1)$, and the limiting values are clear.

For part (6), let $h_1(r) = 4(2\mathcal{E} - r'^2 \mathcal{K})^2/\pi^2 - (1 - r^2)$, $h_2(r) = r^2$ and $h(r) = h_1(r)/h_2(r)$. Then $h(1^-) = 16/\pi^2$, $h_1(0) = h_2(0) = 0$ and

$$\frac{h_1'(r)}{h_2'(r)} = \frac{4}{\pi^2} \left(2\mathcal{E} - r'^2 \mathcal{K} \right) \left(\frac{\mathcal{E} - r'^2 \mathcal{K}}{r^2} \right) + 1. \quad (3.1)$$

It follows from (3.1) and parts (1) and (2) together with Lemma 1.1 that $h(r)$ is strictly increasing on $(0, 1)$ and $h(0^+) = 3/2$.

LEMMA 3.2. *The function*

$$f(r) = \frac{4(1+r^2)(2\mathcal{E} - r'^2 \mathcal{K})^2/\pi^2 - (1-r^2)^2}{r^2}$$

is strictly decreasing from $(0, 1)$ onto $(32/\pi^2, 7/2)$.

Proof. Clearly $f(1^-) = 32/\pi^2$, and by l'Hôpital's rule and Lemma 3.1(1), (2) we have

$$\begin{aligned} \lim_{r \rightarrow 0^+} f(r) &= \lim_{r \rightarrow 0} \left[\frac{4}{\pi^2} (2\mathcal{E} - r'^2 \mathcal{K})^2 + \frac{4}{\pi^2} (1+r^2)(2\mathcal{E} - r'^2 \mathcal{K}) \frac{\mathcal{E} - r'^2 \mathcal{K}}{r^2} + 2(1-r^2) \right] \\ &= \frac{7}{2}. \end{aligned}$$

Differentiating f yields

$$\begin{aligned} f'(r) &= \frac{[8r(2\mathcal{E} - r'^2 \mathcal{K})^2/\pi^2 + 8(1+r^2)(2\mathcal{E} - r'^2 \mathcal{K})(\mathcal{E} - r'^2 \mathcal{K})/r/\pi^2 + 4r(1-r^2)]r^2}{r^4} \\ &\quad - \frac{[4(1+r^2)(\mathcal{E} - r'^2 \mathcal{K})^2/\pi^2 - (1-r^2)^2]2r}{r^4} \\ &= \frac{1}{r^3} \left[-\frac{8}{\pi^2} (2\mathcal{E} - r'^2 \mathcal{K})^2 + \frac{8}{\pi^2} (1+r^2)(2\mathcal{E} - r'^2 \mathcal{K})(\mathcal{E} - r'^2 \mathcal{K}) + 2(1-r^4) \right] \\ &= \frac{2r'^2}{r^3} \left[1+r^2 - \frac{4}{\pi^2} (2\mathcal{E} - r'^2 \mathcal{K})(\mathcal{E} + r^2 \mathcal{K}) \right]. \end{aligned} \quad (3.2)$$

Let

$$f_1(r) = 1 + r^2 - \frac{4}{\pi^2} (2\mathcal{E} - r'^2 \mathcal{K})(\mathcal{E} + r^2 \mathcal{K}), \quad r \in (0, 1). \quad (3.3)$$

Then

$$\lim_{r \rightarrow 0^+} f_1(r) = 0, \quad (3.4)$$

$$\frac{f_1'(r)}{r} = \frac{1}{r} \left[2r - \frac{4}{\pi^2} \frac{\mathcal{E} - r'^2 \mathcal{K}}{r} (\mathcal{E} + r^2 \mathcal{K}) - \frac{4}{\pi^2} (2\mathcal{E} - r'^2 \mathcal{K}) \frac{\mathcal{E} - r'^2 \mathcal{K} + r^2 r'^2 \mathcal{K}}{r r'^2} \right]$$

$$= 2 - \frac{4}{\pi^2} \frac{\mathcal{E} - r'^2 \mathcal{K}}{r^2} (\mathcal{E} + r^2 \mathcal{K}) - \frac{4}{\pi^2} (2\mathcal{E} - r'^2 \mathcal{K}) \left(\frac{\mathcal{E} - r'^2 \mathcal{K}}{r^2 r'^2} + \mathcal{K} \right) = f_2(r). \tag{3.5}$$

The equation (3.5) and Lemma 3.1(1), (2) and (5) imply that $f_2(r)$ is strictly decreasing on $(0, 1)$. Note that $f_2(0) = 0$. Thus $f_2(r) < 0$ for $r \in (0, 1)$. By (3.4) and (3.5) we conclude that $f_1(r)$ is strictly decreasing on $(0, 1)$, and $f_1(r) < 0$ for $r \in (0, 1)$. Therefore, using (3.2) and (3.3), the monotonicity of f follows.

LEMMA 3.3. *Let $c \in \mathbb{R}$. Then the function*

$$f(r) = r'^c \left[\frac{\mathcal{E} - r'^2 \mathcal{K} - r'^2 (\mathcal{K} - \mathcal{E})}{r^4} \right]$$

is strictly decreasing from $(0, 1)$ onto $(0, 3\pi/16)$ if and only if $c \geq 1/2$.

Proof. By power series expansion, one has

$$\begin{aligned} & \frac{2}{\pi} \left[\mathcal{E} - r'^2 \mathcal{K} - r'^2 (\mathcal{K} - \mathcal{E}) \right] \\ &= \frac{2}{\pi} \left[(1 + r'^2) \mathcal{E} - 2r'^2 \mathcal{K} \right] \\ &= (2 - r^2) \sum_{n=0}^{\infty} \frac{(-1/2)_n (1/2)_n}{(n!)^2} r^{2n} - 2(1 - r^2) \sum_{n=0}^{\infty} \frac{(1/2)_n (1/2)_n}{(n!)^2} r^{2n} \\ &= r^2 + (2 - r^2) \sum_{n=1}^{\infty} \frac{(-1/2)_n (1/2)_n}{(n!)^2} r^{2n} - 2(1 - r^2) \sum_{n=1}^{\infty} \frac{(1/2)_n (1/2)_n}{(n!)^2} r^{2n} \\ &= r^2 - \frac{(2 - r^2)}{2} \sum_{n=0}^{\infty} \frac{(1/2)_n (1/2)_{n+1}}{[(n+1)!]^2} r^{2n+2} - 2(1 - r^2) \sum_{n=0}^{\infty} \frac{(1/2)_{n+1} (1/2)_{n+1}}{[(n+1)!]^2} r^{2n+2} \\ &= r^2 - \sum_{n=0}^{\infty} \frac{(1/2)_n (1/2)_{n+1}}{[(n+1)!]^2} r^{2n+2} + \frac{1}{2} \sum_{n=0}^{\infty} \frac{(1/2)_n (1/2)_{n+1}}{[(n+1)!]^2} r^{2n+4} \\ &\quad - 2 \sum_{n=0}^{\infty} \frac{(1/2)_{n+1} (1/2)_{n+1}}{[(n+1)!]^2} r^{2n+2} + 2 \sum_{n=0}^{\infty} \frac{(1/2)_{n+1} (1/2)_{n+1}}{[(n+1)!]^2} r^{2n+4} \\ &= - \sum_{n=0}^{\infty} \frac{(1/2)_{n+1} (1/2)_{n+2}}{[(n+2)!]^2} r^{2n+4} + \frac{1}{2} \sum_{n=0}^{\infty} \frac{(1/2)_n (1/2)_{n+1}}{[(n+1)!]^2} r^{2n+4} \\ &\quad - 2 \sum_{n=0}^{\infty} \frac{(1/2)_{n+2} (1/2)_{n+2}}{[(n+2)!]^2} r^{2n+4} + 2 \sum_{n=0}^{\infty} \frac{(1/2)_{n+1} (1/2)_{n+1}}{[(n+1)!]^2} r^{2n+4} \\ &= \frac{3}{8} \sum_{n=0}^{\infty} \frac{(1/2)_n (3/2)_n}{(3)_n} \frac{r^{2n+4}}{n!} = \frac{3}{8} r^4 F \left(\frac{1}{2}, \frac{3}{2}; 3; r^2 \right). \end{aligned} \tag{3.6}$$

According to (3.6), $f(r)$ can be rewritten as

$$f(r) = \frac{3\pi}{16} r'^c F \left(\frac{1}{2}, \frac{3}{2}; 3; r^2 \right). \tag{3.7}$$

Lemma 2.15(1) in [44] shows that $x \rightarrow (1-x)^d F(a, b; a+b+1, x)$ ($a, b > 0$) is strictly decreasing on $(0, 1)$ if and only if $d \geq ab/(a+b+1)$. Taking $a = 1/2$ and $b = 3/2$, it follows that $f(r)$ is strictly decreasing on $(0, 1)$ if and only if $c \geq 1/2$. Moreover, when $c \geq 1/2$, by (3.7) we get $f(0^+) = 3\pi/16$, and $f(1^+) = 0$.

THEOREM 3.4. *The double inequality*

$$\frac{\alpha}{L(a,b)} + \frac{1-\alpha}{He(a,b)} < \frac{1}{AG(a,b)} < \frac{\beta}{L(a,b)} + \frac{1-\beta}{He(a,b)}$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha \leq 1/2$ and $\beta \geq 2/\pi$, where $He(a, b) = 2A(a, b)/3 + G(a, b)/3$.

Proof. According to the fact that $He(a, b) > L(a, b)$ for all $a, b > 0$ with $a \neq b$, it follows that Theorem 3.4 is equivalent to

$$\alpha < \frac{\frac{1}{AG(a,b)} - \frac{1}{He(a,b)}}{\frac{1}{L(a,b)} - \frac{1}{He(a,b)}} < \beta \tag{3.8}$$

for all $a, b > 0$ with $a \neq b$ if and only if $\alpha \leq 1/2$ and $\beta \geq 2/\pi$.

Since $L(a, b)$, $He(a, b)$ and $AG(a, b)$ are symmetric and homogeneous of degree 1, without loss of generality, we assume that $a = 1 > b$. Let $b = (1-r)/(1+r)$ ($r \in (0, 1)$). Then

$$\frac{A(a,b)}{L(a,b)} = \frac{1/(1+r)}{2r/\log[(1+r)/(1-r)]/(1+r)} = \frac{\text{arth}r}{r}, \tag{3.9}$$

$$\frac{A(a,b)}{AG(a,b)} = \frac{1/(1+r)}{(\pi/2)/\mathcal{K}[2\sqrt{r}/(1+r)]} = \frac{2\mathcal{K}(r)}{\pi}, \tag{3.10}$$

$$\frac{A(a,b)}{He(a,b)} = \frac{1/(1+r)}{2/[3(1+r)] + (1/3)\sqrt{(1-r)/(1+r)}} = \frac{3}{2+r'}, \tag{3.11}$$

$$\frac{\frac{1}{AG(a,b)} - \frac{1}{He(a,b)}}{\frac{1}{L(a,b)} - \frac{1}{He(a,b)}} = \frac{2\mathcal{K}(r)/\pi - 3/(2+r')}{\text{arth}r/r - 3/(2+r')} = \frac{2r\mathcal{K}(r)/\pi - 3r/(2+r')}{\text{arth}r - 3r/(2+r')}. \tag{3.12}$$

Let

$$G(r) = \frac{2r\mathcal{K}(r)/\pi - 3r/(2+r')}{\text{arth}r - 3r/(2+r')}, \quad r \in (0, 1), \tag{3.13}$$

$G_1(r) = 2r\mathcal{K}(r)/\pi - 3r/(2+r')$, $G_2(r) = \text{arth}r - 3r/(2+r')$, $G_3(r) = 2\mathcal{E}(r)/\pi - [3r'(1+2r')]/(2+r')^2$, $G_4(r) = 1 - [3r'(1+2r')]/(2+r')^2$. Then $G(r) = G_1(r)/G_2(r)$, $G_1(0^+) = G_2(0^+) = G_3(0^+) = G_4(0^+) = 0$ and

$$\frac{G'_1(r)}{G'_2(r)} = \frac{2\mathcal{E}(r)/(\pi r'^2) - [3(1+2r')]/[r'(2+r')^2]}{1/r'^2 - [3(1+2r')]/[r'(2+r')^2]} = \frac{G_3(r)}{G_4(r)},$$

$$\begin{aligned} \frac{G'_3(r)}{G'_4(r)} &= \frac{2(\mathcal{E} - \mathcal{K})/(\pi r) + [3r(2 + 7r')]/[r'(2 + r')^3]}{[3r(2 + 7r')]/[r'(2 + r')^3]} \\ &= 1 - \frac{2}{3\pi} \frac{r^{3/4}(\mathcal{K} - \mathcal{E})}{r^2} \cdot \frac{r^{1/4}(2 + r')^3}{7r' + 2}. \end{aligned} \tag{3.14}$$

It is easy to check that $r \rightarrow r(2 + r^4)^3/(7r^4 + 2)$ is strictly increasing from $(0, 1)$ onto $(0, 3)$. Then from (3.14) and Lemma 3.1(4) we know that $G'_3(r)/G'_4(r)$ is strictly increasing on $(0, 1)$, and $\lim_{r \rightarrow 0^+} [G'_3(r)/G'_4(r)] = 1/2$. Applying Lemma 1.1 two times, we obtain that $G(r)$ is strictly increasing on $(0, 1)$, and by l'Hopital's rule, $G(0^+) = \lim_{r \rightarrow 0^+} [G'_3(r)/G'_4(r)] = 1/2$ and $G(1^-) = \lim_{r \rightarrow 1^-} [G_3(r)/G_4(r)] = 2/\pi$. Therefore, Theorem 3.4 directly follows from (3.8)-(3.13).

Combining (1.5) and (3.8), we see that

$$\frac{1}{2L(a,b)} + \frac{1}{2He(a,b)} < \frac{1}{AG(a,b)} < \frac{3}{4L(a,b)} + \frac{1}{4A(a,b)} \tag{3.15}$$

for all $a, b > 0$ with $a \neq b$. The following Theorem 3.5 will be a refinement of (3.15).

THEOREM 3.5. *The double inequality*

$$\begin{aligned} &\alpha \left(\frac{3}{4L(a,b)} + \frac{1}{4A(a,b)} \right) + (1 - \alpha) \left(\frac{1}{2L(a,b)} + \frac{1}{2He(a,b)} \right) \\ &< \frac{1}{AG(a,b)} \beta \left(\frac{3}{4L(a,b)} + \frac{1}{4A(a,b)} \right) + (1 - \beta) \left(\frac{1}{2L(a,b)} + \frac{1}{2He(a,b)} \right) \end{aligned} \tag{3.16}$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha \leq 17/44$ and $\beta \geq 8/\pi - 2$, where $He(a, b) = 2A(a, b)/3 + G(a, b)/3$.

Proof. It suffices to establish the double inequality

$$\alpha < \frac{\frac{1}{AG(a,b)} - \left(\frac{1}{2L(a,b)} + \frac{1}{2He(a,b)} \right)}{\frac{3}{4L(a,b)} + \frac{1}{4A(a,b)} - \left(\frac{1}{2L(a,b)} + \frac{1}{2He(a,b)} \right)} < \beta \tag{3.17}$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha \leq 17/44$ and $\beta \geq 8/\pi - 2$.

Since $L(a, b)$, $A(a, b)$, $He(a, b)$ and $AG(a, b)$ are symmetric and homogeneous of degree 1, without loss of generality, we assume that $a = 1 > b$. Let $b = (1 - r)/(1 + r)$ ($r \in (0, 1)$). Then from (3.9)-(3.12) we have

$$\frac{\frac{1}{AG(a,b)} - \left(\frac{1}{2L(a,b)} + \frac{1}{2He(a,b)} \right)}{\frac{3}{4L(a,b)} + \frac{1}{4A(a,b)} - \left(\frac{1}{2L(a,b)} + \frac{1}{2He(a,b)} \right)}$$

$$\begin{aligned}
 &= \frac{2\mathcal{K}(r)/\pi - \operatorname{ar}th r/(2r) - 3/[2(2+r')]}{\operatorname{ar}th r/(4r) + 1/4 - 3/[2(2+r')]} \\
 &= \frac{4r\mathcal{K}(r)/\pi - \operatorname{ar}th r - 3r/(2+r')}{\operatorname{ar}th r/2 + r/2 - 3r/(2+r')} .
 \end{aligned} \tag{3.18}$$

Let

$$F(r) = \frac{4r\mathcal{K}(r)/\pi - \operatorname{ar}th r - 3r/(2+r')}{\operatorname{ar}th r/2 + r/2 - 3r/(2+r')} , \quad r \in (0, 1), \tag{3.19}$$

$F_1(r) = 4r\mathcal{K}(r)/\pi - \operatorname{ar}th r - 3r/(2+r')$, $F_2(r) = \operatorname{ar}th r/2 + r/2 - 3r/(2+r')$, $F_3(r) = 4\mathcal{E}(r)/\pi - 1 - [3r'(1+2r')]/(2+r')^2$, $F_4(r) = 1/2 + r'^2/2 - [3r'(1+2r')]/(2+r')^2$, $F_5(r) = -4(\mathcal{K} - \mathcal{E})/(\pi r^2) + [3(7r' + 2)]/[r'(2+r')^3]$, $F_6(r) = -1 + [3(7r' + 2)]/[r'(2+r')^3]$. Then $F(r) = F_1(r)/F_2(r)$, $F_1(0^+) = F_2(0^+) = F_3(0^+) = F_4(0^+) = F_5(0^+) = F_6(0^+) = 0$ and

$$\begin{aligned}
 \frac{F'_1(r)}{F'_2(r)} &= \frac{4\mathcal{E}(r)/(\pi r'^2) - 1/r'^2 - [3(1+2r')]/[r'(2+r')^2]}{1/(2r'^2) + 1/2 - [3(1+2r')]/[r'(2+r')^2]} = \frac{F_3(r)}{F_4(r)} \\
 \frac{F'_3(r)}{F'_4(r)} &= \frac{4(\mathcal{E} - \mathcal{K})/(\pi r) + [3r(2+7r')]/[r'(2+r')^3]}{-r + [3r(2+7r')]/[r'(2+r')^3]} = \frac{F_5(r)}{F_6(r)} \\
 \frac{F'_5(r)}{F'_6(r)} &= \frac{-4[\mathcal{E} - r'^2\mathcal{K} - r'^2(\mathcal{K} - \mathcal{E})]/(\pi r^3 r'^2) + 3r(21r'^2 + 8r' + 4)/[r'^3(2+r')^4]}{3r(21r'^2 + 8r' + 4)/[r'^3(2+r')^4]} \\
 &= 1 - \frac{4}{3\pi} \frac{r'^{1/2}[\mathcal{E} - r'^2\mathcal{K} - r'^2(\mathcal{K} - \mathcal{E})]}{r^4} \cdot \frac{r'^{1/2}(2+r')^4}{21r'^2 + 8r' + 4} .
 \end{aligned} \tag{3.20}$$

It is not difficult to verify that $r \rightarrow r(2+r'^2)^4/(21r^4 + 8r^2 + 4)$ is strictly increasing from $(0, 1)$ onto $(0, 27/11)$. Then from (3.20) and Lemma 3.3 we know that $F'_5(r)/F'_6(r)$ is strictly increasing on $(0, 1)$, and $\lim_{r \rightarrow 0^+} [F'_5(r)/F'_6(r)] = 17/44$. Applying Lemma 1.1 three times, we obtain that $F(r)$ is strictly increasing on $(0, 1)$, and by l'Hopital's rule, $F(0^+) = \lim_{r \rightarrow 0^+} [F'_5(r)/F'_6(r)] = 17/44$ and $F(1^-) = \lim_{r \rightarrow 1^-} [F_3(r)/F_4(r)] = 8/\pi - 2$.

Therefore, Theorem 3.5 directly follows from (3.17)-(3.19) together with the monotonicity and range of F on $(0, 1)$.

THEOREM 3.6. *Let $\alpha_1, \alpha_2, \beta_1, \beta_2 \in (0, 1)$. Then:*

(1) *the double inequality*

$$\alpha_1 A(a, b) + (1 - \alpha_1)G(a, b) < E(a, b) < \beta_1 A(a, b) + (1 - \beta_1)G(a, b)$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_1 \leq 3/4$ and $\beta_1 \geq 8/\pi^2$;

(2) *the double inequality*

$$\alpha_2 A(a, b) + (1 - \alpha_2)H(a, b) < E(a, b) < \beta_2 A(a, b) + (1 - \beta_2)H(a, b)$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_2 \leq 8/\pi^2$ and $\beta_2 \geq 7/8$, where $H(a, b) = 2ab/(a+b)$ is classical harmonic mean of a and b .

Proof. It is clear to see that parts (1) and (2) are respectively equivalent to

$$\alpha_1 < \frac{E(a,b) - G(a,b)}{A(a,b) - G(a,b)} < \beta_1,$$

$$\alpha_2 < \frac{E(a,b) - H(a,b)}{A(a,b) - H(a,b)} < \beta_2$$

hold for all $a, b > 0$ with $a \neq b$ if and only if $\alpha_1 \leq 3/4$, $\beta_1 \geq 8/\pi^2$, $\alpha_2 \leq 8/\pi^2$ and $\beta_2 \geq 7/8$.

Without loss of generality, we assume that $a = 1 > b$. Let $b = [(1 - r)/(1 + r)]^2 (r \in (0, 1))$. Then simple computations leads to

$$\frac{E(a,b)}{A(a,b)} = \frac{4 \left[\mathcal{E} \left(\frac{2\sqrt{r}}{1+r} \right) \right]^2 / \pi^2}{(1+r^2)/(1+r)^2} = \frac{4 (2\mathcal{E} - r'^2 \mathcal{K})^2}{\pi^2 (1+r^2)},$$

$$\frac{G(a,b)}{A(a,b)} = \frac{1-r^2}{1+r^2}, \quad \frac{H(a,b)}{A(a,b)} = \left(\frac{1-r^2}{1+r^2} \right)^2.$$

Consequently,

$$\begin{aligned} \frac{E(a,b) - G(a,b)}{A(a,b) - G(a,b)} &= \frac{4(2\mathcal{E} - r'^2 \mathcal{K})^2 / [\pi^2(1+r^2)] - (1-r^2)/(1+r^2)}{1 - (1-r^2)/(1+r^2)} \\ &= \frac{4(2\mathcal{E} - r'^2 \mathcal{K})^2 / \pi^2 - (1-r^2)}{2r^2}, \end{aligned} \tag{3.21}$$

$$\begin{aligned} \frac{E(a,b) - H(a,b)}{A(a,b) - H(a,b)} &= \frac{4(2\mathcal{E} - r'^2 \mathcal{K})^2 / [\pi^2(1+r^2)] - (1-r^2)^2/(1+r^2)^2}{1 - (1-r^2)^2/(1+r^2)^2} \\ &= \frac{4(1+r^2)(2\mathcal{E} - r'^2 \mathcal{K})^2 / \pi^2 - (1-r^2)^2}{4r^2}, \end{aligned} \tag{3.22}$$

Therefore, Theorem 3.6 directly follows from (3.21), (3.22), Lemma 3.1(6) and Lemma 3.2.

THEOREM 3.7. *The double inequality*

$$\alpha \left[\frac{7}{8}A(a,b) + \frac{1}{8}H(a,b) \right] + (1 - \alpha) \left[\frac{3}{4}A(a,b) + \frac{1}{4}G(a,b) \right]$$

$$< E(a,b) < \beta \left[\frac{7}{8}A(a,b) + \frac{1}{8}H(a,b) \right] + (1 - \beta) \left[\frac{3}{4}A(a,b) + \frac{1}{4}G(a,b) \right]$$

holds for all $a, b > 0$ with $a \neq b$ if and only if $\alpha \leq 3/16$ and $\beta \geq 4(16/\pi^2 - 3/2)$.

Proof. With the similar argument in Theorem 3.6, we assume that $a = 1$ and $b = [(1 - r)/(1 + r)]^2 (r \in (0, 1))$. Thus we only need to prove that

$$J(r) = \frac{E(a,b) - [3A(a,b)/4 + G(a,b)/4]}{7A(a,b)/8 + H(a,b)/8 - [3A(a,b)/4 + G(a,b)/4]}$$

$$\begin{aligned}
 &= \frac{E(a,b)/A(a,b) - [3/4 + G(a,b)/(4A(a,b))]}{1/8 + H(a,b)/(8A(a,b)) - [G(a,b)/(4A(a,b))]} \\
 &= \frac{4(2\mathcal{E} - r^2\mathcal{K})^2/\pi^2/(1+r^2) - 3/4 - (1-r^2)/(1+r^2)/4}{1/8 + (1-r^2)^2/(1+r^2)^2/8 - (1-r^2)/(1+r^2)/4} \\
 &= \frac{2(1+r^2) \left[4(2\mathcal{E} - r^2\mathcal{K})^2/\pi^2 - 1 - r^2/2 \right]}{r^4} \tag{3.23}
 \end{aligned}$$

is strictly increasing from $(0, 1)$ onto $(3/16, 4(16/\pi^2 - 3/2))$.

Let $J_1(r) = 2(1+r^2) \left[4(2\mathcal{E} - r^2\mathcal{K})^2/\pi^2 - 1 - r^2/2 \right]$, $J_2(r) = r^4$, $J_3(r) = 4(2\mathcal{E} - r^2\mathcal{K})^2/\pi^2 - 1 - r^2/2 + 1/2(1+r^2)[8(2\mathcal{E} - r^2\mathcal{K})(\mathcal{E} - r^2\mathcal{K})/(\pi^2 r^2) - 1]$ and $J_4(r) = r^2$. Then $J(r) = J_1(r)/J_2(r)$, $J_1(0^+) = J_2(0^+) = J_3(0^+) = J_4(0^+) = 0$ and

$$\begin{aligned}
 \frac{J'_1(r)}{J'_2(r)} &= \frac{4(2\mathcal{E} - r^2\mathcal{K})^2/\pi^2 - 1 - r^2/2}{r^2} + \frac{1+r^2}{2r^2} \left[\frac{8}{\pi^2} (2\mathcal{E} - r^2\mathcal{K}) \frac{\mathcal{E} - r^2\mathcal{K}}{r^2} - 1 \right] \\
 &= \frac{J_3(r)}{J_4(r)}, \tag{3.24}
 \end{aligned}$$

$$\begin{aligned}
 \frac{J'_3(r)}{J'_4(r)} &= \frac{8}{\pi^2} (2\mathcal{E} - r^2\mathcal{K}) \frac{\mathcal{E} - r^2\mathcal{K}}{r^2} - 1 + \frac{2}{\pi^2} (1+r^2) \\
 &\quad \times \left[\left(\frac{\mathcal{E} - r^2\mathcal{K}}{r^2} \right)^2 + (2\mathcal{E} - r^2\mathcal{K}) \frac{\mathcal{K} - \mathcal{E} - (\mathcal{E} - r^2\mathcal{K})}{r^4} \right]. \tag{3.25}
 \end{aligned}$$

The equation (3.25) and Lemma 3.1(1)-(3) show that $J'_3(r)/J'_4(r)$ is strictly increasing from $(0, 1)$ onto $(3/16, +\infty)$. Therefore, the monotonicity of $J(r)$ follows from (3.23) and (3.24) together with Lemma 1.1. Moreover, by l'Hopital's rule, $J(0^+) = \lim_{r \rightarrow 0^+} [J'_3(r)/J'_4(r)] = 3/16$. And $J(1^-) = 4(16/\pi^2 - 3/2)$ is clear.

4. Bounds for $\mathcal{K}(r)$ and $\mathcal{E}(r)$

THEOREM 4.1. *For all $r \in (0, 1)$ we have*

$$\frac{\pi}{2} \left[\alpha \frac{\operatorname{arthr}}{r} + (1-\alpha) \frac{3}{2+r'} \right] < \mathcal{K}(r) < \frac{\pi}{2} \left[\beta \frac{\operatorname{arthr}}{r} + (1-\beta) \frac{3}{2+r'} \right] \tag{4.1}$$

with the best possible constants

$$\alpha = 1/2, \quad \beta = 2/\pi.$$

Proof. Taking $a = 1 + r$ and $b = 1 - r$ in Theorem 3.4, we obtain the result.

THEOREM 4.2. *For all $r \in (0, 1)$ we have*

$$\frac{\pi}{2} \left[\alpha \left(\frac{3}{4L(1, r')} + \frac{1}{4A(1, r')} \right) + (1-\alpha) \left(\frac{1}{2L(1, r')} + \frac{1}{2He(1, r')} \right) \right]$$

$$\langle \mathcal{K}(r) \rangle < \frac{\pi}{2} \left[\beta \left(\frac{3}{4L(1,r')} + \frac{1}{4A(1,r')} \right) + (1-\beta) \left(\frac{1}{2L(1,r')} + \frac{1}{2He(1,r')} \right) \right], \tag{4.2}$$

$$\begin{aligned} & \frac{\pi}{2} \left[\alpha \left(\frac{3}{4} \frac{\operatorname{arth} r}{r} + \frac{1}{4} \right) + (1-\alpha) \left(\frac{\operatorname{arth} r}{2r} + \frac{3}{2(2+r')} \right) \right] \\ \langle \mathcal{K}(r) \rangle & < \frac{\pi}{2} \left[\beta \left(\frac{3}{4} \frac{\operatorname{arth} r}{r} + \frac{1}{4} \right) + (1-\beta) \left(\frac{\operatorname{arth} r}{2r} + \frac{3}{2(2+r')} \right) \right] \end{aligned} \tag{4.3}$$

with the best possible constants

$$\alpha = 17/44, \quad \beta = 8/\pi - 2.$$

Proof. Two inequalities in Theorem 4.2 follow easily from Theorem 3.5 with $a = 1$, $b = r'$ and $a = 1 + r$, $b = 1 - r$, respectively.

THEOREM 4.3. For all $r \in (0, 1)$ we have

$$\frac{\pi}{2} \sqrt{\alpha_1 \frac{1+r'^2}{2} + (1-\alpha_1)r'} < \mathcal{E}(r) < \frac{\pi}{2} \sqrt{\beta_1 \frac{1+r'^2}{2} + (1-\beta_1)r'}, \tag{4.4}$$

$$\frac{\pi}{2} \sqrt{\alpha_2 \frac{1+r'^2}{2} + (1-\alpha_2) \frac{2r'^2}{1+r'^2}} < \mathcal{E}(r) < \frac{\pi}{2} \sqrt{\beta_2 \frac{1+r'^2}{2} + (1-\beta_2) \frac{2r'^2}{1+r'^2}} \tag{4.5}$$

with the best possible constants

$$\alpha_1 = 3/4, \quad \beta_1 = 8/\pi^2, \quad \alpha_2 = 8/\pi^2, \quad \beta_2 = 7/8.$$

Proof. Theorem 4.3 follows from Theorem 3.6 with $a = 1$ and $b = r'^2$. Similarly, letting $a = 1$ and $b = r'^2$ in Theorem 3.7, one has

THEOREM 4.4. For all $r \in (0, 1)$ we have

$$\begin{aligned} & \frac{\pi}{2} \sqrt{\alpha \left[\frac{7+18r'^2+7r'^4}{16(1+r'^2)} \right] + (1-\alpha) \left(\frac{3r'^2+2r'+3}{8} \right)} \\ \langle \mathcal{E}(r) \rangle & < \frac{\pi}{2} \sqrt{\beta \left[\frac{7+18r'^2+7r'^4}{16(1+r'^2)} \right] + (1-\beta) \left(\frac{3r'^2+2r'+3}{8} \right)} \end{aligned} \tag{4.6}$$

with the best possible constants

$$\alpha = 3/16, \quad \beta = 4(16/\pi^2 - 3/2).$$

5. Comparisons

REMARK 5.1. Since $1/AG(1+r, 1-r) = 2\mathcal{K}(r)/\pi$, $1/L(1+r, 1-r) = \text{arth}r/r$ and $1/A(1+r, 1-r) = 1$, the right-hand side of inequality (1.4) can be expressed as

$$\frac{1}{AG(a,b)} < \frac{1}{L(a,b)}$$

with $a = 1+r$ and $b = 1-r$. Thus Theorems 3.4 and 3.5 imply that our upper bounds in (4.1) and (4.3) are better than that in (1.4). On the other hand, it follows from (3.15) and (3.16) that the upper bound in (4.2) is better than that in (1.5).

Next, we shall compare the lower bound in (4.3) with the lower bound in (1.4). For this purpose we need the following two propositions.

PROPOSITION 5.2. For all $r \in (0, 1)$, we have

$$\frac{r^2 \text{arth}r}{r - r'^2 \text{arth}r} > \frac{3}{2} + \frac{1}{5}r^2 + \frac{23}{175}r^4, \tag{5.1}$$

$$35\text{arth}r + \frac{216r'}{(2+r')^2} \left(\frac{3}{2}r + \frac{1}{5}r^3 + \frac{23}{175}r^5 \right) - \frac{162r}{2+r'} - 17r > 0. \tag{5.2}$$

Proof. Clearly, inequality (5.1) can be rewritten as

$$\frac{\text{arth}r}{r} > \frac{r - r'^2 \text{arth}r}{r^3} \left(\frac{3}{2} + \frac{1}{5}r^2 + \frac{23}{175}r^4 \right).$$

Since $\text{arth}r/r = \sum_{n=0}^{\infty} [r^{2n}/(2n+1)]$ and $(r - r'^2 \text{arth}r)/r^3 = [1 - (1-r^2)\text{arth}r/r]/r^2 = \sum_{n=0}^{\infty} (2r^{2n})/[(2n+1)(2n+3)]$, one has

$$\begin{aligned} & \frac{\text{arth}r}{r} - \frac{r - r'^2 \text{arth}r}{r^3} \left(\frac{3}{2} + \frac{1}{5}r^2 + \frac{23}{175}r^4 \right) \\ &= \sum_{n=0}^{\infty} \frac{r^{2n}}{2n+1} - \left(\frac{3}{2} + \frac{1}{5}r^2 + \frac{23}{175}r^4 \right) \sum_{n=0}^{\infty} \frac{2r^{2n}}{(2n+1)(2n+3)} \\ &= \sum_{n=2}^{\infty} \frac{r^{2n}}{2n+1} - \sum_{n=2}^{\infty} \frac{3r^{2n}}{(2n+1)(2n+3)} - \frac{2}{5} \sum_{n=1}^{\infty} \frac{r^{2n+2}}{(2n+1)(2n+3)} - \frac{46}{175} \sum_{n=0}^{\infty} \frac{r^{2n+4}}{(2n+1)(2n+3)} \\ &= \sum_{n=0}^{\infty} \frac{r^{2n+4}}{2n+5} - \sum_{n=0}^{\infty} \frac{3r^{2n+4}}{(2n+5)(2n+7)} - \frac{2}{5} \sum_{n=0}^{\infty} \frac{r^{2n+4}}{(2n+3)(2n+5)} - \frac{46}{175} \sum_{n=0}^{\infty} \frac{r^{2n+4}}{(2n+1)(2n+3)} \\ &= \sum_{n=0}^{\infty} \frac{2n(700n^2 + 2568n + 2213)}{175(2n+1)(2n+3)(2n+5)(2n+7)} r^{2n+4} > 0 \end{aligned}$$

for all $r \in (0, 1)$.

For the inequality (5.2), set

$$f(r) = 35\text{arth}r + \frac{216r'}{(2+r')^2} \left(\frac{3}{2}r + \frac{1}{5}r^3 + \frac{23}{175}r^5 \right) - \frac{162r}{2+r'} - 17r, \quad r \in (0, 1).$$

Then $f(0^+) = 0$, and by tedious computations we have

$$f'(r) = \frac{19872r^8 - 29592r^6 - 30810r^4 + 130480r^2 - 40950 - \sqrt{1-r^2}g(r)}{175(1-r^2)(2+r')^3}, \tag{5.3}$$

where $g(r) = 59616r^6 + 13775r^4 + 89215r^2 - 40950$ is strictly increasing on $(0, 1)$, and there exists a unique zero point $r_0 = 0.62778 \dots$ such that $g(r) < 0$ for $r \in (0, r_0)$, and $g(r) > 0$ for $r \in (r_0, 1)$.

Finally, it is sufficient to prove that $f'(r) > 0$ for $r \in (0, r_0]$ and $r \in (r_0, 1)$.

When $r \in (0, r_0]$, noting that the fact $1 - r^2 < \sqrt{1 - r^2}$ for $r \in (0, 1)$, we have

$$\begin{aligned} & 19872r^8 - 29592r^6 - 30810r^4 + 130480r^2 - 40950 - \sqrt{1-r^2}g(r) \\ & > 19872r^8 - 29592r^6 - 30810r^4 + 130480r^2 - 40950 - (1-r^2)g(r) \\ & = r^2[79488r^6 + r^2(44630 - 75433r^2) + 315] > 0 \end{aligned}$$

for all $r \in (0, r_0]$. Thus it follows from (5.3) that $f'(r) > 0$ for all $r \in (0, r_0]$.

When $r \in (r_0, 1)$, by the inequality $\sqrt{1-r^2} < 1 - r^2/2 - r^4/8$, we get

$$\begin{aligned} & 19872r^8 - 29592r^6 - 30810r^4 + 130480r^2 - 40950 - \sqrt{1-r^2}g(r) \\ & > 19872r^8 - 29592r^6 - 30810r^4 + 130480r^2 - 40950 - (1 - r^2/2 - r^4/8)g(r) \\ & = \frac{(59616r^8 + 411215r^6 - 569349r^4 - 40770r^2 + 166320)r^2}{8}. \end{aligned}$$

It is not difficult to verify that inequality $59616r^8 + 411215r^6 - 569349r^4 - 40770r^2 + 166320 > 0$ takes place for all $r \in (0, 1)$. Thus (5.3) leads to that $f'(r) > 0$ for all $r \in (r_0, 1)$.

PROPOSITION 5.3. *Let*

$$F(r) = \log \left(\frac{105 \operatorname{arth} r}{176} \frac{1}{r} + \frac{81}{88(2+r')} + \frac{17}{176} \right) - \frac{3}{4} \log \left(\frac{\operatorname{arth} r}{r} \right).$$

Then $F(r)$ is strictly increasing from $(0, 1)$ onto $(0, \infty)$.

Proof. Differentiating F yields

$$\begin{aligned} F'(r) &= \frac{105(r - r'^2 \operatorname{arth} r)/(r^2 r'^2) + 162r/[r'(2+r')^2]}{105(\operatorname{arth} r)/r + 162/(2+r') + 17} - \frac{3}{4} \frac{r - r'^2 \operatorname{arth} r}{r r'^2 \operatorname{arth} r} \\ &= \frac{3(r - r'^2 \operatorname{arth} r)F_1(r)}{4r r'^2 \operatorname{arth} r [105(\operatorname{arth} r)/r + 162/(2+r') + 17]} \end{aligned} \tag{5.4}$$

where

$$F_1(r) = 35 \frac{\operatorname{arth} r}{r} + \frac{216r'}{(2+r')^2} \frac{r^2 \operatorname{arth} r}{r - r'^2 \operatorname{arth} r} - \frac{162}{2+r'} - 17. \tag{5.5}$$

It follows from (5.1) and (5.2) that

$$F_1(r) > 35 \frac{\operatorname{arth} r}{r} + \frac{216r'}{(2+r')^2} \left(\frac{3}{2} + \frac{1}{5}r'^2 + \frac{23}{175}r'^4 \right) - \frac{162}{2+r'} - 17 > 0 \quad (5.6)$$

for all $r \in (0, 1)$.

Therefore, the monotonicity of $F(r)$ follows from (5.4)-(5.6). The limiting values are clear.

REMARK 5.4. From Proposition 5.3 we conclude that the lower bound in (4.3) is better than the lower bound in (1.4).

REMARK 5.5. The lower bound in (4.5) is better than the lower bound in (1.6) when $r \rightarrow 1$. Computational and numerical experiments show that the lower bound in (4.6) is tighter than that in (1.6) for $0 \leq r \leq 0.972$. Also, computational and numerical experiments show that the upper bound in (4.6) is tighter than that in (1.7) for $0 \leq r \leq 0.953$.

Acknowledgement. This research was supported by the Natural Science Foundation of China (Grant Nos. 11701176, 11971142, 61673169).

REFERENCES

- [1] I. ABBAS BALOCH, Y.-M. CHU, *Petrović-type inequalities for harmonic h -convex functions*, J. Funct. Spaces **2020** (2020), Article ID 3075390, 7 pages.
- [2] M. ADIL KHAN, S. BEGUM, Y. KHURSHID, Y.-M. CHU, *Ostrowski type inequalities involving conformable fractional integrals*, J. Inequal. Appl. **2018** (2018), Article 70, 14 pages.
- [3] M. ADIL KHAN, Y.-M. CHU, A. KASHURI, R. LIKO, G. ALI, *Conformable fractional integrals versions of Hermite-Hadamard inequalities and their generalizations*, J. Funct. Spaces **2018** (2018), Article ID 6928130, 9 pages.
- [4] M. ADIL KHAN, Y.-M. CHU, T. U. KHAN, J. KHAN, *Some new inequalities of Hermite-Hadamard type for s -convex functions with applications*, Open Math. **15**, 1 (2017), 1414–1430.
- [5] M. ADIL KHAN, M. HANIF, Z. A. KHAN, K. AHMAD, Y.-M. CHU, *Association of Jensen's inequality for s -convex function with Csiszár divergence*, J. Inequal. Appl. **2019** (2019), Article 162, 14 pages.
- [6] M. ADIL KHAN, A. IQBAL, M. SULEMAN, Y.-M. CHU, *Hermite-Hadamard type inequalities for fractional integrals via Green's function*, J. Inequal. Appl. **2018** (2018), Article 161, 15 pages.
- [7] M. ADIL KHAN, Y. KHURSHID, T.-S. DU, Y.-M. CHU, *Generalization of Hermite-Hadamard type inequalities via conformable fractional integrals*, J. Funct. Spaces **2018** (2018), Article ID 5357463, 12 pages.
- [8] M. ADIL KHAN, N. MOHAMMAD, E. R. NWAEZE, Y.-M. CHU, *Quantum Hermite-Hadamard inequality by means of a Green function*, Adv. Difference Equ. **2020** (2020), Article 99, 20 pages.
- [9] M. ADIL KHAN, S.-H. WU, H. ULLAH, Y.-M. CHU, *Discrete majorization type inequalities for convex functions on rectangles*, J. Inequal. Appl. **2019** (2019), Article 16, 18 pages.
- [10] M. ADIL KHAN, S. ZAHEER ULLAH, Y.-M. CHU, *The concept of coordinate strongly convex functions and related inequalities*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **113**, 3 (2019), 2235–2251.
- [11] H. ALZER, S.-L. QIU, *Monotonicity theorems and inequalities for the complete elliptic integrals*, J. Comput. Appl. Math. **172**, 2 (2004), 289–312.

- [12] G. D. ANDERSON, M. K. VAMANAMURTHY, M. VUORINEN, *Functional inequalities for hypergeometric functions and complete elliptic integrals*, SIAM J. Math. Anal. **23**, 2 (1992), 512–524.
- [13] G. D. ANDERSON, M. K. VAMANAMURTHY, M. VUORINEN, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.
- [14] R. W. BARNARD, K. PEARCE, K. C. RICHARDS, *An inequality involving the generalized hypergeometric function and the arc length of an ellipse*, SIAM J. Math. Anal. **31**, 3 (2000), 693–699.
- [15] R. W. BARNARD, K. PEARCE, K. C. RICHARDS, *A monotonicity property involving ${}_3F_2$ and comparisons of the classical approximations of elliptical arc length*, SIAM J. Math. Anal. **32**, 2 (2000), 403–419.
- [16] Y.-M. CHU, M. ADIL KHAN, T. ALI, S. S. DRAGOMIR, *Inequalities for α -fractional differentiable functions*, J. Inequal. Appl. **2017** (2017), Article 93, 12 pages.
- [17] Y.-M. CHU, B.-Y. LONG, *Sharp inequalities between means*, Math. Inequal. Appl. **14**, 3 (2011), 647–655.
- [18] Y.-M. CHU, Y.-F. QIU, M.-K. WANG, *Hölder mean inequalities for the complete elliptic integrals*, Integral Transforms Spec. Funct. **23**, 7 (2012), 521–527.
- [19] Y.-M. CHU, M.-K. WANG, *Inequalities between arithmetic-geometric, Gini, and Toader means*, Abstr. Appl. Anal. **2012** (2012), Article ID 830585, 11 pages.
- [20] Y.-M. CHU, M.-K. WANG, *Optimal Lehmer mean bounds for the Toader mean*, Results Math. **61**, 3–4 (2012), 223–229.
- [21] Y.-M. CHU, M.-K. WANG, Y.-P. JIANG, S.-L. QIU, *Concavity of the complete elliptic integrals of the second kind with respect to Hölder means*, J. Math. Anal. Appl. **395**, 2 (2012), 637–642.
- [22] Y.-M. CHU, M.-K. WANG, S.-L. QIU, *Optimal combinations bounds of root-square and arithmetic means for Toader mean*, Proc. Indian Acad. Sci. Math. Sci. **122**, 1 (2012), 41–51.
- [23] Y.-M. CHU, M.-K. WANG, S.-L. QIU, Y.-P. JIANG, *Bounds for complete elliptic integrals of the second kind with applications*, Comput. Math. Appl. **63**, 7 (2012), 1177–1184.
- [24] Y.-M. CHU, G.-D. WANG, X.-H. ZHANG, *The Schur multiplicative and harmonic convexities of the complete symmetric function*, Math. Nachr. **284**, 5–6 (2011), 653–663.
- [25] Y.-M. CHU, W.-F. XIA, X.-H. ZHANG, *The Schur concavity, Schur multiplicative and harmonic convexities of the second dual form of the Hamy symmetric function with applications*, J. Multivariate Anal. **105** (2012), 412–421.
- [26] G.-J. HAI, T.-H. ZHAO, *Monotonicity properties and bounds involving the two-parameter generalized Grötzsch ring function*, J. Inequal. Appl. **2020** (2020), Article 66, 17 pages.
- [27] X.-H. HE, W.-M. QIAN, H.-Z. XU, Y.-M. CHU, *Sharp power mean bounds for two Sándor-Yang means*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM **113**, 3 (2019), 2627–2638.
- [28] X.-M. HU, J.-F. TIAN, Y.-M. CHU, Y.-X. LU, *On Cauchy-Schwarz inequality for N -tuple diamond-alpha integral*, J. Inequal. Appl. **2020** (2020), Article 8, 15 pages.
- [29] T.-R. HUANG, B.-W. HAN, X.-Y. MA, Y.-M. CHU, *Optimal bounds for the generalized Euler-Mascheroni constant*, J. Inequal. Appl. **2018** (2018), Article 118, 9 pages.
- [30] T.-R. HUANG, S.-Y. TAN, X.-Y. MA, Y.-M. CHU, *Monotonicity properties and bounds for the complete p -elliptic integrals*, J. Inequal. Appl. **2018** (2018), Article 239, 11 pages.
- [31] A. IQBAL, M. ADIL KHAN, S. ULLAH, Y.-M. CHU, *Some new Hermite-Hadamard-type inequalities associated with conformable fractional integrals and their applications*, J. Funct. Spaces **2020** (2020), Article ID 9845407, 18 pages.
- [32] S. KHAN, M. ADIL KHAN, Y.-M. CHU, *Converses of the Jensen inequality derived from the Green functions with applications in information theory*, Math. Methods Appl. Sci. **43**, 5 (2020), 2577–2587.
- [33] Y. KHURSHID, M. ADIL KHAN, Y.-M. CHU, *Conformable integral inequalities of the Hermite-Hadamard type in terms of GG- and GA-convexities*, J. Funct. Spaces **2019** (2019), Article ID 6926107, 8 pages.
- [34] Y. KHURSHID, M. ADIL KHAN, Y.-M. CHU, Z. A. KHAN, *Hermite-Hadamard-Fejér inequalities for conformable fractional integrals via preinvex functions*, J. Funct. Spaces **2019** (2019), Article ID 3146210, 9 pages.
- [35] M. A. LATIF, S. RASHID, S. S. DRAGOMIR, Y.-M. CHU, *Hermite-Hadamard type inequalities for co-ordinated convex and quasi-convex functions and their applications*, J. Inequal. Appl. **2019** (2019), Article 317, 33 pages.
- [36] W.-M. QIAN AND Y.-M. CHU, *Sharp bounds for a special quasi-arithmetic mean in terms of arithmetic and geometric means with two parameters*, J. Inequal. Appl. **2017** (2017), Article 274, 10 pages.

- [37] W.-M. QIAN, Z.-Y. HE, Y.-M. CHU, *Approximation for the complete elliptic integral of the first kind*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM **114**, 2 (2020), Article 57, 12 pages. Available online at <https://doi.org/10.1007/s13398-020-00784-9>.
- [38] W.-M. QIAN, Z.-Y. HE, H.-W. ZHANG, Y.-M. CHU, *Sharp bounds for Neuman means in terms of two-parameter contraharmonic and arithmetic mean*, J. Inequal. Appl. **2019** (2019), Article 168, 13 pages.
- [39] W.-M. QIAN, H.-Z. XU, Y.-M. CHU, *Improvements of bounds for the Sándor-Yang means*, J. Inequal. Appl. **2019** (2019), Article 73, 8 pages.
- [40] W.-M. QIAN, Y.-Y. YANG, H.-W. ZHANG, Y.-M. CHU, *Optimal two-parameter geometric and arithmetic mean bounds for the Sándor-Yang mean*, J. Inequal. Appl. **2019** (2019), Article 287, 12 pages.
- [41] W.-M. QIAN, X.-H. ZHANG, Y.-M. CHU, *Sharp bounds for the Toader-Qi mean in terms of harmonic and geometric means*, J. Math. Inequal. **11**, 1 (2017), 121–127.
- [42] W.-M. QIAN, W. ZHANG, Y.-M. CHU, *Bounding the convex combination of arithmetic and integral means in terms of one-parameter harmonic and geometric means*, Miskolc Math. Notes **20**, 2 (2019), 1157–1166.
- [43] S.-L. QIU, X.-Y. MA, Y.-M. CHU, *Sharp Landen transformation inequalities for hypergeometric functions, with applications*, J. Math. Anal. Appl. **474**, 2 (2019), 1306–1337.
- [44] S.-L. QIU, M. VUORINEN, *Duplication inequalities for the ratios of hypergeometric functions*, Forum Math. **12**, 1 (2000), 109–133.
- [45] Y.-F. QIU, M.-K. WANG, Y.-M. CHU, G.-D. WANG, *Two sharp inequalities for Lehmer mean, identric mean and logarithmic mean*, J. Math. Inequal. **5**, 3 (2011), 301–306.
- [46] S. RAFEEQ, H. KALSOOM, S. HUSSAIN, S. RASHID, Y.-M. CHU, *Delay dynamic double integral inequalities on time scales with applications*, Adv. Difference Equ. **2020** (2020), Article 40, 32 pages.
- [47] S. RASHID, R. ASHRAF, M. A. NOOR, K. I. NOOR, Y.-M. CHU, *New weighted generalizations for differentiable exponentially convex mapping with application*, AIMS Math. **5**, 4 (2020), 3525–3546.
- [48] S. RASHID, F. JARAD, Y.-M. CHU, *A note on reverse Minkowski inequality via generalized proportional fractional integral operator with respect to another function*, Math. Probl. Eng. **2020** (2020), Article ID 7630260, 12 pages.
- [49] S. RASHID, F. JARAD, H. KALSOOM AND Y.-M. CHU, *On Pólya-Szegő and Čebyšev type inequalities via generalized k -fractional integrals*, Adv. Difference Equ. **2020** (2020), Article 125, 18 pages.
- [50] S. RASHID, M. A. NOOR, K. I. NOOR, Y.-M. CHU, *Ostrowski type inequalities in the sense of generalized \mathcal{H} -fractional integral operator for exponentially convex functions*, AIMS Math. **5**, 3 (2020), 2629–2645.
- [51] J. SÁNDOR, *On certain inequalities for means*, J. Math. Anal. Appl. **189**, 2 (1995), 602–606.
- [52] Y.-Q. SONG, M. ADIL KHAN, S. ZAHEER ULLAH, Y.-M. CHU, *Integral inequalities involving strongly convex functions*, J. Funct. Spaces **2018** (2018), Article ID 6595921, 8 pages.
- [53] GH. TOADER, *Some mean values related to the arithmetic-geometric mean*, J. Math. Anal. Appl. **218**, 2 (1998), 358–368.
- [54] M. VUORINEN, *Hypergeometric functions in geometric function theory*, in: *Special Functions and Differential Equations* (Madras, 1997), 119–126, Allied Publ., New Delhi, 1998.
- [55] M.-K. WANG, Y.-M. CHU, *Refinements of transformation inequalities for zero-balanced hypergeometric functions*, Acta Math. Sci. **37B**, 3 (2017), 607–622.
- [56] M.-K. WANG, Y.-M. CHU, *Landen inequalities for a class of hypergeometric functions with applications*, Math. Inequal. Appl. **21**, 2 (2018), 521–537.
- [57] M.-K. WANG, H.-H. CHU, Y.-M. CHU, *Precise bounds for the weighted Hölder mean of the complete p -elliptic integrals*, J. Math. Anal. Appl. **480**, 2 (2019), Article ID 123388, 9 pages, Available online at <https://doi.org/10.1016/j.jmaa.2019.123388>.
- [58] M.-K. WANG, Y.-M. CHU, Y.-P. JIANG, *Ramanujan's cubic transformation inequalities for zero-balanced hypergeometric functions*, Rocky Mountain J. Math. **46**, 2 (2016), 679–691.
- [59] M.-K. WANG, Y.-M. CHU, S.-L. QIU, Y.-P. JIANG, *Convexity of the complete elliptic integrals of the first kind with respect to Hölder means*, J. Math. Anal. Appl. **388**, 2 (2012), 1141–1146.
- [60] M.-K. WANG, Y.-M. CHU, S.-L. QIU, Y.-P. JIANG, *Bounds for the perimeter of an ellipse*, J. Approx. Theory **164**, 7 (2012), 928–937.
- [61] M.-K. WANG, Y.-M. CHU, Y.-F. QIU, S.-L. QIU, *An optimal power mean inequality for the complete elliptic integrals*, Appl. Math. Lett. **24**, 6 (2011), 887–890.

- [62] M.-K. WANG, Y.-M. CHU, W. ZHANG, *Monotonicity and inequalities involving zero-balanced hypergeometric function*, *Math. Inequal. Appl.* **22**, 2 (2019), 601–617.
- [63] M.-K. WANG, Y.-M. CHU, W. ZHANG, *Precise estimates for the solution of Ramanujan's generalized modular equation*, *Ramanujan J.* **49**, 3 (2019), 653–668.
- [64] M.-K. WANG, Z.-Y. HE, Y.-M. CHU, *Sharp power mean inequalities for the generalized elliptic integral of the first kind*, *Comput. Methods Funct. Theory* **20**, 1 (2020), 111–124.
- [65] M.-K. WANG, M.-Y. HONG, Y.-F. XU, Z.-H. SHEN, Y.-M. CHU, *Inequalities for generalized trigonometric and hyperbolic functions with one parameter*, *J. Math. Inequal.* **14**, 1 (2020), 1–21.
- [66] M.-K. WANG, Y.-M. LI, Y.-M. CHU, *Inequalities and infinite product formula for Ramanujan generalized modular equation function*, *Ramanujan J.* **46**, 1 (2018), 189–200.
- [67] B. WANG, C.-L. LUO, S.-H. LI, Y.-M. CHU, *Sharp one-parameter geometric and quadratic means bounds for the Sándor-Yang means*, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM* **114**, 1 (2020), Article 7, 10 pages, Available online at <https://doi.org/10.1007/s13398-019-00734-0>.
- [68] J.-L. WANG, W.-M. QIAN, Z.-Y. HE, Y.-M. CHU, *On approximating the Toader mean by other bivariate means*, *J. Funct. Spaces* **2019** (2019), Article ID 6082413, 7 pages.
- [69] M.-K. WANG, S.-L. QIU, Y.-M. CHU, *Infinite series formula for Hübner upper bound function with applications to Hersch-Pfluger distortion function*, *Math. Inequal. Appl.* **21**, 3 (2018), 629–648.
- [70] M.-K. WANG, S.-L. QIU, Y.-M. CHU, Y.-P. JIANG, *Generalized Hersch-Pfluger distortion function and complete elliptic integrals*, *J. Math. Anal. Appl.* **385**, 1 (2012), 221–229.
- [71] M.-K. WANG, Z.-K. WANG, Y.-M. CHU, *An optimal double inequality between geometric and identric means*, *Appl. Math. Lett.* **25**, 3 (2012), 471–475.
- [72] G.-D. WANG, X.-H. ZHANG, Y.-M. CHU, *A power mean inequality for the Grötzsch ring function*, *Math. Inequal. Appl.* **14**, 4 (2011), 833–837.
- [73] G.-D. WANG, X.-H. ZHANG, Y.-M. CHU, *A power mean inequality involving the complete elliptic integrals*, *Rocky Mountain J. Math.* **44**, 5 (2014), 1661–1667.
- [74] M.-K. WANG, W. ZHANG, Y.-M. CHU, *Monotonicity, convexity and inequalities involving the generalized elliptic integrals*, *Acta Math. Sci.* **39B**, 5 (2019), 1440–1450.
- [75] S.-H. WU, Y.-M. CHU, *Schur m -power convexity of generalized geometric Bonferroni mean involving three parameters*, *J. Inequal. Appl.* **2019** (2019), Article 57, 11 pages.
- [76] W.-F. XIA, Y.-M. CHU, *The Schur convexity of Gini mean values in the sense of harmonic mean*, *Acta Math. Sci.* **31B**, 3 (2011), 1103–1112.
- [77] W.-F. XIA, W. JANOUS, Y.-M. CHU, *The optimal convex combination bounds of arithmetic and harmonic means in terms of power mean*, *J. Math. Inequal.* **6**, 2 (2012), 241–248.
- [78] W.-F. XIA, Y.-M. LI, Y.-M. CHU, S.-Y. XU, W.-M. CHEN, Z.-Q. ZHANG, *Observer-based mixed passive and H_∞ control for uncertain Markovian jump systems with time delays using quantized measurements*, *Nonlinear Anal. Hybrid Syst.* **31** (2019), 233–246.
- [79] W.-F. XIA, S.-Y. XU, J.-W. LUO, Z.-Q. ZHANG, Y.-M. CHU, *Reliable filter design for discrete-time neural networks with Markovian jumping parameters and time-varying delay*, *J. Franklin Inst.* **357**, 5 (2020), 2892–2915.
- [80] H.-Z. XU, Y.-M. CHU, W.-M. QIAN, *Sharp bounds for the Sándor-Yang means in terms of arithmetic and contra-harmonic means*, *J. Inequal. Appl.* **2018** (2018), Article 127, 13 pages.
- [81] Z.-H. YANG, Y.-M. CHU, *A monotonicity property involving the generalized elliptic integral of the first kind*, *Math. Inequal. Appl.* **20**, 3 (2017), 729–735.
- [82] Z.-H. YANG, Y.-M. CHU, W. ZHANG, *High accuracy asymptotic bounds for the complete elliptic integral of the second kind*, *Appl. Math. Comput.* **348** (2019), 552–564.
- [83] Z.-H. YANG, W.-M. QIAN, Y.-M. CHU, *Monotonicity properties and bounds involving the complete elliptic integrals of the first kind*, *Math. Inequal. Appl.* **21**, 4 (2018), 1185–1199.
- [84] Z.-H. YANG, W.-M. QIAN, Y.-M. CHU, W. ZHANG, *Monotonicity rule for the quotient of two functions and its application*, *J. Inequal. Appl.* **2017** (2017), Article 106, 13 pages.
- [85] Z.-H. YANG, W.-M. QIAN, Y.-M. CHU, W. ZHANG, *On rational bounds for the gamma function*, *J. Inequal. Appl.* **2017** (2017), Article 210, 17 pages.
- [86] Z.-H. YANG, W.-M. QIAN, Y.-M. CHU, W. ZHANG, *On approximating the error function*, *Math. Inequal. Appl.* **21**, 2 (2018), 469–479.
- [87] Z.-H. YANG, W.-M. QIAN, Y.-M. CHU, W. ZHANG, *On approximating the arithmetic-geometric mean and complete elliptic integral of the first kind*, *J. Math. Anal. Appl.* **462**, 2 (2018), 1714–1726.

- [88] Z.-H. YANG, W.-M. QIAN, W. ZHANG, Y.-M. CHU, *Notes on the complete elliptic integral of the first kind*, *Math. Inequal. Appl.* **23**, 1 (2020), 77–93.
- [89] Z.-H. YANG, W. ZHANG, Y.-M. CHU, *Sharp Gautschi inequality for parameter $0 < p < 1$ with applications*, *Math. Inequal. Appl.* **20**, 4 (2017), 1107–1120.
- [90] S. ZAHEER ULLAH, M. ADIL KHAN, Y.-M. CHU, *Majorization theorems for strongly convex functions*, *J. Inequal. Appl.* **2019** (2019), Article 58, 13 pages.
- [91] S. ZAHEER ULLAH, M. ADIL KHAN, Y.-M. CHU, *A note on generalized convex functions*, *J. Inequal. Appl.* **2019** (2019), Article 291, 10 pages.
- [92] S. ZAHEER ULLAH, M. ADIL KHAN, Z. A. KHAN, Y.-M. CHU, *Integral majorization type inequalities for the functions in the sense of strong convexity*, *J. Funct. Spaces* **2019** (2019), Article ID 9487823, 11 pages.
- [93] T.-H. ZHAO, Y.-M. CHU, H. WANG, *Logarithmically complete monotonicity properties relating to the gamma function*, *Abstr. Appl. Anal.* **2011** (2011), Article ID 896483, 13 pages.
- [94] T.-H. ZHAO, L. SHI, Y.-M. CHU, *Convexity and concavity of the modified Bessel functions of the first kind with respect to Hölder means*, *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM* **114**, 2 (2020), Article 96, 14 pages, Available online at <https://doi.org/10.1007/s13398-020-00825-3>.
- [95] T.-H. ZHAO, M.-K. WANG, W. ZHANG, Y.-M. CHU, *Quadratic transformation inequalities for Gaussian hypergeometric function*, *J. Inequal. Appl.* **2018** (2018), Article 251, 15 pages.
- [96] T.-H. ZHAO, B.-C. ZHOU, M.-K. WANG, Y.-M. CHU, *On approximating the quasi-arithmetic mean*, *J. Inequal. Appl.* **2019** (2019), Article 42, 12 pages.

(Received February 22, 2017)

Miao-Kun Wang
Department of Mathematics
Huzhou University
Huzhou 313000, P. R. China
e-mail: wmk000@126.com; wangmiaokun@zjhu.edu.cn

Yu-Ming Chu
College of Science
Hunan City University
Yiyang 413000, P. R. China
School of Mathematics and Statistics
Changsha University of Science & Technology
Changsha 410114, P. R. China
e-mail: chuyuming2005@126.com

Yong-Min Li
Department of Mathematics
Huzhou University
Huzhou 313000, P. R. China
e-mail: ymlwww@163.com

Wen Zhang
Friedman Brain Institute
Icahn School of Medicine at Mount Sinai
New York NY 10029, USA
e-mail: zhang.wen81@gmail.com