

NORM OF THE DISCRETE CESÀRO OPERATOR MINUS IDENTITY

GORD SINNAMON

(Communicated by J. Jakšetić)

Abstract. The norm of $C - I$ on ℓ^p , where C is the Cesàro operator, is shown to be $1/(p - 1)$ when $1 < p \leq 2$. This verifies a recent conjecture of G. J. O. Jameson. The norm of $C - I$ on ℓ^p is also determined when $2 < p < \infty$. The two parts together answer a question raised by G. Bennett in 1996. Operator norms in the continuous case, Hardy's averaging operator minus identity, are already known. Norms in the discrete and continuous cases coincide.

The Cesàro operator, C , maps a sequence (x_n) to (y_n) , where

$$y_n = \frac{1}{n} \sum_{k=1}^n x_k.$$

Hilbert space methods, see [7], show that the operator norm of $C - I$, as a map on ℓ^2 , is 1. The question of determining the exact norm of $C - I$ as a map on ℓ^p was posed in 1996 by Grahame Bennett as Problem 10.5 in [4]. Recently, Jameson [9] answered the question in the case $p = 4/3$ by showing that $\|C - I\|_{\ell^{4/3}} = 3$. He conjectured that $\|C - I\|_{\ell^p} = 1/(p - 1)$ for $1 < p \leq 2$. In Theorems 2 and 4, Jameson's method for the case $p = 4/3$ is extended and used to verify his conjecture and to answer Bennett's question for the index range $1 < p \leq 2$.

Jameson also gave the upper bound $\|C - I\|_{\ell^4} \leq 3^{1/4}$ for the operator norm of $C - I$ when $p = 4$. In Theorems 8 and 10 we extend the bound to all $p > 2$ and show it is best possible. This completes the following answer to Bennett's question: If $1 < p \leq \infty$, then

$$\|C - I\|_{\ell^p} = \begin{cases} 1/(p - 1), & 1 < p \leq 2; \\ m_p^{-1/p}, & 2 < p < \infty; \\ 2, & p = \infty. \end{cases}$$

Here m_p is the minimum value taken by the function $pt^{p-1} + (1-t)^p - t^p$ on the interval $[0, \frac{1}{2}]$. See Definition 6 and Lemma 7 below. The minimum value m_p is easy to compute when $p = 3$ and $p = 4$; the former gives $\|C - I\|_{\ell^3} = (2 - \sqrt{2})^{-1/3}$ and the latter recovers Jameson's upper bound. The result $\|C - I\|_{\ell^\infty} = 2$ appears in [9, Proposition 1].

Mathematics subject classification (2020): Primary 26D15; Secondary 47B37.

Keywords and phrases: Cesàro, Hardy, operator norm.

Supported by the Natural Sciences and Engineering Research Council of Canada.

Hardy's averaging operator takes a function x on $(0, \infty)$ to $Px(s) = \frac{1}{s} \int_0^s x$. Many results for $P - I$ were first obtained from work on the Beurling-Ahlfors transform on radial functions. For background and references, see [10]. The operator $P - I$ has been studied as a map on $L^p = L^p(0, \infty)$, on the positive cone of L^p , and on the cone of positive, decreasing functions on L^p . Results for weighted L^p , see [5], and weak- L^p , see [6], are also known. For background and additional references, see [11]. Special cases of these general results reveal that the values for $\|C - I\|_{\ell^p}$, stated above, coincide with those for $\|P - I\|_{L^p}$.

In corollaries following the main theorems below, we imitate the proofs for $C - I$ to give analogous proofs for $P - I$. These quick, elementary proofs recover the known values of $\|P - I\|_{L^p}$ in the case $1 < p < 2$ (see [1, Theorem 4.1], [2, Theorem 5.3], and [12, (3.13)]) and the case $2 < p$ (see [10, (1.2)]).

In what follows we consider only real-valued sequences and functions, but, as is well known, extending a linear operator from real to complex values does not change its ℓ^p or L^p operator norm.

In several of the arguments below, it is important that the power functions involved be defined on all of \mathbb{R} . Let

$$\mathbb{E} = \left\{ \frac{2i}{2j+1} : i, j \in \mathbb{Z} \right\} \cap (2, \infty),$$

a dense subset of $[2, \infty)$. If $p \in \mathbb{E}$, the function $t \rightarrow t^p$ is twice continuously differentiable on \mathbb{R} , it is non-negative, its derivative is strictly increasing and the Mean Value Theorem implies that for all $a, b \in \mathbb{R}$,

$$pa^{p-1}(b-a) \leq b^p - a^p \leq pb^{p-1}(b-a). \quad (1)$$

To investigate $C - I$ in the case $1 < p < 2$ it is convenient to work instead with the transpose Cesàro operator, C^T , in the case $2 < p < \infty$. The transpose maps a sequence $(x_n) \in \ell^p$ to the sequence (y_n) , where

$$y_n = \sum_{k=n}^{\infty} \frac{x_k}{k}.$$

Since $\|C - I\|_{\ell^p} = \|C^T - I\|_{\ell^{p'}}$, the conjecture $\|C - I\|_p = p' - 1$ for $1 < p < 2$ may be equivalently stated as $\|C^T - I\|_{\ell^p} = p - 1$ for $p \geq 2$. (Here and throughout, $1/p + 1/p' = 1$).

LEMMA 1. *Let $p \in \mathbb{E}$. For all $t \in \mathbb{R}$,*

$$(p-1)p^{p-2}t^p + (t+1)^p - p(t+1)^{p-1}t \geq p^{p-2}(p-1)^{1-p}.$$

Proof. Since $p > 2$, $p^{p-2} > 1$ so the left-hand side of the inequality goes to infinity as $|t|$ does. Its derivative is $p(p-1)t((pt)^{p-2} - (t+1)^{p-2})$, which vanishes if and only if $t = 0$ or $t+1 = \pm pt$. This derivative goes from positive to negative as t crosses 0 so there is a local maximum at $t = 0$. The other critical points are $t = 1/(p-1)$ and $t = -1/(p+1)$, so the minimum value taken by the left-hand side is the smaller of,

$$p^{p-2}(p-1)^{1-p} \quad \text{and} \quad p^{p-2}(2p-1)(p+1)^{1-p}.$$

To complete the proof we show that the first of these is smaller. This is equivalent to showing that $h(p) = (p - 1)(\log(p + 1) - \log(p - 1)) - \log(2p - 1) \leq 0$. Since $h(2) = 0$ it suffices to show that $h'(p) \leq 0$ for $p > 2$. Since $\frac{1}{s}$ lies below its secant on $[p - 1, p + 1]$, we have $\frac{1}{s} \leq \frac{2p-s}{p^2-1}$. Thus $\log(p + 1) - \log(p - 1) \leq \int_{p-1}^{p+1} \frac{2p-s}{p^2-1} ds = \frac{2p}{p^2-1}$. Also, $2p - 1 \leq p^2 - 1$ so

$$h'(p) \leq \frac{2p}{p^2-1} + (p-1)\left(\frac{1}{p+1} - \frac{1}{p-1}\right) - \frac{2}{p^2-1} = 0. \quad \square$$

Now we are ready to show that $\|C^T - I\|_{\ell^p} \leq p - 1$ for $p \in \mathbb{E}$. The proof follows the method of [9, Theorem 2].

THEOREM 2. *Let $p \in \mathbb{E}$, let $(x_n) \in \ell^p$ be a real sequence, and set $y_n = \sum_{k=n}^{\infty} \frac{x_k}{k}$. Then*

$$\sum_{n=1}^{\infty} (y_n - x_n)^p \leq (p - 1)^p \sum_{n=1}^{\infty} x_n^p.$$

Proof. By Hölder's inequality, the sum defining y_N converges absolutely and

$$|y_N| \leq \sum_{n=N}^{\infty} \frac{|x_n|}{n} \leq \left(\sum_{n=N}^{\infty} x_n^p\right)^{1/p} \left(\sum_{n=N}^{\infty} n^{-p'}\right)^{1/p'} \sim \left(\sum_{n=N}^{\infty} x_n^p\right)^{1/p} N^{-1/p}.$$

Therefore, as $N \rightarrow \infty$,

$$\sum_{n=1}^N (ny_{n+1}^p - (n-1)y_n^p) = Ny_{N+1}^p \rightarrow 0.$$

Note that for all n , $x_n = n(y_n - y_{n+1})$. For each n , (1) implies

$$ny_{n+1}^p - (n-1)y_n^p = y_n^p - n(y_n^p - y_{n+1}^p) \geq y_n^p - py_n^{p-1}n(y_n - y_{n+1}) = y_n^p - py_n^{p-1}x_n.$$

Now let $z_n = y_n - x_n$ and set $c = p^{p-2}(p-1)^{1-p}$. If $z_n \neq 0$ we may let $t = x_n/z_n$ and apply Lemma 1 in the form $(t+1)^p - p(t+1)^{p-1}t \geq c(1 - (p-1)^p t^p)$ to get

$$y_n^p - py_n^{p-1}x_n = z_n^p((1+t)^p - p(1+t)^{p-1}t) \geq cz_n^p(1 - (p-1)^p t^p) = c(z_n^p - (p-1)^p x_n^p).$$

If $z_n = 0$, then $y_n = x_n$ and the same inequality holds, as

$$y_n^p - py_n^{p-1}x_n = -(p-1)x_n^p \geq -p^{p-2}(p-1)x_n^p = c(z_n^p - (p-1)^p x_n^p).$$

Summing over n , we have

$$c \sum_{n=1}^N (z_n^p - (p-1)^p x_n^p) \leq \sum_{n=1}^N (ny_{n+1}^p - (n-1)y_n^p) \rightarrow 0$$

as $N \rightarrow \infty$ and we conclude that

$$\sum_{n=1}^{\infty} z_n^p \leq (p-1)^p \sum_{n=1}^{\infty} x_n^p. \quad \square$$

The proof simplifies in the continuous case, where instead of the transpose Cesàro operator we work with the dual averaging operator P^T .

COROLLARY 3. *Let $p \in \mathbb{E}$, let $x \in L^p$, and set $y(s) = P^T x(s) = \int_s^\infty x(\theta) \frac{d\theta}{\theta}$. Then $\int_0^\infty (y-x)^p \leq (p-1)^p \int_0^\infty x^p$.*

Proof. It suffices to establish the result for functions x in a dense subset of L^p so suppose x is continuous and compactly supported in $(0, \infty)$. Then $y = P^T x$ satisfies $\lim_{s \rightarrow 0^+} sy(s)^p = 0 = \lim_{s \rightarrow \infty} sy(s)^p$ and $\frac{d}{ds}(sy(s)^p) = y(s)^p - py(s)^{p-1}x(s)$ so $\int_0^\infty (y^p - py^{p-1}x) = 0$. Let $z = y-x$ and $c = p^{p-2}(p-1)^{1-p}$. If $z \neq 0$ and $t = x/z$, then Lemma 1 implies

$$c(z^p - (p-1)^p x^p) = cz^p(1 - (p-1)^p t^p) \leq z^p((1+t)^p - p(1+t)^{p-1}t) = y^p - py^{p-1}x,$$

which also holds when $z = 0$. Integrate to get $\int_0^\infty z^p \leq (p-1)^p \int_0^\infty x^p$. \square

Next we extend Theorem 2 from $p \in \mathbb{E}$ to all $p > 2$ and point out a known lower bound for $\|C^T - I\|_{\ell^p}$.

THEOREM 4. *If $2 \leq p < \infty$ then $\|C^T - I\|_{\ell^p} = p-1$ and if $1 < p \leq 2$ then $\|C - I\|_{\ell^p} = p' - 1 = 1/(p-1)$.*

Proof. As mentioned above, the case $p=2$ is known to hold. For $p > 2$, Theorem 2 shows that for $p \in \mathbb{E}$ with $2 < p < \infty$, $\|C^T - I\|_{\ell^p} \leq p-1$. The Riesz-Thorin Theorem, see [3, Corollary IV.2.3], implies that if $2 < p_0 < p < p_1 < \infty$, with $p_0, p_1 \in \mathbb{E}$, then for some $\theta \in (0, 1)$, depending on p_0, p , and p_1 ,

$$\|C^T - I\|_{\ell^p} \leq \|C^T - I\|_{\ell^{p_0}}^{1-\theta} \|C - I\|_{\ell^{p_1}}^\theta = (p_0 - 1)^{1-\theta} (p_1 - 1)^\theta \leq p_1 - 1.$$

Letting $p_1 \rightarrow p$ through \mathbb{E} , we get $\|C^T - I\|_{\ell^p} \leq p-1$.

The dual discrete Hardy inequality, [8, Theorem 331], shows that for $p > 1$, $\|C^T\|_{\ell^p} = p$. Therefore, $\|C^T - I\|_{\ell^p} \geq \|C^T\|_{\ell^p} - \|I\|_{\ell^p} = p-1$ and we conclude that $\|C^T - I\|_{\ell^p} = p-1$ for all $p > 2$. The second statement of the theorem follows from the first by duality. \square

The continuous case follows in just the same way because $\|P^T\|_{L^p} = p$. See [8, Theorem 328]. The proof is omitted.

COROLLARY 5. *If $2 \leq p < \infty$ then $\|P^T - I\|_{L^p} = p-1$ and if $1 < p \leq 2$ then $\|P - I\|_{L^p} = p' - 1 = 1/(p-1)$.*

Next we consider the case $p > 2$. To begin we introduce m_p , essential for our formula for the operator norm of $C - I$.

DEFINITION 6. Let $p \geq 2$ and set $f_p(t) = pt^{p-1} + (1-t)^p - t^p$. Define m_p to be the minimum value of $f_p(t)$ for $0 \leq t \leq \frac{1}{2}$.

LEMMA 7. *If $p > 2$, then f_p has a unique critical point t_p in $(0, \frac{1}{2})$, $m_p = f_p(t_p)$ and m_p is a continuous function of p . If, in addition, $p \in \mathbb{E}$, then t_p is the unique critical point of f_p on all of \mathbb{R} and $f_p(t) \geq m_p$ for all $t \in \mathbb{R}$.*

Proof. On $(0, \frac{1}{2})$, $f'_p(t) = p((p-1)t^{p-2} - (1-t)^{p-1} - t^{p-1})$. It extends to be continuous on $[0, \frac{1}{2}]$ with $f'_p(0) = -p < 0$, and $f'_p(\frac{1}{2}) = p(p-2)2^{2-p} > 0$. On $(0, \frac{1}{2})$, $0 < t < 1-t$ so $f''_p(t) = p(p-1)((p-2)t^{p-3} + (1-t)^{p-2} - t^{p-2}) > 0$.

Therefore f'_p is strictly increasing on $[0, \frac{1}{2}]$, f_p has a unique critical point t_p in $(0, \frac{1}{2})$ and $f_p(t_p)$ is the minimum value of f_p , namely m_p . For any $p_0 > 2$, the function $(p, t) \mapsto f_p(t)$ is uniformly continuous on $[2, p_0] \times [0, \frac{1}{2}]$. It follows that $p \mapsto m_p$ is continuous on $[2, \infty)$.

If $p \in \mathbb{E}$, then f_p is defined on \mathbb{R} , $(1-t)^p = (t-1)^p$, and

$$f'_p(t) = p((p-1)t^{p-2} + (t-1)^{p-1} - t^{p-1}) = p(p-1) \int_{t-1}^t (t^{p-2} - s^{p-2}) ds.$$

If $t \leq 0$ then $t-1 < s < t$ implies $|t| < |s|$ so $t^{p-2} < s^{p-2}$ and we have $f'_p(t) < 0$. Thus, f_p is strictly decreasing on $(-\infty, 0]$. If $t \geq \frac{1}{2}$ then $t-1 < s < t$ implies $|s| < |t|$ so $s^{p-2} < t^{p-2}$ and we have $f'_p(t) > 0$. Thus f_p is strictly increasing on $[\frac{1}{2}, \infty)$. It follows that t_p is the unique critical point of f_p on \mathbb{R} and $f_p(t) \geq m_p$ for all $t \in \mathbb{R}$. \square

In [9, Theorem 1], the upper bound $\|C - I\|_{\ell^4} \leq 3^{1/4}$ was proved. We employ a similar method to extend it to an upper bound for all $p > 2$.

THEOREM 8. *Let $p \in \mathbb{E}$, let $(x_n) \in \ell^p$ be a real sequence, and set $y_n = \frac{1}{n} \sum_{k=1}^n x_k$. Then*

$$\sum_{k=1}^{\infty} (y_k - x_k)^p \leq \frac{1}{m_p} \sum_{k=1}^{\infty} x_k^p.$$

Proof. Fix a y_0 arbitrarily and observe that $(n-1)(y_{n-1} - y_n) = y_n - x_n$ for $n = 1, 2, \dots$. By (1),

$$(n-1)(y_{n-1}^p - y_n^p) \geq (n-1)py_n^{p-1}(y_{n-1} - y_n) = py_n^{p-1}(y_n - x_n).$$

This becomes

$$ny_n^p - (n-1)y_{n-1}^p \leq py_n^{p-1}x_n - (p-1)y_n^p = y_n^{p-1}(px_n - (p-1)y_n).$$

Take $z_n = y_n - x_n$ and $t = y_n/z_n$ to get $px_n - (p-1)y_n = z_n(t-p)$. By Lemma 7,

$$ny_n^p - (n-1)y_{n-1}^p \leq z_n^p(t^p - pt^{p-1}) \leq z_n^p((t-1)^p - m_p) = x_n^p - m_p z_n^p.$$

Summing from 1 to N gives

$$0 \leq Ny_N^p = \sum_{n=1}^N (ny_n^p - (n-1)y_{n-1}^p) \leq \sum_{n=1}^N x_n^p - m_p \sum_{n=1}^N z_n^p.$$

Letting $N \rightarrow \infty$ we have

$$\sum_{n=1}^{\infty} z_n^p \leq \frac{1}{m_p} \sum_{n=1}^{\infty} x_n^p. \quad \square$$

COROLLARY 9. Let $p \in \mathbb{E}$, let $x \in L^p$ and set $y(s) = Px(s) = \frac{1}{s} \int_0^s x$. Then $\int_0^\infty (y - x)^p \leq \frac{1}{m_p} \int_0^\infty x^p$.

Proof. It suffices to establish the result for functions x in a dense subset of L^p so suppose x is continuous and compactly supported in $(0, \infty)$. Then $y = Px$ satisfies $\lim_{s \rightarrow 0^+} sy(s)^p = 0 = \lim_{s \rightarrow \infty} sy(s)^p$ and $\frac{d}{ds}(y(s)^p) = (1 - p)y(s)^{p-1} + py(s)^{p-1}x(s)$ so $\int_0^\infty ((1 - p)y^p + py^{p-1}x) = 0$. Let $z = y - x$. If $z \neq 0$ and $t = y/z$, then Lemma 7 implies

$$x^p - m_p z^p = z^p((t - 1)^p - m_p) \geq z^p(t^p - pt^{p-1}) = (1 - p)y^p + py^{p-1}x,$$

which also holds when $z = 0$. Integrate to get $\int_0^\infty z^p \leq \frac{1}{m_p} \int_0^\infty x^p$. \square

We again pass from $p \in \mathbb{E}$ to all $p > 2$ using the Riesz-Thorin Theorem. However, this time the lower bound requires some work.

THEOREM 10. If $p \geq 2$ then $\|C - I\|_{\ell^p} = m_p^{-1/p}$.

Proof. It is easy to verify that $m_2 = 1$, so the case $p = 2$ agrees with the known result. Now suppose $p > 2$. Theorem 8 shows that $\|C - I\|_{\ell^p} \leq m_p^{-1/p}$ for all $p \in \mathbb{E}$. The Riesz-Thorin theorem implies that if $2 < p_0 < p < p_1 < \infty$, with $p_0, p_1 \in \mathbb{E}$, then for some $\theta \in (0, 1)$, depending on p_0, p , and p_1 ,

$$\|C - I\|_{\ell^p} \leq \|C - I\|_{\ell^{p_0}}^{1-\theta} \|C - I\|_{\ell^{p_1}}^\theta \leq \max(m_{p_0}^{-1/p_0}, m_{p_1}^{-1/p_1}).$$

Letting p_0 and p_1 approach p through \mathbb{E} , the continuity of $p \mapsto m_p$ implies that $\|C - I\|_{\ell^p} \leq m_p^{-1/p}$.

To prove the reverse inequality we set $r = 1/t_p$, where t_p is the critical point from Lemma 7, and fix an integer $m > 1$. Note that $r > 2$. Define $x_n = -m^{-r}$ for $n \leq m$ and $x_n = (n - 1)^{1-r} - n^{1-r}$ for $n > m$. Then, with $y_n = \frac{1}{n} \sum_{k=1}^n x_k$ and $z_n = y_n - x_n$,

$$y_n = \begin{cases} -m^{-r}, & n \leq m; \\ -n^{-r}, & n \geq m \end{cases} \quad \text{and} \quad z_n = \begin{cases} 0, & n \leq m; \\ n^{1-r} - (n - 1)^{1-r} - n^{-r}, & n > m \end{cases}$$

If $n \geq m + 1$, then

$$0 < x_n = (r - 1) \int_{n-1}^n t^{-r} dt \leq (r - 1)(n - 1)^{-r}$$

so, employing a standard Riemann sum estimate, we get

$$\sum_{n=1}^\infty |x_n|^p \leq m^{1-pr} + (r - 1)^p \sum_{n=m+1}^\infty (n - 1)^{-pr} \leq m^{1-pr} + (r - 1)^p \frac{(m - 1)^{1-pr}}{pr - 1}.$$

Also, if $n \geq m + 1$, then $\frac{n-1}{n} \geq \frac{m}{m+1}$ and

$$-z_n = r(n - 1) \int_{n-1}^n t^{-r-1} dt \geq r(n - 1)n^{-r-1} \geq r \frac{m}{m + 1} n^{-r} > 0$$

so, using another standard Riemann sum estimate, we get

$$\sum_{n=1}^{\infty} |z_n|^p \geq r^p \left(\frac{m}{m+1}\right)^p \sum_{n=m+1}^{\infty} n^{-pr} \geq r^p \left(\frac{m}{m+1}\right)^p \frac{(m+1)^{1-pr}}{pr-1}.$$

We conclude that

$$\frac{\sum_{n=1}^{\infty} |z_n|^p}{\sum_{n=1}^{\infty} |x_n|^p} \geq \frac{r^p \left(\frac{m}{m+1}\right)^p}{(pr-1)\left(\frac{m}{m+1}\right)^{1-pr} + (r-1)^p \left(\frac{m-1}{m+1}\right)^{1-pr}} \rightarrow \frac{r^p}{(pr-1) + (r-1)^p}$$

as $m \rightarrow \infty$. Since $r = 1/t_p$, Lemma 7 shows that the last expression is $1/m_p$. This implies that the operator norm of $C - I$ on ℓ^p cannot be less than $m_p^{-1/p}$, which completes the proof. \square

COROLLARY 11. *If $p \geq 2$ then $\|P - I\|_{L^p} = m_p^{-1/p}$.*

Proof. The upper bound is extended from $p \in \mathbb{E}$ to all $p > 2$ just as in the last theorem. However, proving the reverse inequality is much simpler. With t_p as in Lemma 7 and $r = 1/t_p$, let $x(s) = -1$ on $(0, 1)$ and $x(s) = (r-1)s^{-r}$ on $(1, \infty)$. With $y = Px$ and $z = y - x$ we compute $y(s) = -1$ on $(0, 1)$ and $y(s) = -s^{-r}$ on $(1, \infty)$; and $z(s) = 0$ on $(0, 1)$ and $z(s) = -rs^{-r}$ on $(1, \infty)$. Then

$$\int_0^{\infty} |x|^p = 1 + \frac{(r-1)^p}{pr-1} \quad \text{and} \quad \int_0^{\infty} |z|^p = \frac{r^p}{pr-1}.$$

It follows that $\int_0^{\infty} |z|^p = \frac{1}{m_p} \int_0^{\infty} |x|^p$, which gives the lower bound. \square

The expressions for $\|P - I\|_{L^p}$ given above and in [10] must coincide, but a direct connection is still worth making: In [10], we find $\|P - I\|_{L^p}^p = \sup_{\alpha \leq p'} \frac{|\alpha-1|^p}{p(1-\alpha)-1+|\alpha|^p}$. With $t = 1/(1-\alpha)$ this expression readily reduces to $1/m_p$ for $p \in \mathbb{E}$, by applying Lemma 7. Equality for all p follows by continuity.

Acknowledgements. My thanks to Santiago Boza for providing references to results in the continuous case and to Graham Jameson for an improvement to the proof of Lemma 1.

REFERENCES

- [1] R. BAÑUELOS AND P. JANAKIRAMAN, *On the weak-type constant of the Beurling-Ahlfors transform*, Michigan Math. J. **58**, 2 (2009), 459–477.
- [2] R. BAÑUELOS AND A. OŚĘKOWSKI, *Sharp inequalities for the Beurling-Ahlfors transform on radial functions*, Duke Math. J. **162**, 2 (2013), 417–434.
- [3] C. BENNETT AND R. SHARPLEY, *Interpolation of operators*, Academic Press, Inc., Pure and Applied Mathematics Series **129**, Washington, 1988.
- [4] G. BENNETT, *Factorizing the classical inequalities*, Mem. Amer. Math. Soc. **120**, 1 (1996), 1–130.
- [5] S. BOZA AND J. SORIA, *Solution to a conjecture on the norm of the Hardy operator minus the identity*, J. Funct. Anal. **260**, 4 (2011), 1020–1028.

- [6] S. BOZA AND J. SORIA, *Averaging operators on decreasing or positive functions: equivalence and optimal bounds*, J. Approx. Theory **237**, 1 (2019), 135–152.
- [7] A. BROWN, P. R. HALMOS AND A. L. SHIELDS, *Cesàro operators*, Acta Sci. Math. (Szeged) **26**, 1 (1965), 125–137.
- [8] G. H. HARDY, J. E. LITTLEWOOD, AND G. PÓLYA, *Inequalities*, Reprint of the 1952 edition, Cambridge University Press, Cambridge Mathematical Library Series, Cambridge, 1988.
- [9] G. J. O. JAMESON, *The ℓ_p -norm of $C-I$, where C is the Cesàro operator*, Math. Inequal. Appl. **24**, 2 (2021), 551–557.
- [10] M. STRZELECKI, *The L^p -norms of the Beurling-Ahlfors transform on radial functions*, Ann. Acad. Sci. Fenn. Math. **42**, 1 (2017), 73–93.
- [11] M. STRZELECKI, *Hardy's operator minus identity and power weights*, J. Funct. Anal. **279**, 2 (2020), 108532, 34.
- [12] A. VOLBERG, *Ahlfors-Beurling operator on radial functions*, arXiv:1203.2291 **1**, 1 (2012), 16.

(Received May 29, 2021)

Gord Sinnamon
Department of Mathematics
Western University
1151 Richmond Street, London, Ontario, Canada, N6A 3K7
e-mail: sinnamon@uwo.ca