

HARDY AND SOBOLEV INEQUALITIES FOR DOUBLE PHASE FUNCTIONALS ON THE UNIT BALL

YOSHIHIRO MIZUTA AND TETSU SHIMOMURA *

(Communicated by S. Varošanec)

Abstract. We prove Hardy and Sobolev inequalities for double phase functionals $\Phi(x, t) = t^p + (b(x)t)^q$ on the unit ball \mathbf{B} , as a continuation of our paper [26], where $1 \leq p < q$, $b(\cdot)$ is non-negative and (radially) Hölder continuous of order $\theta \in (0, 1]$. The Sobolev conjugate for Φ is given by $\Phi^*(x, t) = t^{p^*} + (b(x)t)^{q^*}$, where p^* and q^* denote the Sobolev exponent of p and q , respectively, that is, $1/p^* = 1/p - 1/n$ and $1/q^* = 1/q - 1/n$.

1. Introduction

The Hardy-Sobolev inequality says that for $f \geq 0$

$$\left(\int_0^1 \left(\int_0^x f(y) dy \right)^q (1-x)^\alpha dx \right)^{1/q} \leq C \left(\int_0^1 f(y)^p (1-y)^\beta dy \right)^{1/p}, \quad (1)$$

where $1 \leq p \leq q$, $\beta > p - 1$ and $\alpha = \beta q/p - q/p' - 1$ ($1/p + 1/p' = 1$) and

$$\left(\int_0^1 \left(\int_x^1 f(y) dy \right)^q (1-x)^\alpha dx \right)^{1/q} \leq C \left(\int_0^1 f(y)^p (1-y)^\beta dy \right)^{1/p}, \quad (2)$$

where $1 \leq p \leq q$, $\beta < p - 1$ and $\alpha = \beta q/p - q/p' - 1$ (see e.g. [10, 15, 16, 21, 22, 28]).

The double phase functional introduced by Zhikov ([31]) in the 1980s has been studied intensively by many researchers. Regarding regularity theory of differential equations, Baroni, Colombo and Mingione [1, 2, 5, 6] studied a double phase functional

$$\tilde{\Phi}(x, t) = t^p + a(x)t^q, \quad x \in \mathbb{R}^n, \quad t \geq 0,$$

where $1 \leq p < q$, $a(\cdot)$ is non-negative, bounded and Hölder continuous of order $\theta \in (0, 1]$. We refer to [17, 24, 25] for Sobolev inequality for double phase functionals and to [19, 20] for variational problems with nonstandard growth. For other recent works, see e.g. [3, 4, 7, 8, 9, 11, 12, 13, 14, 23, 27, 29].

Mathematics subject classification (2020): 46E30, 26D10, 47G10.

Keywords and phrases: Hardy-Sobolev inequality, double phase functionals.

* Corresponding author.

Let \mathbf{B} be the unit ball $B(0, 1)$ of \mathbb{R}^n . In the present paper, relaxing the continuity of $a(\cdot)$, we consider the case $\Phi(x, t)$ is a double phase functional given by

$$\Phi(x, t) = t^p + (b(x)t)^q,$$

where $1 \leq p < q$, $b(\cdot)$ is non-negative and (radially) Hölder continuous of order $\theta \in (0, 1]$, that is,

$$|b(x) - b(y)| \leq C||x| - |y||^\theta \quad \text{for all } x, y \in \mathbf{B}.$$

Note that if we write

$$\Phi(x, t) = t^p + a(x)t^q$$

with

$$a(x) = b(x)^q,$$

then a is not always Hölder continuous of order θq .

In the previous paper [26], Hardy-Sobolev inequalities were established when

$$\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{\beta/q} |f(y)|) dy \leq 1 \tag{3}$$

with $\beta > q - 1$ and $1/q = 1/p - \theta > 0$ and

$$\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{\beta/p} |f(y)|) dy \leq 1 \tag{4}$$

with $\beta < p - 1$.

The Sobolev conjugate Φ^* for Φ will be given in Section 3 by

$$\Phi^*(x, t) = t^{p^*} + (b(x)t)^{q^*},$$

where p^* and q^* denote the Sobolev exponent of p and q , respectively, that is, $1/p^* = 1/p - 1/n$ and $1/q^* = 1/q - 1/n$. Our aim in this paper is to give a continuation of our paper [26] by the use of $\Phi^*(x, t)$, which was not used in [26], when (3) holds with $\beta > q - 1$ and $1/q = 1/p - \theta > 0$ in Theorem 3.1 and (4) holds with $\beta < p - 1$ in Theorem 3.3. Our strategy is to apply Theorems 2.1 and 2.4, which are extensions of the classical Hardy-Sobolev inequalities (1) and (2) to the unit ball. which are the Hardy-Sobolev inequality in \mathbf{B} . Both Theorems 3.1 and 3.3 are strongly affected by double phase as will be seen in Remark 4.1 below.

As an application of Theorems 2.1, 2.4, 3.1 and 3.3, we discuss behaviors near the sphere for functions in $C^1(\mathbb{R}^n)$ (see Propositions given after the theorems).

Our final aim is to treat the borderline case in Theorems 3.1 and 3.3. For this purpose, we prepare Theorems 5.1 and 5.4, which are the critical case of Theorems 2.1 and 2.4.

Throughout this paper, let C denote various positive constants independent of the variables in question. The symbol $g \sim h$ means that $C^{-1}h \leq g \leq Ch$ for some constant $C > 0$.

2. Hardy-Sobolev inequality in \mathbf{B}

Let $B(x, r)$ denote the open ball centered at $x \in \mathbf{B}$ with radius r .

In \mathbf{B} , Hardy-Sobolev inequality can be stated in the following (cf. [28]):

THEOREM 2.1. ([26, Theorem 2.1]) *Let $1 \leq p \leq q$, $\beta > p - 1$ and $\alpha = \beta q/p - q/p' - 1$, where $1/p + 1/p' = 1$. Then there exists a constant $C > 0$ such that*

$$\left(\int_{\mathbf{B}} \left(\int_{B(0,|x|)} |f(y)| dy \right)^q (1 - |x|)^\alpha dx \right)^{1/q} \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^\beta dy \right)^{1/p}.$$

By Theorem 2.1 we discuss a behavior near the sphere.

PROPOSITION 2.2. (cf. [18, Theorem 6.5], [22, Theorem 4.3.1], [30]) *Let $1 \leq p \leq q$ and $0 < \varepsilon < 1/q$. Let u be a function in $C^1(\mathbb{R}^n)$ such that*

$$\int_{\mathbf{B}} (|\nabla u(y)| (1 - |y|)^\varepsilon)^p (1 - |y|)^{p-1} dy < \infty.$$

Then

$$\lim_{r \rightarrow 1} (1 - r)^\varepsilon \frac{1}{|\mathbf{B} \setminus B(0, r)|} \int_{\mathbf{B} \setminus B(0, r)} |u(x)| dx = 0$$

and

$$\lim_{r \rightarrow 1} (1 - r)^\varepsilon \left(\frac{1}{|S(0, r)|} \int_{S(0, r)} |u(r\xi)| dS(\xi) \right) = 0,$$

where $|S(0, r)|$ denotes the surface area of the spherical surface $S(0, r)$.

Proof. First note

$$\begin{aligned} U_0(r\xi) &:= |u(r\xi) - u(0)| \\ &= \left| \int_0^r \frac{d}{dt} u(t\xi) dt \right| \\ &\leq \int_0^r |\nabla u(t\xi)| dt \end{aligned}$$

for $0 < r < 1$ and $|\xi| = 1$. Then

$$\frac{1}{|S(0, 1)|} \int_{S(0, 1)} U_0(r\xi) dS(\xi) \leq \frac{1}{|S(0, 1)|} \int_{B(0, r)} |\nabla u(y)| |y|^{1-n} dy. \tag{5}$$

Thus for $r > 0$ Theorem 2.1 gives

$$\begin{aligned}
 & (1-r)^\varepsilon \frac{1}{1-r} \int_r^1 \left(\frac{1}{|S(0,1)|} \int_{S(0,1)} U_0(t\xi) dS(\xi) \right) dt \\
 & \leq (1-r)^\varepsilon \left(\frac{1}{1-r} \int_r^1 \left(\frac{1}{|S(0,1)|} \int_{B(0,t)} |\nabla u(y)| |y|^{1-n} dy \right)^q dt \right)^{1/q} \\
 & \leq C \left(\int_{\mathbf{B} \setminus B(0,r)} \left(\int_{B(0,t)} |\nabla u(y)| |y|^{1-n} dy \right)^q (1-|x|)^\alpha |x|^{1-n} dx \right)^{1/q} \\
 & \leq Cr^{(1-n)/q} \left(\int_{\mathbf{B} \setminus B(0,r)} (|\nabla u(y)| |y|^{1-n})^p (1-|y|)^\beta dy \right)^{1/p} \\
 & \leq C \int_{\mathbf{B} \setminus B(0,r)} (|\nabla u(y)| (1-|y|)^\varepsilon)^p (1-|y|)^{p-1} dy
 \end{aligned}$$

since $(1-n)/q + 1 - n < 0$, where $\beta = (p-1) + \varepsilon p > p-1$ and $\alpha = \beta q/p - q/p' - 1 = \varepsilon q - 1 < 0$. Here note

$$\begin{aligned}
 & (1-r)^\varepsilon \frac{1}{1-r} \int_r^1 \left(\frac{1}{|S(0,1)|} \int_{S(0,1)} U_0(t\xi) dS(\xi) \right) dt \\
 & \sim (1-r)^\varepsilon \frac{1}{|\mathbf{B} \setminus B(0,r)|} \int_{\mathbf{B} \setminus B(0,r)} U_0(x) dx \quad \text{for } 0 < r < 1,
 \end{aligned}$$

which yields

$$\lim_{r \rightarrow 1} (1-r)^\varepsilon \frac{1}{|\mathbf{B} \setminus B(0,r)|} \int_{\mathbf{B} \setminus B(0,r)} U_0(x) dx = 0.$$

Moreover, it is seen from (5) that

$$\lim_{r \rightarrow 1} (1-r)^\varepsilon \left(\frac{1}{|S(0,r)|} \int_{S(0,r)} U_0(r\xi) dS(\xi) \right) = 0.$$

Noting that $|u(x)| \leq |U_0(x)| + |u(0)|$, we establish the required result. \square

REMARK 2.3. Consider the function

$$u(x) = (1-|x|)^{-a} - 1$$

for $a > 0$. Then

(1) $|\nabla u(x)| = a(1-|x|)^{-a-1}$ and

$$\int_{\mathbf{B}} (|\nabla u(x)| (1-|x|)^\varepsilon)^p (1-|x|)^{p-1} dx < \infty$$

if and only if $(-a-1+\varepsilon)p + p > 0$;

(2)

$$\lim_{r \rightarrow 1} (1-r)^\varepsilon \left(\frac{1}{|S(0,r)|} \int_{S(0,r)} u(r\xi) dS(\xi) \right) = 0$$

if and only if $-a + \varepsilon > 0$.

This implies the best possibility as to the power ε in Proposition 2.2.

THEOREM 2.4. ([26, Theorem 2.2]) *Let $1 \leq p \leq q$, $\beta < p - 1$ and $\alpha = \beta q/p - q/p' - 1$. Then there exists a constant $C > 0$ such that*

$$\left(\int_{\mathbf{B}} \left(\int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| dy \right)^q (1-|x|)^\alpha dx \right)^{1/q} \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1-|y|)^\beta dy \right)^{1/p}.$$

By Theorem 2.4 we discuss a behavior near the sphere, as in Proposition 2.2.

PROPOSITION 2.5. *Let $p \geq 1$ and $\varepsilon > 0$. Let u be a function in $C^1(\mathbb{R}^n)$ such that*

$$\int_{\mathbf{B}} (|\nabla u(y)| (1-|y|)^{-\varepsilon})^p (1-|y|)^{p-1} dy < \infty.$$

Set

$$U_1(x) = U_1(r\xi) = \limsup_{t \rightarrow 1} |u(r\xi) - u(t\xi)|.$$

Then

$$\lim_{r \rightarrow 1} (1-r)^{-\varepsilon} \frac{1}{|\mathbf{B} \setminus B(0,r)|} \int_{\mathbf{B} \setminus B(0,r)} U_1(x) dx = 0$$

and

$$\lim_{r \rightarrow 1} (1-r)^{-\varepsilon} \left(\frac{1}{|S(0,r)|} \int_{S(0,r)} U_1(r\xi) dS(\xi) \right) = 0.$$

Proof. To show this, note

$$\begin{aligned} |u(r\xi) - u(t\xi)| &= \left| \int_r^t \frac{d}{dt} u(t\xi) dt \right| \\ &\leq \int_r^t |\nabla u(t\xi)| dt \end{aligned}$$

for $0 < r < t < 1$ and $|\xi| = 1$. Then

$$\frac{1}{|S(0,1)|} \int_{S(0,1)} U_1(r\xi) dS(\xi) \leq \frac{1}{|S(0,1)|} \int_{\mathbf{B} \setminus B(0,r)} |\nabla u(y)| |y|^{1-n} dy. \tag{6}$$

Thus for $r > 0$ Theorem 2.4 gives

$$\begin{aligned} & (1-r)^{-\varepsilon} \frac{1}{1-r} \int_r^1 \left(\frac{1}{|S(0,1)|} \int_{S(0,1)} U_1(t\xi) dS(\xi) \right) dt \\ & \leq (1-r)^{-\varepsilon} \left(\frac{1}{1-r} \int_r^1 \left(\frac{1}{|S(0,1)|} \int_{\mathbf{B} \setminus B(0,t)} |\nabla u(y)| |y|^{1-n} dy \right)^q dt \right)^{1/q} \\ & \leq C \left(\int_{\mathbf{B} \setminus B(0,r)} \left(\int_{\mathbf{B} \setminus B(0,t)} |\nabla u(y)| |y|^{1-n} dy \right)^q (1-|x|)^\alpha |x|^{1-n} dx \right)^{1/q} \\ & \leq Cr^{(1-n)/q} \left(\int_{\mathbf{B} \setminus B(0,r)} (|\nabla u(y)| |y|^{1-n})^p (1-|y|)^\beta dy \right)^{1/p} \\ & \leq C \int_{\mathbf{B} \setminus B(0,r)} (|\nabla u(y)| (1-|y|)^{-\varepsilon})^p (1-|y|)^{p-1} dy, \end{aligned}$$

where $1 \leq p \leq q$, $\beta = (p-1) - \varepsilon p < p-1$ and $\alpha = \beta q/p - q/p' - 1 = -\varepsilon q - 1$. Here note

$$\begin{aligned} & (1-r)^{-\varepsilon} \frac{1}{1-r} \int_r^1 \left(\frac{1}{|S(0,1)|} \int_{S(0,1)} U_1(t\xi) dS(\xi) \right) dt \\ & \sim (1-r)^{-\varepsilon} \frac{1}{|\mathbf{B} \setminus B(0,r)|} \int_{\mathbf{B} \setminus B(0,r)} U_1(x) dx \quad \text{for } 0 < r < 1, \end{aligned}$$

which yields

$$\lim_{r \rightarrow 1} (1-r)^{-\varepsilon} \frac{1}{|\mathbf{B} \setminus B(0,r)|} \int_{\mathbf{B} \setminus B(0,r)} U_1(x) dx = 0.$$

Moreover, it is seen from (6) that

$$\lim_{r \rightarrow 1} (1-r)^{-\varepsilon} \left(\frac{1}{|S(0,r)|} \int_{S(0,r)} U_1(r\xi) dS(\xi) \right) = 0. \quad \square$$

REMARK 2.6. Consider the function

$$u(x) = (1-|x|)^a$$

for $a > 0$. Then

(1) $|\nabla u(x)| = a(1-|x|)^{a-1}$ and

$$\int_{\mathbf{B}} (|\nabla u(x)| (1-|x|)^{-\varepsilon})^p (1-|x|)^{p-1} dx < \infty$$

if and only if $(a-1-\varepsilon)p + p > 0$;

(2)

$$\lim_{r \rightarrow 1} (1-r)^{-\varepsilon} \left(\frac{1}{|S(0,r)|} \int_{S(0,r)} u(r\xi) dS(\xi) \right) = 0$$

if and only if $a - \varepsilon > 0$.

This implies the best possibility as to the power ε in Proposition 2.5.

3. Hardy-Sobolev inequality in \mathbf{B} for double phase functionals

In this section, we give Hardy-Sobolev inequality in \mathbf{B} when Φ is a double phase functional.

Let p^* and q^* denote the Sobolev exponent of p and q , respectively, that is, $1/p^* = 1/p - 1/n$ and $1/q^* = 1/q - 1/n$. The Sobolev conjugate for Φ is given by

$$\Phi^*(x, t) = t^{p^*} + (b(x)t)^{q^*}.$$

THEOREM 3.1. *Let $1 \leq p < q < n$, $\beta > q - 1$ and $1/q = 1/p - \theta > 0$. Set $F(x) = \int_{B(0,|x|)} |f(y)|dy$. Then there exists a constant $C > 0$ such that*

$$\int_{\mathbf{B}} \Phi^* \left(x, (1 - |x|)^{\beta/q-1/n'} F(x) \right) dx \leq C$$

when $\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{\beta/q}|f(y)|)dy \leq 1$.

Proof. First note from Theorem 2.1

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{\beta/q-1/n'} \int_{B(0,|x|)} |f(y)|dy \right)^{p^*} dx \right)^{1/p^*} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{\beta p/q} dy \right)^{1/p} \end{aligned}$$

when $(\beta/q - 1/n')p^* = (\beta p/q)p^*/p - p^*/p' - 1$ and $\beta p/q > p - 1$ since $\beta > q - 1$. In this case,

$$\left(\int_{\mathbf{B}} \left((1 - |x|)^{\beta/q-1/n'} F(x) \right)^{p^*} dx \right)^{1/p^*} \leq C \left(\int_{\mathbf{B}} \left((1 - |y|)^{\beta/q} |f(y)| \right)^p dy \right)^{1/p}.$$

Next note

$$\begin{aligned} b(x)F(x) &= \int_{B(0,|x|)} |f(y)|\{b(x) - b(y)\}dy + \int_{B(0,|x|)} |f(y)|b(y)dy \\ &\leq C \int_{B(0,|x|)} |f(y)|(1 - |y|)^\theta dy + \int_{B(0,|x|)} |f(y)|b(y)dy. \end{aligned}$$

Theorem 2.1 gives

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{\beta/q-1/n'} \int_{B(0,|x|)} |f(y)|(1 - |y|)^\theta dy \right)^{q^*} dx \right)^{1/q^*} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)|(1 - |y|)^\theta |p(1 - |y|)^{(\beta/q-\theta)p} dy \right)^{1/p} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{\beta p/q} dy \right)^{1/p} \end{aligned}$$

when $(\beta/q - 1/n')q^* = (\beta p/q - \theta p)q^*/p - q^*/p' - 1$ and $\beta p/q - \theta p > p - 1$ since $\beta > q - 1$. Moreover,

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{\beta/q - 1/n'} \int_{B(0,|x|)} |f(y)|b(y)dy \right)^{q^*} dx \right)^{1/q^*} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)b(y)|^q (1 - |y|)^\beta dy \right)^{1/q} \end{aligned}$$

since $(\beta/q - 1/n')q^* = \beta q^*/q - q^*/q' - 1$ and $\beta > q - 1$. Thus

$$\int_{\mathbf{B}} \left((1 - |x|)^{\beta/q - 1/n'} b(x)F(x) \right)^{q^*} dx \leq C$$

when $\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{\beta/q} |f(y)|) dy \leq 1$. This completes the proof. \square

In the same way as Proposition 2.2, we have the following result.

PROPOSITION 3.2. *Let $1 \leq p < q < n$, $\beta > q - 1$ and $1/q = 1/p - \theta > 0$. Let u be a function in $C^1(\mathbb{R}^n)$ such that*

$$\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{\beta/q} |\nabla u(y)|) dy < \infty.$$

Then

$$\begin{aligned} & \lim_{r \rightarrow 1} \left((1 - r)^{\beta/q - 1/p'} \frac{1}{|\mathbf{B} \setminus B(0, r)|} \int_{\mathbf{B} \setminus B(0, r)} |u(x)| dx \right. \\ & \left. + (1 - r)^{\beta/q - 1/q'} \frac{1}{|\mathbf{B} \setminus B(0, r)|} \int_{\mathbf{B} \setminus B(0, r)} b(x)|u(x)| dx \right) = 0 \end{aligned}$$

and

$$\begin{aligned} & \lim_{r \rightarrow 1} \left((1 - r)^{\beta/q - 1/p'} \frac{1}{|S(0, r)|} \int_{S(0, r)} |u(r\xi)| dS(\xi) \right. \\ & \left. + (1 - r)^{\beta/q - 1/q'} \frac{1}{|S(0, r)|} \int_{S(0, r)} b(r\xi)|u(r\xi)| dS(\xi) \right) = 0. \end{aligned}$$

For this, note that

$$(\beta/q - 1/n')p^* = (\beta p/q)p^*/p - p^*/p' - 1 \quad \text{and} \quad \beta p/q = (\beta/q - 1/p')p + p - 1$$

and

$$(\beta/q - 1/n')q^* = \beta q^*/q - q^*/q' - 1 \quad \text{and} \quad \beta = (\beta/q - 1/q')q + q - 1.$$

Next we consider the dual Hardy operator.

THEOREM 3.3. *Let $1 \leq p < q < n$, $\beta < p - 1$ and $1/q = 1/p - \theta > 0$. Set $G(x) = \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| dy$. Then there exists a constant $C > 0$ such that*

$$\int_{\mathbf{B}} \Phi^* \left(x, (1 - |x|)^{\beta/p-1/n'} G(x) \right) dx \leq C$$

when $\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{\beta/p} |f(y)|) dy \leq 1$.

Proof. First note from Theorem 2.4

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{\beta/p-1/n'} \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| dy \right)^{p^*} dx \right)^{1/p^*} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^\beta dy \right)^{1/p} \end{aligned}$$

when $(\beta/p - 1/n')p^* = \beta p^*/p - p^*/p' - 1$ and $\beta < p - 1$. In this case,

$$\left(\int_{\mathbf{B}} \left((1 - |x|)^{\beta/p-1/n'} G(x) \right)^{p^*} dx \right)^{1/p^*} \leq C \left(\int_{\mathbf{B}} \left((1 - |y|)^{\beta/p} |f(y)| \right)^p dy \right)^{1/p}.$$

Next note

$$\begin{aligned} b(x)G(x) &= \int_{\mathbf{B} \setminus B(0,|x|)} \{b(x) - b(y)\} |f(y)| dy + \int_{\mathbf{B} \setminus B(0,|x|)} b(y) |f(y)| dy \\ &\leq C(1 - |x|)^\theta \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| dy + \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| b(y) dy. \end{aligned}$$

Theorem 2.4 gives

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{\beta/p-1/n'+\theta} \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| dy \right)^{q^*} dx \right)^{1/q^*} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^\beta dy \right)^{1/p} \end{aligned}$$

when $(\beta/p - 1/n' + \theta)q^* = \beta q^*/p - q^*/p' - 1$ and $\beta < p - 1$. Moreover,

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{\beta/p-1/n'} \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| b(y) dy \right)^{q^*} dx \right)^{1/q^*} \\ & \leq C \left(\int_{\mathbf{B}} |f(y) b(y)|^q (1 - |y|)^{\beta q/p} dy \right)^{1/q} \end{aligned}$$

when $(\beta/p - 1/n')q^* = (\beta q/p)q^*/q - q^*/q' - 1$ and $\beta q/p < q - 1$ since $\beta < p - 1$. Thus

$$\int_{\mathbf{B}} \left((1 - |x|)^{\beta/p-1/n'} b(x)G(x) \right)^{q^*} dx \leq C$$

when $\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{\beta/p} |f(y)|) dy \leq 1$. The proof is now completed. \square

In the same way as Proposition 2.5, we have the following result.

PROPOSITION 3.4. *Let $1 \leq p < q < n$, $\beta < p - 1$ and $1/q = 1/p - \theta > 0$. Let u be a function in $C^1(\mathbb{R}^n)$ such that*

$$\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{\beta/p} |\nabla u(y)|) dy < \infty.$$

Then

$$\begin{aligned} & \lim_{r \rightarrow 1} \left((1 - r)^{-(\beta/q - 1/p')} \frac{1}{|\mathbf{B} \setminus B(0, r)|} \int_{\mathbf{B} \setminus B(0, r)} U_1(x) dx \right. \\ & \left. + (1 - r)^{-(\beta/q - 1/q')} \frac{1}{|\mathbf{B} \setminus B(0, r)|} \int_{\mathbf{B} \setminus B(0, r)} b(x) U_1(x) dx \right) = 0 \end{aligned}$$

and

$$\begin{aligned} & \lim_{r \rightarrow 1} \left((1 - r)^{-(\beta/q - 1/p')} \frac{1}{|S(0, r)|} \int_{S(0, r)} U_1(r\xi) dS(\xi) \right. \\ & \left. + (1 - r)^{-(\beta/q - 1/q')} \frac{1}{|S(0, r)|} \int_{S(0, r)} b(r\xi) U_1(r\xi) dS(\xi) \right) = 0. \end{aligned}$$

4. Sharpness

We discuss the sharpness of Theorem 3.1 in the double phase setting.

REMARK 4.1. In Theorem 3.1, the single condition that

$$\int_{\mathbf{B}} (|f(y)|b(y))^q (1 - |y|)^\beta dy \leq 1$$

may not imply

$$\int_{\mathbf{B}} \left(b(x) (1 - |x|)^{\beta/q - 1/n'} F(x) \right)^{q^*} dx \leq C.$$

In fact, for $0 < r < 1$ and $a > 0$ consider

$$b(x) = \begin{cases} |x|^\theta - r^\theta & \text{on } \mathbf{B} \setminus B(0, r); \\ 0 & \text{on } B(0, r); \end{cases}$$

and

$$f_a(y) = \begin{cases} 0 & \text{on } \mathbf{B} \setminus B(0, r); \\ a & \text{on } B(0, r). \end{cases}$$

Then note

$$(1) \int_{\mathbf{B}} (|f_a(y)|b(y))^q (1 - |y|)^\beta dy = 0 ;$$

- (2) $\int_{B(0,|x|)} |f_a(y)|dy \geq Cr^n a$ for $x \in \mathbf{B} \setminus B(0, r)$;
- (3) $\lim_{a \rightarrow \infty} \int_{\mathbf{B}} \left(b(x)(1 - |x|)^{\beta/q-1/n'} \int_{B(0,|x|)} |f_a(y)|dy \right)^{q^*} dx = \infty$.

5. The borderline case

The borderline case $\beta = p - 1$ in Theorem 2.1 is known.

THEOREM 5.1. ([26, Theorem 5.1]) *Let $1 \leq p \leq q$. Then there exists a constant $C > 0$ such that*

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left(\int_{B(0,|x|)} |f(y)|dy \right)^q (1 - |x|)^{-1} (\log(e/(1 - |x|)))^{-a} dx \right)^{1/q} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{p-1} (\log(e/(1 - |y|)))^{p/p'+(1-a)p/q} dy \right)^{1/p} \end{aligned}$$

when $a > 1$.

REMARK 5.2. Let p, q be as in Theorem 5.1. Suppose $a_1 > a > 1$ and $1 + (1 - a_1)/q < \gamma < 1 + (1 - a)/q$.

Consider

$$f(y) = |y|^{-1} (\log(e/(1 - |y|)))^{-\gamma} \text{ on } \mathbf{B}.$$

Then

- (1) $\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{p-1} (\log(e/(1 - |y|)))^{p/p'+(1-a_1)p/q} dy < \infty$;
- (2) $\int_{\mathbf{B}} \left(\int_{B(0,|x|)} |f(y)|dy \right)^q (1 - |x|)^{-1} (\log(e/(1 - |x|)))^{-a} dx = \infty$.

This implies that in Theorem 5.1, the exponent $p/p' + (1 - a)p/q$ could not be replaced by a smaller $p/p' + (1 - a_1)p/q$.

The borderline case $\beta = q - 1$ in Theorem 3.1 is treated in the following.

THEOREM 5.3. *Let $1 \leq p < q < n$ and $1/q = 1/p - \theta > 0$. Set*

$$F(x) = \int_{B(0,|x|)} |f(y)|dy$$

as before and

$$L(y) = 1 + (\log(e/(1 - |y|)))^{q/q'+(1-a)q/q^*} + (\log(e/(1 - |y|)))^{p/p'+(1-a)p/q^*}.$$

If $a > 1$, then there exists a positive constant $C > 0$ such that

$$\int_{\mathbf{B}} \Phi^* \left(x, (1 - |x|)^{-1/q+1/n} F(x) \right) (\log(e/(1 - |x|)))^{-a} dx \leq C$$

when $\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{(q-1)/q} |f(y)|) \tilde{L}(y) dy \leq 1$.

Proof. In view of Theorem 2.1 we have

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{-1/q+1/n} F(x) \right)^{p^*} dx \right)^{1/p^*} \\ &= \left(\int_{\mathbf{B}} F(x)^{p^*} (1 - |x|)^{(-1/q+1/n)p^*} dx \right)^{1/p^*} \\ &\leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{p(q-1)/q} dy \right)^{1/p} \\ &\leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{p(q-1)/q} L(y) dy \right)^{1/p} \end{aligned}$$

when $(-1/q + 1/n)p^* = (p(q - 1)/q)p^*/p - p^*/p' - 1$ and $p(q - 1)/q > p - 1$.

As in the proof of Theorem 3.1, note

$$b(x)F(x) \leq C \int_{B(0,|x|)} |f(y)|(1 - |y|)^\theta dy + \int_{B(0,|x|)} |f(y)|b(y)dy.$$

Theorem 5.1 gives

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{-1/q+1/n} \int_{B(0,|x|)} |f(y)|(1 - |y|)^\theta dy \right)^{q^*} (\log(e/(1 - |x|)))^{-a} dx \right)^{1/q^*} \\ &\leq C \left(\int_{\mathbf{B}} \left(|f(y)|(1 - |y|)^\theta \right)^p (1 - |y|)^{p-1} (\log(e/(1 - |y|)))^{p/p'+(1-a)p/q^*} dy \right)^{1/p} \\ &= C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{p-1+\theta p} (\log(e/(1 - |y|)))^{p/p'+(1-a)p/q^*} dy \right)^{1/p} \\ &\leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{p(q-1)/q} L(y) dy \right)^{1/p}. \end{aligned}$$

Moreover,

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{-1/q+1/n} \int_{B(0,|x|)} |f(y)|b(y)dy \right)^{q^*} (\log(e/(1 - |x|)))^{-a} dx \right)^{1/q^*} \\ &\leq C \left(\int_{\mathbf{B}} (|f(y)|b(y))^q (1 - |y|)^{q-1} (\log(e/(1 - |y|)))^{q/q'+(1-a)q/q^*} dy \right)^{1/q} \\ &\leq C \left(\int_{\mathbf{B}} (|f(y)|b(y))^q (1 - |y|)^{q-1} L(y) dy \right)^{1/q} \end{aligned}$$

since $a > 1$.

Hence we find

$$\int_{\mathbf{B}} \left((1 - |x|)^{-1/q+1/n} b(x)F(x) \right)^{q^*} (\log(e/(1 - |x|)))^{-a} dx \leq C$$

when $a > 1$ and $\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{(q-1)/q} |f(y)|) L(y) dy \leq 1$.

Thus we obtain the required inequality. \square

Finally we treat the dual Hardy operator. The borderline case $\beta = p - 1$ in Theorem 2.4 is known.

THEOREM 5.4. ([26, Theorem 5.5]) *Let $1 \leq p \leq q$ and $a < 1$. Then there exists a constant $C > 0$ such that*

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left(\int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| dy \right)^q (1 - |x|)^{-1} (\log(e/(1 - |x|)))^{-a} dx \right)^{1/q} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{p-1} (\log(e/(1 - |y|)))^{p/p'+(1-a)p/q} dy \right)^{1/p}. \end{aligned}$$

With the aid of Theorem 5.4, we obtain the following result as in Theorem 3.3.

THEOREM 5.5. *Let $1 \leq p < q < n$ and $1/q = 1/p - \theta > 0$. Set*

$$G(x) = \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| dy$$

as before. If $a < 1$, then

$$\int_{\mathbf{B}} \Phi^* \left(x, (1 - |x|)^{-1/p+1/n} G(x) \right) (\log(e/(1 - |x|)))^{-a} dx \leq C$$

when $\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{(p-1)/p} |f(y)|) (\log(e/(1 - |y|)))^{p/p'+(1-a)p/p^*} dy \leq 1$.

Proof. In view of Theorem 5.4 we have

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{-1/p+1/n} G(x) \right)^{p^*} (\log(e/(1 - |x|)))^{-a} dx \right)^{1/p^*} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{p-1} (\log(e/(1 - |y|)))^{p/p'+(1-a)p/p^*} dy \right)^{1/p}. \end{aligned}$$

As in the proof of Theorem 3.3 note

$$b(x)G(x) \leq C(1 - |x|)^\theta \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| dy + \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| b(y) dy.$$

By Theorem 5.4 we obtain

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{-1/p+1/n} (1 - |x|)^\theta \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| dy \right)^{q^*} (\log(e/(1 - |x|)))^{-a} dx \right)^{1/q^*} \\ & = \left(\int_{\mathbf{B}} \left((1 - |x|)^{-1/q+1/n} \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| dy \right)^{q^*} (\log(e/(1 - |x|)))^{-a} dx \right)^{1/q^*} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{p-1} (\log(e/(1 - |y|)))^{p/p'+(1-a)p/q^*} dy \right)^{1/p} \\ & \leq C \left(\int_{\mathbf{B}} |f(y)|^p (1 - |y|)^{p-1} (\log(e/(1 - |y|)))^{p/p'+(1-a)p/p^*} dy \right)^{1/p} \end{aligned}$$

when $a < 1$. Moreover, Theorem 2.4 gives

$$\begin{aligned} & \left(\int_{\mathbf{B}} \left((1 - |x|)^{-1/p+1/n} \int_{\mathbf{B} \setminus B(0,|x|)} |f(y)| b(y) dy \right)^{q^*} dx \right)^{1/q^*} \\ & \leq C \left(\int_{\mathbf{B}} (|f(y)| b(y))^q (1 - |y|)^{q(p-1)/p} dy \right)^{1/q} \end{aligned}$$

when $(-1/p + 1/n)q^* = (q(p-1)/p)q^*/q - q^*/q' - 1$ and $q(p-1)/p < q - 1$.

Hence we find

$$\int_{\mathbf{B}} ((1 - |x|)^{-1/p+1/n} b(x) G(x))^{q^*} (\log(e/(1 - |y|)))^{-a} dx \leq C$$

when $\int_{\mathbf{B}} \Phi(y, (1 - |y|)^{(p-1)/p} |f(y)|) (\log(e/(1 - |y|)))^{p/p'+(1-a)p/p^*} dy \leq 1$.

Thus Theorem 5.5 is proved. \square

REFERENCES

- [1] P. BARONI, M. COLOMBO AND G. MINGIONE, *Regularity for general functionals with double phase*, Calc. Var. Partial Differential Equations **57** (2018), no. 2, Paper No. 62, 48 pp.
- [2] P. BARONI, M. COLOMBO AND G. MINGIONE, *Non-autonomous functionals, borderline cases and related function classes*, St Petersburg Math. J. **27** (2016), 347–379.
- [3] S.-S. BYUN, J. OH, *Regularity results for generalized double phase functionals*, Anal. PDE **13** (2020), no. 5, 1269–1300.
- [4] S.-S. BYUN, S. RYU AND P. SHIN, *Calderon-Zygmund estimates for ω -minimizers of double phase variational problems*, Appl. Math. Letters **86** (2018), 256–263.
- [5] M. COLOMBO AND G. MINGIONE, *Regularity for double phase variational problems*, Arch. Ration. Mech. Anal. **215** (2015), no. 2, 443–496.
- [6] M. COLOMBO AND G. MINGIONE, *Bounded minimizers of double phase variational integrals*, Arch. Rat. Mech. Anal. **218** (2015), 219–273.
- [7] C. DE FILIPPIS AND G. MINGIONE, *Lipschitz bounds and nonautonomous Integrals*, Arch. Ration. Mech. Anal. **242** (2021), no. 2, 973–1057.
- [8] C. DE FILIPPIS AND J. OH, *Regularity for multi-phase variational problems*, J. Differential Equations **267** (2019), no. 3, 1631–1670.
- [9] C. DE FILIPPIS AND G. PALATUCCI, *Hölder regularity for nonlocal double phase equations*, J. Differential Equations **267** (2019), no. 1, 547–586.
- [10] G. H. HARDY, J. E. LITTLEWOOD AND G. POLYA, *Inequalities. 2d ed.* Cambridge, at the University Press, 1952. xii+324 pp.
- [11] P. HARJULEHTO AND P. HÄSTÖ, *Orlicz Spaces and Generalized Orlicz Spaces*, Lecture Notes in Mathematics, **2236**, Springer, Cham, 2019.
- [12] P. HARJULEHTO, P. HÄSTÖ AND M. LEE, *Hölder continuity of quasiminimizers and ω -minimizers of functionals with generalized Orlicz growth*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **XXII** (2021), 549–582.
- [13] P. HÄSTÖ, *The maximal operator on generalized Orlicz spaces*, J. Funct. Anal. **269** (2015), no. 12, 4038–4048; Corrigendum to “The maximal operator on generalized Orlicz spaces”, J. Funct. Anal. **271** (2016), no. 1, 240–243.
- [14] P. HÄSTÖ AND J. OK, *Maximal regularity for local minimizers of non-autonomous functionals*, to appear in J. Eur. Math. Soc.
- [15] A. KUFNER AND B. OPIC, *Hardy-type inequalities*, Pitman Research Notes in Mathematics Series, vol. 219, Longman Group UK Limited, London, 1990.
- [16] A. KUFNER, L.-E. PERSSON AND N. SAMKO, *Weighted inequalities of Hardy type*, second edition, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2017.
- [17] F.-Y. MAEDA, Y. MIZUTA, T. OHNO AND T. SHIMOMURA, *Sobolev’s inequality for double phase functionals with variable exponents*, Forum Math. **31** (2019), 517–527.
- [18] F.-Y. MAEDA, Y. MIZUTA AND T. SHIMOMURA, *Growth properties of Musielak-Orlicz integral means for Riesz potentials*, Nonlinear Anal. **112** (2015), 69–83.
- [19] P. MARCELLINI, *On the definition and the lower semicontinuity of certain quasiconvex integrals*, Ann. Inst. H. Poincaré Anal. Non Linéaire **3** (1986), no. 5, 391–409.
- [20] P. MARCELLINI, *Regularity of minimizers of integrals of the calculus of variations with nonstandard growth conditions*, Arch. Rational Mech. Anal. **105** (1989), no. 3, 267–284.
- [21] V. G. MAZ’YA, *Sobolev Spaces*, Springer, Heidelberg, 2011.
- [22] Y. MIZUTA, *Potential theory in Euclidean spaces*, Gakkōtoshō, Tokyo, 1996.

- [23] Y. MIZUTA, E. NAKAI, T. OHNO AND T. SHIMOMURA, *Campanato-Morrey spaces for the double phase functionals*, Rev. Mat. Complut. **33** (2020), 817–834.
- [24] Y. MIZUTA, T. OHNO AND T. SHIMOMURA, *Sobolev's theorem for double phase functionals*, Math. Ineq. Appl. **23** (2020), 17–33.
- [25] Y. MIZUTA AND T. SHIMOMURA, *Hardy-Sobolev inequalities in the half space for double phase functionals*, to appear in Rocky Mountain J. Math.
- [26] Y. MIZUTA AND T. SHIMOMURA, *Hardy-Sobolev inequalities in the unit ball for double phase functionals*, J. Math. Anal. Appl. **501** (2021), 124133, 17 pp.
- [27] Y. MIZUTA AND T. SHIMOMURA, *Boundary growth of Sobolev functions of monotone type for double phase functionals*, Ann. Fenn. Math. **47** (2022), 23–37.
- [28] L.-E. PERSSON AND S. G. SAMKO, *A note on the best constants in some Hardy inequalities*, J. Math. Inequal. **9** (2015), no. 2, 437–447.
- [29] M. A. RAGUSA AND A. TACHIKAWA, *Regularity for minimizers for functionals of double phase with variable exponents*, Adv. Nonlinear Anal. **9** (2020), no. 1, 710–728.
- [30] T. SHIMOMURA, *L^q -mean limits for Taylor's expansion of Riesz potentials of functions in Orlicz classes*, Hiroshima Math. J. **27** (1997), 159–175.
- [31] V. V. ZHIKOV, *Averaging of functionals of the calculus of variations and elasticity theory*, Izv. Akad. Nauk SSSR Ser. Mat. **50** (1986), 675–710.

(Received August 20, 2021)

Yoshihiro Mizuta
Graduate School of Advanced Science and Engineering
Hiroshima University
Higashi-Hiroshima 739-8521, Japan
e-mail: yomizuta@hiroshima-u.ac.jp

Tetsu Shimomura
Department of Mathematics
Graduate School of Humanities and Social Sciences
Hiroshima University
Higashi-Hiroshima 739-8524, Japan
e-mail: tshimo@hiroshima-u.ac.jp