

## REFINEMENTS OF TWO DETERMINANTAL INEQUALITIES FOR POSITIVE SEMIDEFINITE MATRICES

YAN HONG AND FENG QI\*

Dedicated to retired Professor Junesang Choi at Dongguk University in South Korea

(Communicated by I. Perić)

*Abstract.* Let  $A, B, C \in \mathbb{C}^{n \times n}$  be positive semidefinite matrices and let  $|A|, |B|, |C|$  be determinants of  $A, B, C \in \mathbb{C}^{n \times n}$  respectively. In this paper, the authors prove two determinantal inequalities

$$|A + B + C| + |C| \geq |A + C| + |B + C| + (2^n - 2)|AB|^{1/2} + 3(3^{n-1} - 2^n + 1)|ABC|^{1/3}$$

and

$$|A + B + C| + |A| + |B| + |C| \geq |A + B| + |A + C| + |B + C| + 3(3^{n-1} - 2^n + 1)|ABC|^{1/3}.$$

These two inequalities refine known ones.

### 1. Introduction

Let  $A \in \mathbb{C}^{n \times n}$  be a square matrix of order  $n$ . In this paper, we will denote the determinant of the matrix  $A$  by  $|A|$  and denote eigenvalues of the matrix  $A$  by

$$\lambda_1(A), \lambda_2(A), \dots, \lambda_n(A).$$

If eigenvalues of the matrix  $A$  are real numbers, we specify

$$\lambda_1(A) \geq \lambda_2(A) \geq \dots \geq \lambda_n(A).$$

Let  $A, B \in \mathbb{C}^{n \times n}$  be positive semidefinite matrices. In [1, p. 465, Corollary], Hertfiel obtained the determinantal inequality

$$|A + B| \geq |A| + |B| + (2^n - 2)(|A||B|)^{1/2}. \quad (1)$$

In [5, p. 215], Zhang established a determinantal inequality below.

*Mathematics subject classification* (2020): Primary 15A45; Secondary 15A39, 15A42.

*Keywords and phrases:* Determinantal inequality, eigenvalue, positive semidefinite matrix, determinant.

\* Corresponding author.

**THEOREM 1.** ([5, p. 215]) *Let  $A, B, C \in \mathbb{C}^{n \times n}$  be positive semidefinite matrices. Then*

$$|A + B + C| + |C| \geq |A + C| + |B + C|. \quad (2)$$

In [3], Lin established the following determinantal inequality.

**THEOREM 2.** ([3, Theorem 1.1]) *Let  $A, B, C \in \mathbb{C}^{n \times n}$  be positive semidefinite matrices. Then*

$$|A + B + C| + |A| + |B| + |C| \geq |A + B| + |A + C| + |B + C|. \quad (3)$$

In this paper, we will refine determinantal inequalities (2) and (3) above.

## 2. Lemmas

To refine determinantal inequalities (2) and (3) in Theorems 1 and 2, we need the following lemmas.

**LEMMA 1.** ([4, p. 333]) *Let  $A, B \in \mathbb{C}^{n \times n}$  be positive semidefinite matrices. Then*

$$\prod_{i=1}^n [\lambda_i(A) + \lambda_{n-i+1}(B)] \geq |A + B| \geq \prod_{i=1}^n [\lambda_i(A) + \lambda_i(B)].$$

By the proof of Theorem 1.1 in [3], we have the following inequality.

**LEMMA 2.** ([3, Theorem 1.1]) *Let  $A, B \in \mathbb{C}^{n \times n}$  be positive semidefinite matrices. Then*

$$\prod_{i=1}^n \lambda_i(A + B + I_n) - \prod_{i=1}^n \lambda_i(A + B) \geq \prod_{i=1}^n [\lambda_i(A) + \lambda_i(B) + 1] - \prod_{i=1}^n [\lambda_i(A) + \lambda_i(B)].$$

**LEMMA 3.** *Let  $a_i, b_i, c_i \geq 0$  for  $i \in \mathbb{N}$ . Then*

$$\begin{aligned} \prod_{i=1}^n (a_i + b_i + c_i) + \prod_{i=1}^n a_i + \prod_{i=1}^n b_i + \prod_{i=1}^n c_i - \prod_{i=1}^n (a_i + b_i) - \prod_{i=1}^n (a_i + c_i) - \prod_{i=1}^n (b_i + c_i) \\ \geq 3(3^{n-1} - 2^n + 1) \prod_{i=1}^n (a_i b_i c_i)^{1/3}. \end{aligned} \quad (4)$$

*Proof.* If  $n = 1, 2$ , the inequality (4) obviously holds. If  $n \geq 3$ , let  $i_1, \dots, i_k, j_1, \dots, j_t, \ell_1, \dots, \ell_s$  be non-negative positive numbers such that

$$\{i_1, \dots, i_k, j_1, \dots, j_t, \ell_1, \dots, \ell_s\} \setminus \{0\} = \{1, 2, \dots, n\}.$$

Since

$$\prod_{i=1}^n (a_i + b_i + c_i) = \sum_{\substack{0 \leq i_1 < \dots < i_k, 0 \leq j_1 < \dots < j_t, m=1 \\ 0 \leq \ell_1 < \dots < \ell_s, i_k + j_t + \ell_s = n}} \prod_{m=1}^k a_{i_m} \prod_{p=1}^t b_{j_p} \prod_{q=1}^s c_{\ell_q},$$

by virtue of the inequality between the geometric and arithmetic means, we acquire

$$\begin{aligned}
& \prod_{i=1}^n (a_i + b_i + c_i) + \prod_{i=1}^n a_i + \prod_{i=1}^n b_i + \prod_{i=1}^n c_i - \prod_{i=1}^n (a_i + b_i) - \prod_{i=1}^n (a_i + c_i) - \prod_{i=1}^n (b_i + c_i) \\
&= \sum_{\substack{1 \leq i_1 < \dots < i_k, 1 \leq j_1 < \dots < j_t, \\ 1 \leq \ell_1 < \dots < \ell_s, i_k + j_t + \ell_s = n}} \prod_{m=1}^k a_{i_m} \prod_{p=1}^t b_{j_p} \prod_{q=1}^s c_{\ell_q} \\
&\geq 3(3^{n-1} - 2^n + 1) \prod_{i=1}^n (a_i b_i c_i)^{\frac{3^{n-1} - 2^n + 1}{3(3^{n-1} - 2^n + 1)}} \\
&= 3(3^{n-1} - 2^n + 1) \prod_{i=1}^n (a_i b_i c_i)^{1/3}.
\end{aligned}$$

The proof of Lemma 3 is complete.  $\square$

REMARK 1. Taking  $c_i = 1$  for  $i = 1, 2, \dots, n$  in Lemma 3 results in

$$\begin{aligned}
& \prod_{i=1}^n (a_i + b_i + 1) + \prod_{i=1}^n a_i + \prod_{i=1}^n b_i + 1 - \prod_{i=1}^n (a_i + b_i) - \prod_{i=1}^n (a_i + 1) - \prod_{i=1}^n (b_i + 1) \\
&\geq 3(3^{n-1} - 2^n + 1) \prod_{i=1}^n (a_i b_i)^{1/3}, \quad (5)
\end{aligned}$$

where  $a_i, b_i \geq 0$  for  $i \in \mathbb{N}$ .

LEMMA 4. Let  $a_i, b_i, c_i \geq 0$  for  $i \in \mathbb{N}$ . Then

$$\begin{aligned}
& \prod_{i=1}^n (a_i + b_i + c_i) + \prod_{i=1}^n c_i - \prod_{i=1}^n (a_i + c_i) - \prod_{i=1}^n (b_i + c_i) \\
&\geq (2^n - 2) \prod_{i=1}^n (a_i b_i)^{1/2} + 3(3^{n-1} - 2^n + 1) \prod_{i=1}^n (a_i b_i c_i)^{1/3}. \quad (6)
\end{aligned}$$

*Proof.* Using the inequality (1), we obtain

$$\prod_{i=1}^n (a_i + b_i) - \prod_{i=1}^n a_i - \prod_{i=1}^n b_i \geq (2^n - 2) \prod_{i=1}^n (a_i b_i)^{1/2}.$$

Therefore, by the inequality (4), we arrive at

$$\begin{aligned}
& \prod_{i=1}^n (a_i + b_i + c_i) + \prod_{i=1}^n c_i - \prod_{i=1}^n (a_i + c_i) - \prod_{i=1}^n (b_i + c_i) \\
&\geq \prod_{i=1}^n (a_i + b_i) - \prod_{i=1}^n a_i - \prod_{i=1}^n b_i + 3(3^{n-1} - 2^n + 1) \prod_{i=1}^n (a_i b_i c_i)^{1/3} \\
&\geq (2^n - 2) \prod_{i=1}^n (a_i b_i)^{1/2} + 3(3^{n-1} - 2^n + 1) \prod_{i=1}^n (a_i b_i c_i)^{1/3}.
\end{aligned}$$

The proof of Lemma 4 is complete.  $\square$

REMARK 2. Setting  $c_i = 1$  for  $i = 1, 2, \dots, n$  in the inequality (6) in Lemma 4 leads to

$$\begin{aligned} & \prod_{i=1}^n (a_i + b_i + 1) + 1 - \prod_{i=1}^n (a_i + 1) - \prod_{i=1}^n (b_i + 1) \\ & \geq (2^n - 2) \prod_{i=1}^n (a_i b_i)^{1/2} + 3(3^{n-1} - 2^n + 1) \prod_{i=1}^n (a_i b_i)^{1/3}, \end{aligned} \quad (7)$$

where  $a_i, b_i \geq 0$  for  $i \in \mathbb{N}$ .

REMARK 3. From the inequality (7), it is easy to see that,

1. when  $\prod_{i=1}^n a_i b_i \geq 1$ , we have

$$\prod_{i=1}^n (a_i + b_i + 1) + 1 - \prod_{i=1}^n (a_i + 1) - \prod_{i=1}^n (b_i + 1) \geq (3^n - 2^{n+1} + 1) \prod_{i=1}^n (a_i b_i)^{1/3}.$$

2. when  $0 \leq \prod_{i=1}^n a_i b_i \leq 1$ , we have

$$\prod_{i=1}^n (a_i + b_i + 1) + 1 - \prod_{i=1}^n (a_i + 1) - \prod_{i=1}^n (b_i + 1) \geq (3^n - 2^{n+1} + 1) \prod_{i=1}^n (a_i b_i)^{1/2}.$$

### 3. Refinements of two determinantal inequalities

In this section, we refine determinantal inequalities (2) and (3).

**THEOREM 3.** Let  $A, B, C \in \mathbb{C}^{n \times n}$  be positive semidefinite matrices. Then

$$\begin{aligned} & |A + B + C| + |A| + |B| + |C| \\ & \geq |A + B| + |A + C| + |B + C| + 3(3^{n-1} - 2^n + 1)|ABC|^{1/3}. \end{aligned} \quad (8)$$

*Proof.* If  $|C| = 0$ , the inequality (8) evidently holds.

If  $|C| \neq 0$ , putting

$$A_1 = C^{-1/2} A C^{-1/2} \quad \text{and} \quad B_1 = C^{-1/2} B C^{-1/2}.$$

Using Lemma 2 and the inequality (5), we deduce

$$\begin{aligned}
& (|A + B + C| + |A| + |B| + |C| - |A + B| - |A + C| - |B + C|)|C|^{-1} \\
&= |A_1 + B_1 + I_n| + |A_1| + |B_1| + 1 - |A_1 + B_1| - |A_1 + I_n| - |B_1 + I_n| \\
&= \prod_{i=1}^n \lambda_i(A_1 + B_1 + I_n) + \prod_{i=1}^n \lambda_i(A_1) + \prod_{i=1}^n \lambda_i(B_1) + 1 \\
&\quad - \prod_{i=1}^n \lambda_i(A_1 + B_1) - \prod_{i=1}^n \lambda_i(A_1 + I_n) - \prod_{i=1}^n \lambda_i(B_1 + I_n) \\
&= \prod_{i=1}^n [\lambda_i(A_1 + B_1) + 1] + \prod_{i=1}^n \lambda_i(A_1) + \prod_{i=1}^n \lambda_i(B_1) + 1 \\
&\quad - \prod_{i=1}^n [\lambda_i(A_1 + B_1)] - \prod_{i=1}^n [\lambda_i(A_1) + 1] - \prod_{i=1}^n [\lambda_i(B_1) + 1] \\
&\geq \prod_{i=1}^n [\lambda_i(A_1) + \lambda_i(B_1) + 1] + \prod_{i=1}^n \lambda_i(A_1) + \prod_{i=1}^n \lambda_i(B_1) + 1 \\
&\quad - \prod_{i=1}^n [\lambda_i(A_1) + \lambda_i(B_1)] - \prod_{i=1}^n [\lambda_i(A_1) + 1] - \prod_{i=1}^n [\lambda_i(B_1) + 1] \\
&\geq 3(3^{n-1} - 2^n + 1)|ABC|^{1/3}|C|^{-1}.
\end{aligned}$$

The proof of Theorem 3 is complete.  $\square$

**THEOREM 4.** *Let  $A, B, C \in \mathbb{C}^{n \times n}$  be positive semidefinite matrices. Then*

$$|A + B + C| + |C| \geq |A + C| + |B + C| + (2^n - 2)|AB|^{1/2} + 3(3^{n-1} - 2^n + 1)|ABC|^{1/3}. \quad (9)$$

*Proof.* By the inequality (8) and the inequality (1), we obtain

$$\begin{aligned}
& |A + B + C| + |C| - |A + C| - |B + C| \\
&\geq |A + B| - |A| - |B| + 3(3^{n-1} - 2^n + 1)|ABC|^{1/3} \\
&\geq (2^n - 2)|AB|^{1/2} + 3(3^{n-1} - 2^n + 1)|ABC|^{1/3}.
\end{aligned}$$

The proof of Theorem 4 is complete.  $\square$

**REMARK 4.** Theorem 4 can be proved by Lemma 1 and the inequality (7), as done in the proof of Theorem 3.

**REMARK 5.** This paper is a corrected and revised version of the preprint [2].

*Acknowledgements.* The authors are thankful to anonymous referees for their careful corrections to and valuable comments on the original version of this paper.

*Funding.* This work was partially supported by the National Natural Science Foundation of China (Grant No. 11901322) and by the Foundation of the Research Program of Science and Technology at Universities of Inner Mongolia Autonomous Region (Grant No. NJZY20119), China.

## REFERENCES

- [1] D. J. HARTFIEL, *An extension of Haynsworth's determinant inequality*, Proc. Amer. Math. Soc. **41** (1973), no. 2, 463–465; available online at <https://doi.org/10.2307/2039114>.
- [2] Y. HONG AND F. QI, *Two determinantal inequalities for positive semidefinite matrices*, Research Square (2021), available online at <https://doi.org/10.21203/rs.3.rs-360043/v1>.
- [3] M. LIN, *A determinantal inequality for positive semidefinite matrices*, Electron. J. Linear Algebra **27** (2014), 821–826; available online at <https://doi.org/10.13001/1081-3810.2830>.
- [4] A. W. MARSHALL, I. OLKIN, AND B. C. ARNOLD, *Inequalities: Theory of Majorization and its Applications*, 2nd ed., Springer Verlag, New York/Dordrecht/Heidelberg/London, 2011; available online at <http://dx.doi.org/10.1007/978-0-387-68276-1>.
- [5] F. ZHANG, *Matrix Theory: Basic Results and Techniques*, 2nd ed., Universitext, Springer, New York, 2011; available online at <https://doi.org/10.1007/978-1-4614-1099-7>.

(Received December 31, 2021)

Yan Hong

College of Mathematics and Physics  
Inner Mongolia Minzu University  
Tongliao 028043, Inner Mongolia, China  
e-mail: nmhongyan@163.com  
<https://orcid.org/0000-0002-9916-9097>

Feng Qi

Institute of Mathematics  
Henan Polytechnic University  
Jiaozuo 454003, Henan, China  
and  
School of Mathematical Sciences  
Tiangong University  
Tianjin 300387, China  
e-mail: qifeng618@gmail.com  
qifeng618@hotmail.com  
qifeng618@qq.com  
<https://qifeng618.wordpress.com>  
<https://orcid.org/0000-0001-6239-2968>